

A



A-level

Physics data and formulae

For use in exams from the June 2017 Series onwards

[Turn over]

DATA - FUNDAMENTAL CONSTANTS AND VALUES

Quantity	Symbol	Value	Units
speed of light in vacuo	c	3.00×10^8	m s^{-1}
permeability of free space	μ_0	$4\pi \times 10^{-7}$	H m^{-1}
permittivity of free space	ϵ_0	8.85×10^{-12}	F m^{-1}
magnitude of the charge of electron	e	1.60×10^{-19}	C
the Planck constant	h	6.63×10^{-34}	J s
gravitational constant	G	6.67×10^{-11}	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	N_A	6.02×10^{23}	mol^{-1}
molar gas constant	R	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	k	1.38×10^{-23}	J K^{-1}
the Stefan constant	σ	5.67×10^{-8}	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	α	2.90×10^{-3}	m K
electron rest mass (equivalent to 5.5×10^{-4} u)	m_e	9.11×10^{-31}	kg

electron charge/mass ratio	$\frac{e}{m_e}$	1.76×10^{11}	C kg^{-1}
proton rest mass (equivalent to 1.00728 u)	m_p	$1.67(3) \times 10^{-27}$	kg
proton charge/mass ratio	$\frac{e}{m_p}$	9.58×10^7	C kg^{-1}
neutron rest mass (equivalent to 1.00867 u)	m_n	$1.67(5) \times 10^{-27}$	kg
gravitational field strength	g	9.81	N kg^{-1}
acceleration due to gravity	g	9.81	m s^{-2}
atomic mass unit (1u is equivalent to 931.5 MeV)	u	1.661×10^{-27}	kg

[Turn over]

ALGEBRAIC EQUATION

quadratic equation $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

ASTRONOMICAL DATA

Body	Mass/kg	Mean radius/m
Sun	1.99×10^{30}	6.96×10^8
Earth	5.97×10^{24}	6.37×10^6

GEOMETRICAL EQUATIONS

arc length $= r\theta$

circumference of circle $= 2\pi r$

area of circle $= \pi r^2$

curved surface area of cylinder $= 2\pi rh$

area of sphere $= 4\pi r^2$

volume of sphere $= \frac{4}{3} \pi r^3$

PARTICLE PHYSICS

Class	Name	Symbol	Rest energy/MeV
photon	photon	γ	0
lepton	neutrino	ν_e	0
		ν_μ	0
	electron	e^\pm	0.510999
	muon	μ^\pm	105.659
mesons	π meson	π^\pm	139.576
		π^0	134.972
	K meson	K^\pm	493.821
		K^0	497.762
baryons	proton	p	938.257
	neutron	n	939.551

[Turn over]

PROPERTIES OF QUARKS

antiquarks have opposite signs

Type	Charge	Baryon number	Strangeness
u	$+\frac{2}{3}e$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}e$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}e$	$+\frac{1}{3}$	-1

PROPERTIES OF LEPTONS

		Lepton number
Particles:	$e^-, \nu_e; \mu^-, \nu_\mu$	+ 1
Antiparticles:	$e^+, \bar{\nu}_e, \mu^+, \bar{\nu}_\mu$	- 1

PHOTONS AND ENERGY LEVELS

photon energy	$E = hf = \frac{hc}{\lambda}$
photoelectricity	$hf = \phi + E_{k(\max)}$
energy levels	$hf = E_1 - E_2$
de Broglie wavelength	$\lambda = \frac{h}{p} = \frac{h}{mv}$

[Turn over]

WAVES

wave speed $c = f\lambda$ period $f = \frac{1}{T}$

first harmonic $f = \frac{1}{2l} \sqrt{\frac{T}{\mu}}$

fringe spacing $w = \frac{\lambda D}{s}$ diffraction grating $d \sin \theta = n\lambda$

refractive index of a substance s , $n = \frac{c}{c_s}$

for two different substances of refractive indices n_1 and n_2 ,

law of refraction $n_1 \sin \theta_1 = n_2 \sin \theta_2$

critical angle $\sin \theta_c = \frac{n_2}{n_1}$ for $n_1 > n_2$

MECHANICS**moments**

$$\text{moment} = Fd$$

**velocity and
acceleration**

$$v = \frac{\Delta s}{\Delta t}$$

$$a = \frac{\Delta v}{\Delta t}$$

**equations of
motion**

$$v = u + at$$

$$s = \left(\frac{u + v}{2} \right) t$$

$$v^2 = u^2 + 2as$$

$$s = ut + \frac{at^2}{2}$$

force

$$F = ma$$

force

$$F = \frac{\Delta(mv)}{\Delta t}$$

impulse

$$F \Delta t = \Delta(mv)$$

**work, energy
and power**

$$W = F s \cos \theta$$

$$E_k = \frac{1}{2} m v^2$$

$$\Delta E_p = mg\Delta h$$

$$P = \frac{\Delta W}{\Delta t}, P = Fv$$

$$\text{efficiency} = \frac{\text{useful output power}}{\text{input power}}$$

[Turn over]

MATERIALS

density $\rho = \frac{m}{v}$

Hooke's law $F = k \Delta L$

Young modulus = $\frac{\text{tensile stress}}{\text{tensile strain}}$

tensile stress = $\frac{F}{A}$

tensile strain = $\frac{\Delta L}{L}$

energy stored $E = \frac{1}{2} F \Delta L$

ELECTRICITY

current and pd $I = \frac{\Delta Q}{\Delta t}$ $V = \frac{W}{Q}$ $R = \frac{V}{I}$

resistivity $\rho = \frac{RA}{L}$

resistors in series $R_T = R_1 + R_2 + R_3 + \dots$

resistors in parallel $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$

power $P = VI = I^2 R = \frac{V^2}{R}$

emf $\varepsilon = \frac{E}{Q}$ $\varepsilon = I(R + r)$

CIRCULAR MOTION

magnitude of
angular speed

$$\omega = \frac{v}{r}$$

$$\omega = 2\pi f$$

centripetal
acceleration

$$a = \frac{v^2}{r} = \omega^2 r$$

centripetal
force

$$F = \frac{mv^2}{r} = m\omega^2 r$$

SIMPLE HARMONIC MOTION

acceleration

$$a = -\omega^2 x$$

displacement

$$x = A \cos(\omega t)$$

speed

$$v = \pm \omega \sqrt{(A^2 - x^2)}$$

maximum speed

$$v_{\max} = \omega A$$

maximum acceleration

$$a_{\max} = \omega^2 A$$

for a mass-spring system

$$T = 2\pi \sqrt{\frac{m}{k}}$$

for a simple pendulum

$$T = 2\pi \sqrt{\frac{l}{g}}$$

[Turn over]

THERMAL PHYSICS

energy to change
temperature

$$Q = mc\Delta\theta$$

energy to change
state

$$Q = ml$$

gas law

$$pV = nRT$$

$$pV = NkT$$

kinetic theory
model

$$pV = \frac{1}{3}Nm(c_{\text{rms}})^2$$

kinetic energy of
gas molecule

$$\frac{1}{2}m(c_{\text{rms}})^2 = \frac{3}{2}kT = \frac{3RT}{2N_A}$$

GRAVITATIONAL FIELDS

force between two
masses

$$F = \frac{Gm_1m_2}{r^2}$$

gravitational field
strength

$$g = \frac{F}{m}$$

magnitude of
gravitational field
strength in a radial
field

$$g = \frac{GM}{r^2}$$

work done

$$\Delta W = m\Delta V$$

gravitational
potential

$$V = -\frac{GM}{r}$$

$$g = -\frac{\Delta V}{\Delta r}$$

ELECTRIC FIELDS AND CAPACITORS

force between two point charges

$$F = \frac{1}{4\pi\epsilon_0} \frac{Q_1Q_2}{r^2}$$

force on a charge

$$F = EQ$$

field strength for a uniform field

$$E = \frac{V}{d}$$

work done

$$\Delta W = Q\Delta V$$

field strength for a radial field

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

electric potential

$$V = \frac{1}{4\pi\epsilon_0} \frac{Q}{r}$$

field strength

$$E = \frac{\Delta V}{\Delta r}$$

capacitance

$$C = \frac{Q}{V}$$

$$C = \frac{A\epsilon_0\epsilon_r}{d}$$

capacitor energy stored

$$E = \frac{1}{2}QV = \frac{1}{2}CV^2 = \frac{1}{2} \frac{Q^2}{C}$$

[Turn over]

capacitor charging $Q = Q_0(1 - e^{-\frac{t}{RC}})$

decay of charge $Q = Q_0 e^{-\frac{t}{RC}}$

time constant RC

MAGNETIC FIELDS

force on a current $F = BIl$

force on a moving charge $F = BQv$

magnetic flux $\Phi = BA$

magnetic flux linkage $N\Phi = BAN \cos \theta$

magnitude of induced emf $\varepsilon = N \frac{\Delta\Phi}{\Delta t}$

$$N\Phi = BAN \cos \theta$$

emf induced in a rotating coil $\varepsilon = BAN\omega \sin \omega t$

alternating current $I_{\text{rms}} = \frac{I_0}{\sqrt{2}} \quad V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$

transformer equations $\frac{N_s}{N_p} = \frac{V_s}{V_p}$

$$\text{efficiency} = \frac{I_s V_s}{I_p V_p}$$

NUCLEAR PHYSICS

inverse square law
for γ radiation

$$I = \frac{k}{x^2}$$

radioactive decay

$$\frac{\Delta N}{\Delta t} = -\lambda N, N = N_0 e^{-\lambda t}$$

activity

$$A = \lambda N$$

half-life

$$T_{1/2} = \frac{\ln 2}{\lambda}$$

nuclear radius

$$R = R_0 A^{1/3}$$

energy-mass
equation

$$E = mc^2$$

[Turn over]

OPTIONS

ASTROPHYSICS

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ light year} = 9.46 \times 10^{15} \text{ m}$$

$$1 \text{ parsec} = 2.06 \times 10^5 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$\text{Hubble constant, } H = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$\text{telescope in normal adjustment} \quad M = \frac{f_o}{f_e}$$

$$\text{Rayleigh criterion} \quad \theta \approx \frac{\lambda}{D}$$

$$\text{magnitude equation} \quad m - M = 5 \log \frac{d}{10}$$

$$\text{Wien's law} \quad \lambda_{\text{max}} T = 2.9 \times 10^{-3} \text{ m K}$$

$$\text{Stefan's law} \quad P = \sigma AT^4$$

$$\text{Schwarzschild radius} \quad R_s \approx \frac{2GM}{c^2}$$

Doppler shift for $v \ll c$ $\frac{\Delta f}{f} = -\frac{\Delta \lambda}{\lambda} = \frac{v}{c}$

red shift $z = -\frac{v}{c}$

Hubble's law $v = Hd$

[Turn over]

MEDICAL PHYSICS

lens equations $P = \frac{1}{f}$

$$m = \frac{v}{u}$$

$$\frac{1}{f} = \frac{1}{u} + \frac{1}{v}$$

threshold of hearing $I_0 = 1.0 \times 10^{-12} \text{ W m}^{-2}$

intensity level $\textit{intensity level} = 10 \log \frac{I}{I_0}$

absorption $I = I_0 e^{-\mu x}$

$$\mu_m = \frac{\mu}{\rho}$$

ultrasound imaging $Z = p c$

$$\frac{I_r}{I_i} = \left(\frac{Z_2 - Z_1}{Z_2 + Z_1} \right)^2$$

half-lives $\frac{1}{T_E} = \frac{1}{T_B} + \frac{1}{T_P}$

ENGINEERING PHYSICS

moment of inertia $I = \Sigma mr^2$

angular kinetic energy $E_k = \frac{1}{2} I\omega^2$

equations of angular motion $\omega_2 = \omega_1 + \alpha t$

$$\omega_2^2 = \omega_1^2 + 2\alpha\theta$$

$$\theta = \omega_1 t + \frac{\alpha t^2}{2}$$

$$\theta = \frac{(\omega_1 + \omega_2) t}{2}$$

torque $T = I \alpha$

$$T = F r$$

angular momentum angular momentum = $I \omega$

angular impulse $T\Delta t = \Delta(I\omega)$

work done $W = T\theta$

power $P = T\omega$

thermodynamics $Q = \Delta U + W$

$$W = p\Delta V$$

adiabatic change $pV^\gamma = \text{constant}$

isothermal change $pV = \text{constant}$

[Turn over]

heat engines

$$\text{efficiency} = \frac{W}{Q_H} = \frac{Q_H - Q_C}{Q_H}$$

$$\text{maximum theoretical efficiency} = \frac{T_H - T_C}{T_H}$$

work done per cycle = area of loop

input power = calorific value × fuel flow rate

**indicated power = (area of $p - V$ loop)
 × (number of cycles per second)
 × (number of cylinders)**

output or brake power $P = T\omega$

friction power = indicated power – brake power

heat pumps and refrigerators

$$\text{refrigerator: } COP_{\text{ref}} = \frac{Q_C}{W} = \frac{Q_C}{Q_H - Q_C}$$

$$\text{heat pump: } COP_{\text{hp}} = \frac{Q_H}{W} = \frac{Q_H}{Q_H - Q_C}$$

TURNING POINTS IN PHYSICS**electrons in fields**

$$F = \frac{eV}{d}$$

$$F = Bev$$

$$r = \frac{mv}{Be}$$

$$\frac{1}{2}mv^2 = eV$$

**Millikan's
experiment**

$$\frac{QV}{d} = mg$$

$$F = 6\pi\eta rv$$

Maxwell's formula

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$$

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2meV}}$$

[Turn over]

special relativity

$$t = \frac{t_0}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$l = l_0 \sqrt{1 - \frac{v^2}{c^2}}$$

$$E = mc^2 = \frac{m_0 c^2}{\sqrt{1 - \frac{v^2}{c^2}}}$$

ELECTRONICSresonant
frequency

$$f_0 = \frac{1}{2\pi \sqrt{LC}}$$

Q-factor

$$Q = \frac{f_0}{f_B}$$

operational
amplifiers: open
loop

$$V_{\text{out}} = A_{\text{OL}} (V_+ - V_-)$$

inverting amplifier

$$\frac{V_{\text{out}}}{V_{\text{in}}} = - \frac{R_f}{R_{\text{in}}}$$

non-inverting amplifier $\frac{V_{\text{out}}}{V_{\text{in}}} = 1 + \frac{R_f}{R_1}$

summing amplifier $V_{\text{out}} = -R_f \left(\frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} + \dots \right)$

difference amplifier $V_{\text{out}} = (V_+ - V_-) \frac{R_f}{R_1}$

Bandwidth requirement:

for AM **bandwidth = $2f_M$**

for FM **bandwidth = $2(\Delta f + f_M)$**

END OF FORMULAE

There are no formulae printed on this page

Copyright Information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2016 AQA and its licensors. All rights reserved.

IB/M/Jun17/CD/7408/INS/E5