



Surname \_\_\_\_\_

Other Names \_\_\_\_\_

Centre Number \_\_\_\_\_

Candidate Number \_\_\_\_\_

Candidate Signature \_\_\_\_\_

**AS**

**CHEMISTRY**

**Paper 1: Inorganic and Physical Chemistry**

**7404/1**

**Friday 27 May 2016          Morning**

**Time allowed: 1 hour 30 minutes**

**For this paper you must have:**

- **the Periodic Table/Data Sheet, provided as an insert (enclosed)**
- **a ruler with millimetre measurement**
- **a calculator, which you are expected to use where appropriate.**

**At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.**

**[Turn over]**



## INSTRUCTIONS

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer ALL questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.



## **INFORMATION**

- **The maximum mark for this paper is 80.**
- **The Periodic Table/Data Sheet is provided as in insert.**

## **ADVICE**

- **You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.**

**DO NOT TURN OVER UNTIL TOLD TO DO SO**



**SECTION A**

Answer ALL questions in this section.

1 This question is about electron configuration.

**0 1 . 1** Give the full electron configuration of an Al atom and of a  $\text{Cr}^{3+}$  ion. [2 marks]

Al atom

---

---

$\text{Cr}^{3+}$  ion

---

---

**0 1 . 2** Deduce the formula of the ion that has a charge of  $2+$  with the same electron configuration as krypton. [1 mark]

---

---



**0 1 . 3** Deduce the formula of the compound that contains 2+ ions and 3- ions that both have the same electron configuration as argon.  
[1 mark]

---

---

[Turn over]



2 This question is about Period 3 of the Periodic Table.

**0 2** . **1** Deduce which of  $\text{Na}^+$  and  $\text{Mg}^{2+}$  is the smaller ion.  
Explain your answer. [2 marks]

Smaller ion \_\_\_\_\_

\_\_\_\_\_

Explanation \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



**0 2 . 2** Write an equation to represent the process that occurs when the first ionisation energy for sodium is measured. [1 mark]

---


---

**[Turn over]**



- 0 2 . 3 The first ionisation energies of some Period 3 elements are shown in Figure 1.

FIGURE 1





- 3 This question is about a white solid,  $\text{MHCO}_3$ , that dissolves in water and reacts with hydrochloric acid to give a salt.



A student was asked to design an experiment to determine a value for the  $M_r$  of  $\text{MHCO}_3$ . The student dissolved 1464 mg of  $\text{MHCO}_3$  in water and made the solution up to  $250 \text{ cm}^3$ .

$25.0 \text{ cm}^3$  samples of the solution were titrated with  $0.102 \text{ mol dm}^{-3}$  hydrochloric acid. The results are shown in Table 1.

TABLE 1

	Rough	1	2	3
Initial burette reading / $\text{cm}^3$	0.00	10.00	19.50	29.25
Final burette reading / $\text{cm}^3$	10.00	19.50	29.25	38.90
Titre / $\text{cm}^3$	10.00	9.50	9.75	9.65





- 03 . 2** Calculate the amount, in moles, of  $\text{MHCO}_3$  in  $250 \text{ cm}^3$  of the solution. Then calculate the experimental value for the  $M_r$  of  $\text{MHCO}_3$ . Give your answer to the appropriate number of significant figures. [3 marks]

---

---

---

---

---

---

---

---

---

---



**03 . 3** The student identified use of the burette as the largest source of uncertainty in the experiment.

Using the same apparatus, suggest how the procedure could be improved to reduce the percentage uncertainty in using the burette.

Justify your suggested improvement.  
[2 marks]

Suggestion \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Justification \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

[Turn over]







- 4 Table 2 shows some data about the elements bromine and magnesium.

TABLE 2

Element	Melting point / K	Boiling point / K
Bromine	266	332
Magnesium	923	1383

- 0 4 . 1 In terms of structure and bonding explain why the boiling point of bromine is different from that of magnesium. Suggest why magnesium is a liquid over a much greater temperature range compared to bromine. [5 marks]

---

---

---

---

---

---

---

---

---

---






- 5 Figure 2 represents two glass flasks, P and Q, connected via a tap.

Flask Q (volume =  $1.00 \times 10^3 \text{ cm}^3$ ) is filled with ammonia ( $\text{NH}_3$ ) at 102 kPa and 300 K. The tap is closed and there is a vacuum in flask P.  
(Gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

FIGURE 2



**0 5 . 1** Calculate the mass of ammonia in flask Q.  
Give your answer to the appropriate number  
of significant figures. [3 marks]

---

---

---

---

---

---

---

---

---

---

---

[Turn over]





6

**0 6 . 1** Explain how ions are accelerated, detected and have their abundance determined in a time of flight (TOF) mass spectrometer.  
**[3 marks]**

---

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



- 0 6** . **2** Calculate the mass, in kg, of a single  $^{52}\text{Cr}^+$  ion.  
Assume that the mass of a  $^{52}\text{Cr}^+$  ion is the same  
as that of a  $^{52}\text{Cr}$  atom.

(The Avogadro constant  $L = 6.022 \times 10^{23} \text{ mol}^{-1}$ )  
[1 mark]

---

---

---

- 0 6** . **3** In a TOF mass spectrometer the kinetic energy  
(KE) of a  $^{52}\text{Cr}^+$  ion was  $1.269 \times 10^{-13} \text{ J}$

Calculate the velocity of the ion using the  
equation.

$$\text{KE} = \frac{1}{2}mv^2$$

( $m$  = mass/kg and  $v$  = velocity/ $\text{ms}^{-1}$ )  
[2 marks]

---


---

---



- 
- 
- 0 6 . 4** Bromine has two isotopes,  $^{79}\text{Br}$  and  $^{81}\text{Br}$ , in approximately equal abundance. In a TOF mass spectrometer bromine forms ions with formula  $[\text{Br}_2]^+$

Sketch the pattern of peaks you would expect to see in the mass spectrum of a sample of bromine. [2 marks]



[Turn over]



- 06** . **5** A sample of xenon has  $A_r = 131.31$ . The sample consists of four isotopes. The abundances of three of the isotopes are shown in Table 3. The data for one of the isotopes,  ${}^m\text{Xe}$ , is missing.

TABLE 3

Isotope	${}^{129}\text{Xe}$	${}^{131}\text{Xe}$	${}^{132}\text{Xe}$	${}^m\text{Xe}$
% abundance	28.0	25.0	27.0	To be calculated

Use the data to calculate the abundance of isotope  ${}^m\text{Xe}$  and calculate  $m$ , the mass number of  ${}^m\text{Xe}$ . Show your working. [4 marks]

---

---

---

---

---

---

---

---

---

---



---

---

---

---

---

---

---

---


---

---

**[Turn over]**



- 7 Ammonia reacts with aluminium chloride as shown by the equation:



- 0 7** . **1** Draw diagrams to illustrate the shapes of  $\text{NH}_3$  molecules and of  $\text{AlCl}_3$  molecules.

Include in your diagrams any lone pairs of electrons that influence the shape.

Indicate the values of the bond angles.  
[3 marks]



- 07** . **2** Name the type of bond formed between N and Al in  $\text{H}_3\text{NAlCl}_3$  and explain how this bond is formed. [2 marks]

Type of bond \_\_\_\_\_

\_\_\_\_\_

Explanation \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

[Turn over]



- 07 . 3** Explain how the value of the Cl-Al-Cl bond angle in  $\text{AlCl}_3$  changes, if at all, on formation of the compound  $\text{H}_3\text{NAlCl}_3$  [2 marks]

---

---

---

---

---

---

---

---

- 8** A student oxidised a solution of hydrochloric acid with a few drops of sodium chlorate(I) solution. The reaction mixture effervesced and turned pale green. The gas formed bleached universal indicator paper.



**0 8** . **1** Write a half-equation for the oxidation of chloride ions. [ 1mark]

---

**0 8** . **2** Write a half-equation for the reduction of chlorate(I) ions to chlorine in acidic conditions. [1 mark]

---

**0 8** . **3** Write an overall equation for the redox reaction of chlorate(I) ions with hydrochloric acid. [1 mark]

---

[Turn over]



**0 8 . 4** A solution of sodium chlorate(I) was added to a colourless solution of potassium iodide. Suggest what is observed.

**Explain the reaction that leads to this observation. [3 marks]**

---

---

---

---

---

---

---

---

---

---

---



9

**0 9 . 1** A student was given a powder made from a mixture of anhydrous barium chloride and anhydrous magnesium chloride. The student dissolved 1.056 g of the powder in water in a conical flask and added an excess of sulfuric acid.

A white precipitate formed and was filtered off, washed and dried.

The mass of this solid was 0.764 g.

Identify the white precipitate and calculate the percentage, by mass, of magnesium chloride in the powder. [4 marks]

---

---

---

---

---

---

---

---

---

---

**[Turn over]**






**SECTION B**




Answer ALL questions in the spaces provided

Only ONE answer per question is allowed.  
For each answer completely fill in the circle alongside  
the appropriate answer.


**CORRECT METHOD**




**WRONG METHODS**



If you want to change your answer you must  
cross out your original answer as shown.



If you wish to return to an answer previously  
crossed out, ring the answer you now wish to  
select as shown.



You may do your working out in the blank spaces  
around the questions but this will not be marked. Do  
NOT use additional sheets for this working.

[Turn over]



**1 0** Which element is in the d-block of the Periodic Table? [1 mark]

**A** Selenium

**B** Antimony

**C** Tantalum

**D** Lead

**1 1** Which species contains an element with an oxidation state of +4? [1 mark]

**A**  $\text{NO}_2^+$

**B**  $\text{ClO}_3^-$

**C**  $\text{H}_2\text{SO}_3$

**D**  $\text{PCl}_5$



- 1 2** There are 392 mol of pure gold in a bar measuring 10 cm by 10 cm by 40 cm.  
What is the density of gold in  $\text{kg dm}^{-3}$ ? [1 mark]

**A 193**

**B 19.3**

**C 1.93**

**D 0.193**

**[Turn over]**



**1 3** Ions of two isotopes of iron are



Which statement is correct? [1 mark]

**A** The ions of both the isotopes have the electronic configuration  $1s^22s^22p^63s^23p^64s^23d^6$

**B** The ions of both the isotopes contains 26 neutrons

**C**  $^{53}\text{Fe}^{2+}$  has fewer protons than  $^{56}\text{Fe}^{2+}$

**D** After acceleration to the same kinetic energy  $^{56}\text{Fe}^{2+}$  will move more slowly than  $^{53}\text{Fe}^{2+}$



**BLANK PAGE**


**[Turn over]**



1 4

The successive ionisation energies for element X are shown in Figure 3.

Ionisation energy/  
 $\text{kJ mol}^{-1}$



39

Which element in X? [1 mark]

A Nitrogen

B Phosphorous

C Aluminium

D Boron

1 5 Which of these decreases down Group 2?  
[1 mark]

A First ionisation energy

B Atomic radius


C Number of protons

D Reactivity with water

[Turn over]



Refer to the unbalanced equation below when answering questions 16 and 17.



16 In the balanced equation the mole ratio for sulfuric acid to water is [1 mark]

A 1 : 4

B 1 : 2

C 4 : 7

D 4 : 9



**1 7** What is the reducing agent in this reaction?  
[1 mark]

**A**  $\text{H}^+$

**B**  $\text{C}_2\text{O}_4^{2-}$

**C**  $\text{K}^+$

**D**  $\text{Cr}_2\text{O}_7^{2-}$

**1 8** Which substance exists as a macromolecule?  
[1 mark]

**A** **Cu**

**B**  $\text{SiO}_2$


**C**  $\text{P}_4\text{O}_{10}$

**D** **MgO**

[Turn over]



- 1 9** A pale brown mixture of  $\text{NO}_2$  and  $\text{N}_2\text{O}_4$  is allowed to reach equilibrium in a sealed gas syringe according to the following equation.



When the plunger is pushed further into the syringe the pressure increases and the mixture becomes paler in colour.

When the syringe is placed in a hot oven the mixture becomes darker in colour.

Which of the following statements is correct?  
[1 mark]

**A**  $\text{NO}_2$  is brown and the forward reaction is exothermic


**B**  $\text{NO}_2$  is brown and the forward reaction is endothermic.

**C**  $\text{NO}_2$  is colourless and the forward reaction is exothermic.


**D**  $\text{NO}_2$  is colourless and the forward reaction is endothermic.




**2 0** Which molecule has the largest dipole?  
[1 mark]




**A**  $\text{ClF}_3$



**B**  $\text{BF}_3$



**C**  $\text{SF}_6$



**D**  $\text{CF}_4$

[Turn over]



**2 1** In a molecule of a hydrocarbon, the fraction by mass of carbon is  $\frac{9}{11}$

What is the empirical formula of the hydrocarbon? [1 mark]

A CH


B CH<sub>3</sub>

C C<sub>3</sub>H<sub>8</sub>

D C<sub>5</sub>H<sub>12</sub>



- 2 2** 30 cm<sup>3</sup> of xenon are mixed with 20 cm<sup>3</sup> of fluorine. The gases react according to the following equation. Assume that the temperature and pressure remain constant.



What is the final volume of gas after the reaction is complete?

**A** 50 cm<sup>3</sup>

**B** 40 cm<sup>3</sup>

**C** 30 cm<sup>3</sup>

**D** 20 cm<sup>3</sup>

[Turn over]



**2 3** Which of the following solutions would react exactly with a solution containing 0.0500 mol sulfuric acid? [1 mark]

**A** 50.0 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> KOH

**B** 100.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> KOH

**C** 100.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> Ba(OH)<sub>2</sub>


**D** 50.0 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> Ba(OH)<sub>2</sub>



- 2 4** In a car airbag, sodium azide ( $\text{NaN}_3$ ) decomposes to form sodium metal and nitrogen gas.



The sodium metal then reacts with potassium nitrate to produce more nitrogen gas.



If 2.00 mol of sodium azide react in this way, how many molecules of  $\text{N}_2$  will be formed?

(The Avogadro constant  $L = 6.022 \times 10^{23} \text{ mol}^{-1}$ )  
[1 mark]

**A**  $2.41 \times 10^{24}$

**B**  $1.93 \times 10^{24}$

**C**  $1.81 \times 10^{24}$

**D**  $9.63 \times 10^{23}$

**END OF QUESTIONS**



**There are no questions printed on this page**

**Copyright Information**

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk) after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2016 AQA and its licensors. All rights reserved.

**IB/M/Jun16/CD/7404/1/E5**

