



Surname _____

Other Names _____

Centre Number _____

Candidate Number _____

Candidate Signature _____

AS

FURTHER MATHEMATICS

Paper 2 – Mechanics

7366/2M

Thursday 17 May 2018 Afternoon

Time allowed: 1 hour 30 minutes

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.

[Turn over]



For this paper:

- **You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.**
- **You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)**
- **You must ensure you have the other optional Question Paper/Answer Book for which you are entered (either Discrete or Statistics). You will have 1 hour 30 minutes to complete both papers.**

INSTRUCTIONS

- **Use black ink or black ball-point pen. Pencil should only be used for drawing.**
- **Answer ALL questions.**
- **You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).**



- **Show all necessary working; otherwise marks for method may be lost.**
- **Do all rough work in this book. Cross through any work that you do not want to be marked.**
- **Do not write on blank pages.**

INFORMATION

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 40.**

ADVICE

- **Unless stated otherwise, you may quote formulae, without proof, from the booklet.**
- **You do not necessarily need to use all the space provided.**

**DO NOT TURN OVER UNTIL TOLD
TO DO SO**



4

Answer ALL questions in the spaces provided.

1 A particle A , of mass 0.2 kg, collides with a particle B , of mass 0.3 kg

Immediately before the collision, the

velocity of A is $\begin{bmatrix} 4 \\ 12 \end{bmatrix} \text{ m s}^{-1}$

and the velocity of B is $\begin{bmatrix} -1 \\ -3 \end{bmatrix} \text{ m s}^{-1}$

As a result of the collision the particles coalesce to become a single particle.

Find the velocity of the single particle.



5

Circle your answer. [1 mark]

$$\begin{bmatrix} 0.5 \\ 1.5 \end{bmatrix} \text{ m s}^{-1} \qquad \begin{bmatrix} 2 \\ 6 \end{bmatrix} \text{ m s}^{-1}$$

$$\begin{bmatrix} 1 \\ 3 \end{bmatrix} \text{ m s}^{-1} \qquad \begin{bmatrix} 3 \\ 9 \end{bmatrix} \text{ m s}^{-1}$$

- 2 A train is travelling at maximum speed with its engine using its maximum power of 1800 kW
- When travelling at this speed the train experiences a total resistive force of 40 000 N
- Find the maximum speed of the train.
- Circle your answer. [1 mark]

$$22 \text{ m s}^{-1} \quad 45 \text{ m s}^{-1} \quad 54 \text{ m s}^{-1} \quad 90 \text{ m s}^{-1}$$

[Turn over]



6

- 3 The kinetic energy, E , of a compound pendulum is given by

$$E = \frac{1}{2} I \omega^2$$

where ω is the angular speed and I is a quantity called the moment of inertia.

- 3 (a) Show that for this formula to be dimensionally consistent then I must have dimensions ML^2 , where M represents mass and L represents length. [2 marks]



7

- 3(b) The time, T , taken for one complete swing of a pendulum is thought to depend on its moment of inertia, I , its weight, W , and the distance, h , of the centre of mass of the pendulum from the point of suspension.

The formula being proposed is

$$T = kI^\alpha W^\beta h^\gamma$$

where k is a dimensionless constant.

Determine the values of α , β and γ . [3 marks]

[Turn over]



8

4 Two smooth spheres A and B of equal radius are free to move on a smooth horizontal surface.

The masses of A and B are m and $4m$ respectively.

The coefficient of restitution between the spheres is e .

The spheres are projected directly towards each other, each with speed u , and subsequently collide.

4 (a) Show that the speed of B immediately after the impact with A is

$$\frac{u(3 - 2e)}{5}$$

[4 marks]



10

4(b) Find the speed of A in terms of u and e . [2 marks]



4 (c) Comment on the direction of motion of the spheres after the collision, justifying your answer. [2 marks]

[Turn over]



14

5 A car travels around a roundabout at a constant speed. The surface of the roundabout is horizontal.

The car has mass 990 kg and the path of the car is a circular arc of radius 48 metres.

A simple model assumes that the car is a particle and the only horizontal force acting on it as it travels around the roundabout is friction.

On a dry day typical values of friction, F , between the surface of the roundabout and the tyres of the car are

$$7300 \text{ N} \leq F \leq 9200 \text{ N}$$

5 (a) Using this model calculate a safe speed limit, in miles per hour, for the car as it travels around the roundabout.

Explain your reasoning fully.



BLANK PAGE



17

5(b) Gary assumes that on a wet day typical values for friction, F , are

$$5400 \text{ N} \leq F \leq 10\,000 \text{ N}$$

Comment on the validity of Gary's revised assumption. [2 marks]

[Turn over]



18

6 At a fairground a dodgem car is moving in a straight horizontal line towards a side wall that is perpendicular to the velocity of the car.

The speed of the car is 1.8 m s^{-1}

It collides with the side wall and rebounds along its original path with a speed of 1.2 m s^{-1}

The total mass of the dodgem car and the passengers is 250 kg

6 (a) Find the magnitude of the impulse on the car during the collision with the side wall. [2 marks]



BLANK PAGE



7 Use g as 9.8 m s^{-2} in this question.

Dominic, a bungee jumper of mass 75 kg, has his ankles attached to one end of a cord. The other end of the cord is attached to a bridge which is 50 metres above the surface of a river.

The cord can be modelled as a light elastic cord of natural length 25 metres and modulus of elasticity 3200 N. Dominic is modelled as a particle.

Dominic steps off the bridge at the point where the cord is attached and falls vertically downwards.

7 (a) Find Dominic's speed at the point when the cord initially becomes taut. [2 marks]



30

**Question
number**

**Additional page, if required.
Write the question numbers in
the left-hand margin.**

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
TOTAL	

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

PB/Jun18/7366/2M/E1



3 0