

1. March/2022/Paper_9709/62/No.6

In a game a ball is rolled down a slope and along a track until it stops. The distance, in metres, travelled by the ball is modelled by the random variable X with probability density function

$$f(x) = \begin{cases} -k(x-1)(x-3) & 1 \leq x \leq 3, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (a) Without calculation, explain why $E(X) = 2$.

[1]

Quadratic with roots $x=1$ and $x=3$ so symmetrical about $x = \frac{1+3}{2} = 2$.

- (b) Show that $k = \frac{3}{4}$.

[3]

$$\Rightarrow \int_1^3 -k(x-1)(x-3) dx = 1.0 \quad \text{Since } f(x) \text{ is a p.d.f}$$

$$\Rightarrow -k \int_1^3 (x^2 - 4x + 3) dx = 1$$

$$\Rightarrow -k \left[\frac{x^3}{3} - 2x^2 + 3x \right]_1^3 = 1$$

$$\Rightarrow \left[\frac{x^3}{3} - 2x^2 + 3x \right]_1^3 = \frac{1}{-k}$$

$$\left[\frac{(3)^3}{3} - 2(3)^2 + 3(3) \right] - \left[\frac{(1)^3}{3} - 2(1)^2 + 3(1) \right] = \frac{1}{-k}$$

$$\left[\frac{1}{3} - 2 + 3 \right] = \frac{1}{-k}$$

$$\left[9 - 18 + 9 \right] - \left[\frac{1}{3} - 2 + 3 \right] = \frac{1}{-k}$$

$$(-9 + 9) - \left(\frac{4}{3} \right) = \frac{1}{-k}$$

$$\frac{-4}{3} = \frac{1}{-k}$$

$$\frac{-4k}{-4} = \frac{-3}{-4}$$

$$k = \frac{3}{4}$$

Shown.

(c) Find $\text{Var}(X)$.


$$\begin{aligned} \text{Var}(X) &= \frac{-3}{4} \int_1^3 x^2 \cdot f(x) dx - [E(X)]^2 = \frac{-3}{4} \left[\frac{-27}{5} - \frac{1}{5} \right] - 4 \\ &= \frac{-3}{4} \int_1^3 (x^4 - 4x^3 + 3x^2) dx - [E(X)]^2 = \frac{-3}{4} \left(-\frac{28}{5} \right) - 4 \\ \text{Recall } \text{Var}(X) &= E(X^2) - [E(X)]^2 = 4 \cdot 2 - 4 \\ &= \frac{-3}{4} \left\{ \left[\frac{x^5}{5} - x^4 - x^3 \right]_1^3 \right\} - 2^2 \\ &= \frac{-3}{4} \left(\frac{243}{5} - 81 + 27 \right) - \left(\frac{1}{5} - 1 + 1 \right) \\ &= -4 \end{aligned}$$

One turn consists of rolling the ball 3 times and noting the largest value of X obtained. If this largest value is greater than 2.5, the player scores a point.

(d) Find the probability that on a particular turn the player scores a point.

$$\begin{aligned} &= \frac{-3}{4} \int_{2.5}^3 (x^2 - 4x + 3) dx \\ &= \frac{-3}{4} \left[\frac{x^3}{3} - 2x^2 + 3x \right]_{2.5}^3 \\ &= \frac{-3}{4} \left[\left(9 - 18 + 9 \right) - \left(\frac{125}{24} - \frac{25}{2} + 7.5 \right) \right] \\ &= \frac{-3}{4} \left(-\frac{5}{24} \right) \\ &= \frac{5}{32} \end{aligned}$$

$$P(X > 2.5) = P(A)$$



$$\begin{aligned} P(\text{score a point}) &= 1 - \left(1 - \frac{5}{32} \right)^3 \\ &= 1 - 0.60068 \\ &= 0.399 \\ &= 0.399 \end{aligned}$$