

Cambridge IGCSE™

CHEMISTRY**0620/41**

Paper 4 Theory (Extended)

May/June 2025

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of **14** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.

2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.

3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).

4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 'List rule' guidance

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards *n*.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.












Annotations guidance for centres


Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
	information missing or insufficient for credit
	benefit of doubt given
	contradiction in response, mark not awarded
	incorrect point or mark not awarded
	key point attempted / working towards marking point / incomplete answer / response seen but not credited / blank page seen
	error carried forward applied
	incorrect or insufficient response, mark not awarded
	response has not answered question
	benefit of doubt was considered, but the response was decided to not be sufficiently close for benefit of doubt to be applied
	incorrect point or mark not awarded
	blank page seen

Annotation	Meaning
	correct point of mark awarded

Question	Answer	Marks
1(a)	calcium oxide	1
1(b)	graphite	1
1(c)	ethanol	1
1(d)	methane	1
1(e)	bauxite	1
1(f)	sulfur dioxide	1
1(g)	propane	1
1(h)	calcium oxide	1
1(i)	oxygen	1
1(j)	propene	1

Question	Answer	Marks								
2(a)	<table border="1" style="display: inline-table; vertical-align: top;"> <thead> <tr> <th>M1 charge (1)</th> <th>M2 mass(1)</th> </tr> </thead> <tbody> <tr> <td>-1</td> <td></td> </tr> <tr> <td></td> <td>1</td> </tr> <tr> <td>+1</td> <td>1</td> </tr> </tbody> </table> <p>1 mark for each correct column</p>	M1 charge (1)	M2 mass(1)	-1			1	+1	1	2
M1 charge (1)	M2 mass(1)									
-1										
	1									
+1	1									

Question	Answer	Marks									
2(b)(i)	<table border="1"> <thead> <tr> <th>electron</th> <th>neutron</th> <th>proton</th> </tr> </thead> <tbody> <tr> <td>19</td> <td>20</td> <td>19</td> </tr> <tr> <td>18</td> <td>22</td> <td>19</td> </tr> </tbody> </table> <p>1 mark for each correct column</p>	electron	neutron	proton	19	20	19	18	22	19	3
electron	neutron	proton									
19	20	19									
18	22	19									
2(b)(ii)	<p>M1 $(41 \times 10) + (39 \times 90) = 3920$ (1)</p> <p>M2 $(3920 \div 100) = 39.2$(1)</p>	2									
2(b)(iii)	aluminium has only 1 isotope which has nucleon number 27	1									
2(c)	<p>M1 Ar (1)</p> <p>M2 P^{3-} or S^{2-} or Cl^{-} (1)</p>	2									

Question	Answer	Marks
3(a)(i)	<p>M1 positive ions / cations(1)</p> <p>M2 sea of electrons / mobile electrons / delocalised electrons(1)</p> <p>M3 attraction between positive ions and electrons(1)</p>	3
3(a)(ii)	<p>electrons move / electrons mobile / electrons flow</p>	1
3(b)(i)	harder or stronger	1
3(b)(ii)	zinc	1

Question	Answer	Marks
3(c)(i)	Any two from:- <ul style="list-style-type: none"> • solid disappears / solid dissolves • fizzing / bubbling / effervescence • blue solution 	2
3(c)(ii)	all the sulfuric acid had reacted	1
3(c)(iii)	copper(II) carbonate	1
3(c)(iv)	copper(II) oxide / copper(II) hydroxide	1
3(c)(v)	saturated (solution)	1
3(c)(vi)	M1 more particles per unit volume M2 more collisions per unit time / increased collision frequency	2
3(c)(vii)	hydrated	1

Question	Answer	Marks
4(a)	M1 3 dots and 3 crosses shared M2 one non-bonding pair of 2 dots on one N atom and 2 crosses on the other N atom to complete the octet on both	2

Question	Answer	Marks
4(b)(i)	<p>Conditions can be in any order</p> <p>M1 (temperature) 450 and °C</p> <p>M2(pressure) 200 and atmospheres / 20 000 and kPa</p> <p>M3 iron and catalyst</p>	3
4(b)(ii)	$\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$	1
4(c)(i)	$5\text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$	1
4(c)(ii)	<p>M1 –3</p> <p>M2 +2</p>	2
4(c)(iii)	increase in oxidation number	1
4(c)(iv)	<p>$4\text{NO} + 3\text{O}_2 + 2\text{H}_2\text{O} \rightarrow 4\text{HNO}_3$</p> <p>M1 $\text{HNO}_3(1)$</p> <p>M2 equation completely correct</p>	2
4(d)	<p>M1 15</p> <p>M2 7.5</p> <p>M3 990</p>	3

Question	Answer	Marks		
5(a)	M1 concentration is no longer changing M2 rates of forward reaction and of the reverse reaction are equal	2		
5(b)(i)	M1 rate decreases M2 yield of methanol decreases	2		
5(b)(ii)	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 2px;">no change(1)</td> <td style="width: 50%; padding: 2px;">increases (1)</td> </tr> </table>	no change(1)	increases (1)	2
no change(1)	increases (1)			
5(b)(iii)	exothermic	1		
5(b)(iv)	M1 copper M2 transition element	2		
5(c)(i)	any 2 from: <ul style="list-style-type: none"> • same or similar chemical properties or reactions • (same) general formula • (consecutive members) differ by CH₂ • same functional group • physical properties OR example of a physical property e.g. melting point / boiling point / volatility vary in predictable manner or show trends or gradually change 	2		
5(c)(ii)	C ₅ H ₁₂ O	1		

Question	Answer	Marks
5(d)(i)	<p>M1 displayed formula of ester linkage, i.e.</p> $\text{H or C} - \overset{\text{O}}{\parallel}{\text{C}} - \text{O} - \text{C}$ <p>M2 completely correct displayed formula of methyl ethanoate or ethyl methanoate i.e.</p> $\begin{array}{c} \text{H} & \text{O} & \text{H} \\ & & \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{C}-\text{H} \\ & & \\ \text{H} & & \text{H} \end{array}$ <p>OR</p> $\begin{array}{c} \text{O} & \text{H} & \text{H} \\ & & \\ \text{H}-\text{C}-\text{O}-\text{C}-\text{C}-\text{H} \\ & & \\ & \text{H} & \text{H} \end{array}$	2
5(d)(ii)	<p>M1 butan-1-ol</p> <p>M2 ethanoic acid</p>	2

Question	Answer	Marks
5(e)	M1 C 64.87 / 12 H 13.51 / 1 O 21.62 / 16 OR 5.41 13.51 1.35 M2 C ₄ H ₁₀ O	2

Question	Answer	Marks
6(a)	halogen(s)	1
6(b)	fluorine / F ₂	1
6(c)	M1 solid M2 grey-black	2
6(d)(i)	$\text{Br}_2 + 2\text{I}^- \rightarrow 2\text{Br}^- + \text{I}_2$ M1 I ⁻ as a reactant and Br ⁻ as a product M2 equation fully correct	2
6(d)(ii)	M1 343 M2 350 M3 -7	3

Question	Answer	Marks
6(e)(i)	any two from:- <ul style="list-style-type: none">• solid dissolves / solid disappears• bubbling / effervescence / fizzing• melts / forms a ball• floats• moves	2
6(e)(ii)	blue	1