

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Ordinary Level

MARK SCHEME for the October/November 2014 series

5070 CHEMISTRY

5070/42

Paper 4 (Alternative to Practical), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5070	42

- 1 (a) measuring cylinder (1) [1]
- (b) $26(1) \text{ cm}^3$ [1]
- (c) (i) (turns) red (1) [1]
- (ii) bubbles/effervescence **OR** solid dissolves / disappears / forms a solution (1) [1]
- (d) propanol / propan-1-ol (1) [1]
- (e) ethyl propanoate (1)
 $\text{C}_2\text{H}_5\text{COOC}_2\text{H}_5$ or $\text{C}_2\text{H}_5\text{CO}_2\text{C}_2\text{H}_5$ (1) [2]

[Total: 7]

- 2 (a) hydrogen / H_2 **NOT** H (1)
 burning splint pops or pops in a flame (1) [2]
- (b) $\text{Mg} + 2\text{HCl} \rightarrow \text{MgCl}_2 + \text{H}_2$ (1) [1]
- (c) final temperature 35.2
 initial temperature 26.3
 change in temperature 8.9
 all three correct scores 2 marks; two correct scores 1 mark [2]
- (d) exothermic (1) [1]

[Total: 6]

- 3 (a) limewater turns milky (1) [1]
- (b) heat to constant mass (1) [1]
- (c) (i) $0.16(1) \text{ g}$ [1]
- (ii) $0.004(1) \text{ moles}$ [1]
- (iii) $0.004(1) \text{ moles}$ [1]
- (iv) $40(1)$ [1]
- (v) $((\text{iv}) - 16) = 24(1)$ [1]

[Total: 7]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5070	42

4 (d) (1) [Total: 1]

5 (a) (1) [Total: 1]

6 (d) (1) [Total: 1]

7 (b) (1) [Total: 1]

8 (c) (1) [Total: 1]

9 (a) 3.35 (1)g [1]

(b) volumetric flask (1) [1]

(c) (i) pipette (1) [1]

(ii) yellow to red/orange/pink (1) [1]

(d)

23.8	47.8	33.3
<u>0.0</u>	<u>24.3</u>	<u>10.0</u>
23.8	23.5	23.3

 1 mark for each correct row or column to the benefit of the candidate (3)

average volume of 0.100 mol/dm³ HCl = 23.4 (1) cm³ [4]

(e) 0.00234 (1) moles [1]

(f) 0.00117 (1) moles [1]

(g) 0.0117 (1) moles [1]

(h) 286 (1) [1]

(i) (h) – 106 (1)
 $x = (\text{answer} / 18) = 10$ (1)
 answer need not be a whole number but may be rounded up to a whole number [2]

[Total: 14]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5070	42

10 (a) (**Z** is a) compound of a transition metal or transition element or **Z** contains transition metal ions (1)

(b) (i) green ppt (1)

(ii) insoluble (1)

(c) (i) green ppt (1)

(ii) insoluble (1)

(d) (dilute) hydrochloric or nitric acid (1)
aqueous barium chloride or nitrate (1)
white ppt (1)

Conclusion: The formula for **Z** is FeSO_4 . (1)

[Total: 9]

11 (a) gas escapes/lost from apparatus (1) [1]

(b) to allow the gas / vapour to escape (1)
to prevent the liquid from splashing out **OR** to prevent an explosion / flask from bursting / pressure build up / to release the pressure (1) [2]

(c) all points plotted correctly (1)
two smooth curves drawn (1)
curves pass through all points (1) [3]

(d) (i) 0.46(5) (1) g [1]

(ii) $89.55 - 89.47$ (1) = 0.08 (1) g [2]

(iii) manganese (IV) oxide: graph is steeper (at the start) in experiment 1 (1) [1]

(e) all the hydrogen peroxide is used up or has reacted (1) [1]

(f) 89.45 (1) g [1]

[Total: 12]