

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2014 series**

**0610 BIOLOGY**

**0610/61**

Paper 6 (Alternative to Practical), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0610	61

### Abbreviations used in the Mark Scheme

- ; separates marking points
- / separates alternatives within a marking point
- R reject
- I ignore (mark as if this material was not present)
- A accept (a less than ideal answer which should be marked correct)
- AW alternative wording (accept other ways of expressing the same idea)
- underline words underlined (or grammatical variants of them) must be present
- max indicates the maximum number of marks that can be awarded when there are more marking points listed.
- mark independently the second mark may be given even if the first mark is wrong
- A, S, P, L Axes, Size, Plots and Line for graphs
- O, S, D, L Outline, Size, Detail and Label for drawings
- (n)ecf (no) error carried forward (credit a correct operation from a previous wrong response.)
- ( ) the word / phrase in brackets is not required, but sets the context.
- ora or reverse argument.
- AVP Any valid point

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – October/November 2014</b>	<b>0610</b>	<b>61</b>

	<b>Answer</b>	<b>Marks</b>	<b>Guidance for Examiners</b>																											
<b>1 (a)</b>	table drawn with (ruled) lines and cells; headings correct (time, volume and (syringe) 1, 2, 3); units correct in both headings; results recorded in table;;; (1 mark per column completed)	<b>6</b>	<b>A</b> any orientation, outer border not needed  <b>R</b> units within the table  <table border="1"> <thead> <tr> <th rowspan="2">time / min</th> <th colspan="3">volume / cm<sup>3</sup></th> </tr> <tr> <th>(syringe) 1</th> <th>(syringe) 2</th> <th>(syringe) 3</th> </tr> </thead> <tbody> <tr> <td>0</td> <td></td> <td></td> <td></td> </tr> <tr> <td>5</td> <td></td> <td></td> <td></td> </tr> <tr> <td>10</td> <td></td> <td></td> <td></td> </tr> <tr> <td>15</td> <td></td> <td></td> <td></td> </tr> <tr> <td>20</td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	time / min	volume / cm <sup>3</sup>			(syringe) 1	(syringe) 2	(syringe) 3	0				5				10				15				20			
time / min	volume / cm <sup>3</sup>																													
	(syringe) 1	(syringe) 2	(syringe) 3																											
0																														
5																														
10																														
15																														
20																														
<b>(b) (i)</b>	to make the results more reliable/to find anomalies/to calculate an average;	<b>1</b>																												
<b>(ii)</b>	syringe 2 (reading at 15 min/20 min) much lower than others/ syringes 1 and 3 are similar;	<b>1</b>																												
<b>(iii)</b>	16;	<b>1</b>	(18 + 12 + 19 = 49 , 49 / 3 = 16.33 = 16)																											
<b>(c) (i)</b>	30;	<b>1</b>	(35 – 5 = 30)																											

<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – October/November 2014</b>	<b>0610</b>	<b>61</b>

<b>(ii)</b>	<p>A – axes labelled <b>and</b> scaled evenly;</p> <p>S – size,</p> <p>P – all points plotted accurately <math>\pm\frac{1}{2}</math> small square;</p> <table border="1" data-bbox="297 491 1171 616"> <tr> <td>10</td> <td>20</td> <td>30</td> <td>40</td> <td>50</td> <td>60</td> <td>70</td> </tr> <tr> <td>1</td> <td>5</td> <td>15</td> <td>30</td> <td>42</td> <td>25</td> <td>2</td> </tr> </table> <p>L – line drawn;</p>	10	20	30	40	50	60	70	1	5	15	30	42	25	2	<b>4</b>	<p>x-axis: temperature / °C y-axis: average increase in volume / cm<sup>3</sup> <b>I</b> orientation plots to fill half, or more than half, of grid along both axes</p> <p>P = 0 if no scale <b>A</b> ecf <b>(d)(i)</b> <b>A</b> ecf of correct plots on an uneven scale</p> <p>if plot average volume and not average increase in volume = max 3</p> <p><b>A</b> either best fit or point to point, ruled lines or smooth curve <b>R</b> extrapolation &gt; <math>\frac{1}{2}</math> small square <b>R</b> histogram or bar chart</p>
10	20	30	40	50	60	70											
1	5	15	30	42	25	2											
<b>(iii)</b>	<p>as the temperature increases the (average) increase in volume increases to a peak / up to 50 °C;</p> <p>up to 50 °C the (average) increase in volume starts slowly, then increases;</p> <p>above 50 °C the (average increase in) volume slows / increases less / decreases;</p>	<b>max 2</b>	<p><b>A</b> trend– as temperature increases, volume increases then decreases = max 1</p> <p><b>A</b> non-linear / changes gradient</p> <p><b>R</b> volume decreases <b>A</b> ecf for wrong optimum temperature</p>														

<b>Page 5</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – October/November 2014</b>	<b>0610</b>	<b>61</b>

<b>(iv)</b>	yeast activity increases with temperature up to 50 °C; optimum temperature is 50 °C; (some of ) yeast is killed /enzymes become denatured above 50 °C;	<b>max 1</b>	<b>A</b> enzyme activity /metabolism /respiration <b>I</b> volume /growth of yeast <b>R</b> yeast is denatured /enzyme is killed
		<b>[Total: 17]</b>	

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0610	61

2	(a) (i)	<p>drawing of leaf R (monocot):</p> <p><b>O</b> – outline is single clear line (and no shading anywhere);</p> <p><b>S</b> – drawing occupies at least half of the space provided;</p> <p><b>D</b> – detail at least mid-rib and 3 veins each side;</p> <p><b>L</b> – label on midrib;</p>	<p>wrong leaf drawn = max 3 (O, S and L)</p> <p>occupies at least half of the space provided / leaf longer than 50 mm</p> <p><b>R</b> if drawing touches / extends into printed words</p> <p>minimum 7 lines, central line extends full length of leaf, other veins need not connect to base of midrib / petiole</p> <p><b>R</b> ruled lines</p> <p>label lines must make contact with midrib</p>	4
	(ii)	<p>line drawn for widest part of leaf <math>R \pm 1</math> (mm);</p> <p>measurement of widest part of leaf <math>R = 15 \pm 1</math> (mm);</p> <p>mm recorded for at least one measurement;</p>		3
	(iii)	<p>formula: <math>\frac{\text{widest part of drawing}}{\text{widest part of specimen}}</math> ;</p> <p>calculation: magnification correct from their figures;</p>	<p>measurements should be same as in (a)(ii)</p> <p><b>A</b> ecf for cm measurements</p> <p><b>A</b> words or figures</p> <p>answer must be whole number</p>	2

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0610	61

<b>(b) (i)</b>		<b>R</b>	<b>S</b>	<b>A</b> comparative answers on one side only
	shape	narrow / thin / AW	oval / round / wide / AW;	
	venation	parallel / straight / AW	netted / branched / curved / AW;	
	leaf stalk	no petiole	petiole;	
	appearance	shiny / bright / light	dull / dark;	
	edge	smooth	irregular / toothed;	
<b>(ii)</b>	<b>R</b> is monocotyledon as has parallel veins / AW;			<b>1</b>

<b>Page 8</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – October/November 2014</b>	<b>0610</b>	<b>61</b>

<b>(c) (i)</b>	temperature;	keep in the same room / put into an environmental chamber / AW;	<b>max 4</b>	<b>A</b> description e.g. lamp and a heat shield
	idea of no air currents / wind / draughts;	keep all windows and doors closed / idea of a screen around the balance / AW;		
	(sun) light (intensity);	use a light source at a fixed distance / same light source / AW;		<b>A</b> keep in dark
	leaf surface area;	use leaves of same size of leaf / surface area;		
	mark as pairs, one mark for a correct variable and one mark for a suitable method			
<b>(ii)</b>	method of collecting liquid / water / water vapour;		<b>3</b>	<b>A</b> e.g. clip paper to leaf, collect water / liquid / water vapour in bag / tube / box
	test for water: use (dry) cobalt chloride paper / test (liquid) boiling point / freezing point for water;			<b>A</b> any other anhydrous salt
	result: cobalt chloride changes in colour from blue to pink / boiling point 100 °C / freezing point 0 °C;			



<b>Page 9</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – October/November 2014</b>	<b>0610</b>	<b>61</b>

<b>(iii)</b>	<p>similarities: (max 2)</p> <p>both leaves lose water / mass;</p> <p>both leaves lose more water at the start / water loss slows with time;</p> <p>actual loss as percentage of leaf mass is almost the same;</p> <p>differences: (max 2)</p> <p>leaf <b>W</b> loses more water than leaf <b>V</b> / ora;</p> <p>calculation of data;</p> <p>leaf <b>V</b> appears to have anomalous result (at 10/15 min) / leaf <b>V</b> increase in mass between 10 and 15 min / AW;</p> <p>mass leaf <b>V</b> stops losing mass / stays constant at 50 mins;</p>	<b>max 4</b>	<p><b>A W</b> loses water at a faster rate than <b>V</b>.</p> <p><b>A</b> 65% loss for <b>V</b> and 64% loss for <b>W</b>  <b>A</b> leaf <b>W</b> loses 4.8g / leaf <b>V</b> loses 3.4 g / <b>W</b> loses 1.4g more than <b>V</b></p> <p><b>A</b> At 15 min <b>V</b> increases by 1.5g</p>
		<b>[Total: 23]</b>	