



# **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/53

Paper 5 Practical Test October/November 2014

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use
Total

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 7 printed pages and 1 blank page.



You are going to investigate the addition of four different solids, **H**, **J**, **K** and **L**, to wait. The same mass of solid, 4g, will be used in each experiment.

Read all the instructions below carefully before starting the experiments.

#### Instructions

You are going to carry out five experiments.

## (a) Experiment 1

Use a measuring cylinder to pour 25 cm³ of distilled water into the polystyrene cup provided. Support the polystyrene cup in the 250 cm³ beaker. Measure the initial temperature of the water and record it in the table below.

Add all of solid **H** to the water in the cup and stir the mixture with the thermometer. Measure the temperature of the liquid mixture after 90 seconds. Record your result in the table. Remove the thermometer and rinse the thermometer and the cup with water.

### (b) Experiment 2

Repeat Experiment 1, using solid **J** instead of solid **H**.

Measure and record the initial and final temperatures in the table below. **Keep** some of this solution in a test-tube for Experiment 5.

## (c) Experiment 3

Repeat Experiment 1, using solid **K**. Record the temperatures in the table.

### (d) Experiment 4


Repeat Experiment 1 using solid **L**. Record the temperatures in the table and complete the table. **Keep** this solution for Experiment 5.

experiment	solid	initial temperature /°C	final temperature /°C	temperature difference/°C
1	н			
2	J			
3	K			
4	L			

г.	7

	۱ _ ۱	<b>\</b> Experimer	. <i>t</i> r	
1		ı Evnarımar	7 <i>T F</i>	٠

Pour about 3 cm³ of the solution from Experiment 2 into a test-tube. Use a teat pipette to about 3 cm³ of the solution from Experiment 4 to the test-tube. Record your observations.	add



experiment

		bservations			

(g)	(i)	Which experiment produced the smallest temperature change?

(ii) Which solids dissolve in water to produce an endothermic change? Explain your choice.

	121

[4]

(h)	Sug	ggest the temperature change that would occur if	1
	(i)	ggest the temperature change that would occur if  Experiment 3 was repeated using 50 cm³ of distilled water,	708
			7
	(ii)	2g of solid <b>L</b> were used in Experiment 4.	
	iii)	Explain your answer to (h)(ii).	[1]
			[1]
(i)	Pre	edict the temperature of the solution in Experiment 2 after one hour. Explain your answer	
(j)	Suc	ggest an explanation for the observations in Experiment 5.	
•			
			[2]
		[Total: 2	20]

5	I and N. ling all of your observations in the table.  observations
	abo.
You are provided with two aqueous solutions, M Carry out the following tests on M and N, record Conclusions must not be written in the table.	and <b>N</b> . ling all of your observations in the table.
tests	observations
tests on solution <b>M</b>	
Divide solution ${\bf M}$ into four equal portions in separate test-tubes.	
(a) Describe the appearance of solution M.	
Test the pH of the first portion of <b>M</b> .	[1]
(b) Add a spatula measure of calcium carbonate to	
the second portion of <b>M</b> .  Test the gas given off with a splint.	[2]
(c) To the third portion of <b>M</b> , add magnesium ribbon.	
Test the gas given off with a splint.	
	[3]
(d) To the fourth portion of M, add a few drops of	
dilute nitric acid and about 1 cm <sup>3</sup> of aqueous silver nitrate.	[2]

[Turn over

	Age.
tests	observations
tests on solution N	observations Add Cannibridge
Divide solution <b>N</b> into three equal portions in separate test-tubes.	
(e) Describe the appearance of solution N.	[1]
Test the pH of the first portion of solution N.	[1]
(f) Use a teat pipette to add three to four drops of aqueous zinc sulfate to the second portion of N. Shake the mixture.	
Now add excess aqueous zinc sulfate to the mixture and shake.	[3]
(g) To the third portion of <b>N</b> , add a spatula measure of ammonium chloride. Warm the mixture and test the gas given off with damp litmus paper.	[2]
(h) (i) Identify the gas given off in test (c).	[41]
	[1]
(ii) Identify the gas given off in test (g).	
	[1]
(i) Identify solution M.	
	[2]
(j) What conclusion can you draw about solution	on <b>N</b> ?
	[1]
	[Total: 20]

7

**BLANK PAGE** 

### **NOTES FOR USE IN QUALITATIVE ANALYSIS**

### **Test for anions**

Test for anions	8 NOTES FOR USE IN QUALITATIVE	E ANALYSIS  test result  effervescence, carbon dioxide
anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

# Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> +)	ammonia produced on warming	
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

# **Test for gases**

gas	test and test results
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.