



**Cambridge International Examinations**  
Cambridge International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**CHEMISTRY**

**0620/42**

Paper 4 Theory (Extended)

**October/November 2017**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **14** printed pages and **2** blank pages.



1 (a) Dust particles in the air move around in a random way.

(i) What term describes the random movement of the dust particles?

..... [1]

(ii) Identify the particles in the air which cause the random movement of the dust particles.

..... [2]

(iii) Explain why the dust particles move in this way.

.....


.....

..... [2]

(b) When chlorine gas,  $Cl_2$ , is put into a gas jar, it spreads out to fill the gas jar.

When bromine gas,  $Br_2$ , is put into a gas jar, it also spreads out to fill the gas jar.

The process takes longer for bromine gas than for chlorine gas.



(i) What term describes the way that the gas particles spread out?

..... [1]

(ii) Use **data** from the Periodic Table to explain why bromine gas takes longer to fill a gas jar than chlorine gas.

.....

.....

..... [2]

(iii) Explain why increasing the temperature increases the rate at which the gas particles spread out.

.....

..... [1]

[Total: 9]

- 2 (a) Complete the table to show the electronic structure of the atoms and ions.

	electronic structure
F	2,7
Si	
Ca <sup>2+</sup>	
N <sup>3-</sup>	

[3]

- (b) Predict the formula of the compound formed between Ca<sup>2+</sup> and N<sup>3-</sup>.

..... [1]

- (c) Draw a dot-and-cross diagram to show the electron arrangements in the **two** ions present in lithium chloride, LiCl.  
Show outer shell electrons only. Include the charges on the ions.

[3]

- (d) Sulfur dichloride, SCl<sub>2</sub>, is a covalent compound. It has the structure Cl–S–Cl.

Draw a dot-and-cross diagram to show the electron arrangement in a molecule of sulfur dichloride.  
Show outer shell electrons only.

[3]

(e) In terms of attractive forces, explain why  $\text{LiCl}$  has a higher melting point than  $\text{SCl}_2$ .


.....  
.....  
.....  
.....  
..... [3]

(f) Suggest the identity of a **covalent compound** with a higher melting point than  $\text{LiCl}$ .

..... [1]


[Total: 14]

- 3 The chemical equation for the complete combustion of ethanol,  $\text{C}_2\text{H}_5\text{OH}$ , is shown.



The energy released when one mole of ethanol undergoes complete combustion is 1280 kJ.

Part of the energy level diagram for this reaction is shown.




- (a) Complete the energy level diagram to show
- the products of the reaction,
  - the overall energy change of the reaction.

[3]


- (b) What does **X** represent?

..... [1]

(c) The chemical equation for the complete combustion of methanol, CH<sub>3</sub>OH, is shown.



The equation can be represented as shown.



Use the bond energies in the table to determine the energy change,  $\Delta H$ , for the complete combustion of **one** mole of methanol.

bond	bond energy in kJ/mol
C-H	410
C-O	360
O-H	460
O=O	500
C=O	805

- energy needed to break bonds

..... kJ

- energy released when bonds are formed

..... kJ

- energy change,  $\Delta H$ , for the complete combustion of **one** mole of methanol

..... kJ/mol  
[4]

- (d) Dodecane is an alkane containing 12 carbon atoms. Ethanol can be manufactured from dodecane in a two-stage process.

In **stage 1**, each molecule of dodecane is converted into three molecules of ethene and one molecule of another hydrocarbon.

- (i) Name the process which occurs in **stage 1**.

..... [1]

- (ii) Write a chemical equation for the reaction which occurs in **stage 1**.

..... [2]

In **stage 2**, ethene reacts with steam to produce ethanol.

- (iii) State **two** conditions needed for **stage 2**.

1 .....

2 ..... [2]

- (iv) Name the type of reaction which occurs in **stage 2**.

..... [1]

- (v) Suggest how to test the purity of the ethanol produced.

.....

..... [2]


(e) Ethanol can also be manufactured by the fermentation of glucose,  $C_6H_{12}O_6$ .

(i) State **two** conditions needed for the fermentation of glucose.

1 .....

2 ..... [2]

(ii) Complete the chemical equation for the fermentation of glucose.



(iii) One disadvantage of fermentation is that the maximum concentration of ethanol produced is about 15%.

Suggest why the concentration of ethanol produced by fermentation does **not** exceed 15%.

.....

..... [1]

(iv) Give **one** other disadvantage of manufacturing ethanol by fermentation.

..... [1]

(v) Give **one** advantage, other than cost, of manufacturing ethanol by fermentation.


..... [1]

(vi) Suggest the name of a process to obtain ethanol from a mixture of ethanol and water.

..... [1]



(f) Ethane-1,2-diol has the following structure.



(i) Write the empirical formula of ethane-1,2-diol.

..... [1]

(ii) Ethane-1,2-diol can undergo condensation polymerisation but cannot undergo addition polymerisation.

Explain why ethane-1,2-diol **cannot** undergo addition polymerisation.


.....  
 ..... [1]

(iii) Ethane-1,2-diol undergoes condensation polymerisation with molecule Y.

The diagrams represent the structures of ethane-1,2-diol and molecule Y.



ethane-1,2-diol



molecule Y

Draw the condensation polymer formed between ethane-1,2-diol and molecule Y.  
 Show **one** repeat unit. Show all of the atoms and all of the bonds in the linkage.


[3]

(iv) Name the type of condensation polymer formed between ethane-1,2-diol and molecule Y.

..... [1]

[Total: 30]

- 4 A student sets up the following electrolysis experiment.



- (a) Define the term *electrolysis*.

.....  
 ..... [2]

- (b) The student observes bubbles of colourless gas forming at each electrode.

- (i) Name the main gas produced at the positive electrode (anode).

..... [1]

- (ii) Describe a test for the gas produced in (b)(i).

test .....

result ..... [2]

- (iii) Write the ionic half-equation for the reaction taking place at the negative electrode (cathode).

..... [2]

- (c) Charge is transferred during electrolysis.

Name the type of particle responsible for the transfer of charge in

the wires, .....

the electrolyte. ....

[2]

- (d) The student replaces the dilute aqueous sodium chloride with **concentrated** aqueous sodium chloride.

Suggest **two** differences that the student observes.

1 .....


2 .....

[2]

- (e) The student has a small piece of impure copper. The main impurities in the copper are small quantities of silver and zinc.

The student uses electrolysis to extract pure copper from the small piece of impure copper.

- (i) Complete the labels on the diagram of the student's electrolysis experiment.



[3]

- (ii) Use your knowledge of the reactivity series to suggest what happens to the silver and zinc impurities. Explain your answers.

silver impurities .....

.....

.....

zinc impurities .....

.....

.....

[3]


[Total: 17]

5 Some chemical reactions are reversible.

(a) Aqueous potassium chromate(VI),  $K_2CrO_4$ , is a yellow solution.

Aqueous potassium dichromate(VI),  $K_2Cr_2O_7$ , is an orange solution.

The two compounds interconvert when the pH of the solution changes.



Solution **Y** is a mixture of aqueous potassium chromate(VI) and aqueous potassium dichromate(VI) at equilibrium.

- Explain, in terms of the position of the equilibrium, what you would **see** if sulfuric acid were added to solution **Y**.

.....

.....

.....

- Explain, in terms of the position of the equilibrium, what you would **see** if sodium hydroxide were added to solution **Y**.

.....


.....

.....

.....

[5]

- (b) Hydrogen can be manufactured using a reversible reaction between methane and steam.



At 900 °C, in the presence of a nickel catalyst, the yield of hydrogen is 70%.

- (i) What volume of hydrogen is produced from 100 cm<sup>3</sup> of methane under these conditions?

..... cm<sup>3</sup> [2]

Under different conditions, different yields of hydrogen are obtained.

- (ii) If the pressure is increased, the yield of hydrogen becomes less than 70%.

Explain why, in terms of the position of the equilibrium.

.....  
..... [1]

- (iii) If the temperature is decreased, the yield of hydrogen decreases.

What does this information indicate about the reaction between methane and steam?

..... [1]

- (iv) Why is a catalyst used in this reaction?

..... [1]

[Total: 10]



**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

## The Periodic Table of Elements

		Group							
I	II	III	IV	V	VI	VII	VIII		
1	2	3	4	5	6	7	8	9	10
H hydrogen 1	He helium 4	B boron 11	C carbon 12	N nitrogen 14	O oxygen 16	F fluorine 19	Ne neon 20		
<b>Key</b>									
atomic number atomic symbol name relative atomic mass									
11	12	13	14	15	16	17	18		
Na sodium 23	Mg magnesium 24	Al aluminium 27	Si silicon 28	P phosphorus 31	S sulfur 32	Cl chlorine 35.5	Ar argon 40		
19	20	21	22	23	24	25	26	27	28
K potassium 39	Ca calcium 40	Sc scandium 45	Ti titanium 48	V vanadium 51	Cr chromium 52	Mn manganese 55	Fe iron 56	Co cobalt 59	Ni nickel 59
37	38	39	40	41	42	43	44	45	46
Rb rubidium 85	Sr strontium 88	Y yttrium 89	Zr zirconium 91	Nb niobium 93	Mo molybdenum 96	Tc technetium —	Ru ruthenium 101	Rh rhodium 103	Pd palladium 106
55	56	57–71	72	73	74	75	76	77	78
Cs caesium 133	Ba barium 137	lanthanoids	Hf hafnium 178	Ta tantalum 181	W tungsten 184	Re rhenium 186	Os osmium 190	Ir iridium 192	Pt platinum 195
87	88	89–103	104	105	106	107	108	109	110
Fr francium —	Ra radium —	actinoids	Rf rutherfordium —	Db dubnium —	Sg seaborgium —	Bh bohrium —	Hs hassium —	Mt meitnerium —	Ds darmstadtium —
atomic number atomic symbol name relative atomic mass									
81	82	83	84	85	86	87	88	89	90
Tl thallium 204	Pb lead 207	Bi bismuth 209	Po polonium —	At astatine —	Rn radon —	Cn copernicium —	Nh nihonium —	Dl dubnium —	Fl flerovium —
91	92	93	94	95	96	97	98	99	100
Ac actinium —	Th thorium 232	Pa protactinium 231	U uranium 238	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —
101	102	103	104	105	106	107	108	109	110
Lr lawrencium —	No nobelium —	Lr lawrencium —	111	112	113	114	115	116	117
atomic number atomic symbol name relative atomic mass									
111	112	113	114	115	116	117	118	119	120
111	112	113	114	115	116	117	118	119	120

lanthanoids

actinoids

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La lanthanum 139	Ce cerium 140	Pr praseodymium 141	Nd neodymium 144	Pm promethium —	Sm samarium 150	Eu europium 152	Gd gadolinium 157	Tb terbium 159	Dy dysprosium 163	Ho holmium 165	Er erbium 167	Tm thulium 169	Yb ytterbium 173	Lu lutetium 175
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac actinium —	Th thorium 232	Pa protactinium 231	U uranium 238	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —	Es einsteinium —	Fm fermium —	Md mendelevium —	No nobelium —	Lr lawrencium —

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).