

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

VS VS CONTINUE CONTIN

CENTRE NUMBER
COMBINE Paper 5 P

**CANDIDATE** NAME CENTRE **CANDIDATE** 

NUMBER

0653/52 COMBINED SCIENCE

Paper 5 Practical Test May/June 2010

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in Instructions to Supervisors.

### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Chemistry practical notes for this paper are printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
Total		

This document consists of 11 printed pages and 5 blank pages.



- 1 This question is about variation in leaves.
- For iner's tiole evenly on (a) You are provided with 20 leaves of the same species. Measure the length *l* of each leaves in millimetres as shown in Fig. 1.1a. If the lamina does not meet the petiole evenly on either side of the leaf use the longer measurement. See Fig. 1.1b.

Enter your measurements in Table 1.1.




Fig. 1.1a




Fig. 1.1b

Table 1.1

length of leaf // mm				
1	11			
2	12			
3	13			
4	14			
5	15			
6	16			
7	17			
8	18			
9	19			
10	20			

[2]

**(b)** Calculate the average (mean) length of the 20 leaves. Show your working.

leaf area = \_\_\_\_\_cm<sup>2</sup>

[3]

		4	100
(c)	The	e difference between the greatest length and the smallest length is the ra	nge. Parta Ca
	Cor	mplete the following.	
	the	greatest length =mm	
	the	smallest length =mm	
	the	range =mmm	[1]
(d)		e the grid provided on page 5 to estimate the area of <b>one</b> of the leaves. ch square is 1 cm <sup>2</sup> .	The area of
	•	Place the leaf on the grid provided.	
	•	Carefully draw round the leaf then remove it.	
	•	Write the letter <b>C</b> in the <b>complete</b> squares. Count the number of squares.	of complete
		number of complete ( <b>C</b> ) squares =	
	•	Write the letter ${\bf P}$ in any incomplete squares that have an area of half more.	a square or
		number of incomplete ( <b>P</b> ) squares =	
	•	Ignore the rest of the squares.	

Add **C** + **P** to estimate the area of the leaf.

(e)	The leaves in the sample were all of the same species yet they showed variation in
	length.

Suggest and explain a reason for this.

reason	
explanation	
	[2]

[2]

		www.xtra	papers.co
		6	
		going to find the specific heat capacity of the material of a can. The specific of a material is the heat energy required to raise 1 g of the material by 1 °C. In the mass of the can to the nearest gram.	For iner's
(a)	Find	d the mass of the can to the nearest gram.	Tage
	Red	cord its mass below.	.col
		mass of can, $\mathbf{m}_1$ , =g [1]	
(b)		ce the lagging around the can. Place the thermometer inside the can and leave fo minutes. Read the temperature, $\mathbf{t_1}$ , to the nearest 0.5 °C and record it below.	r
		temperature of can, $\mathbf{t_1} = \underline{}^{\circ} \mathbf{C}$ [1	]
(c)	(i)	Heat enough water in a beaker to about one-third fill the can. When the temperature is just above 70 °C, remove the Bunsen. As soon as the temperature of the water has cooled to exactly $70.0^{\circ}$ C pour the water into the can. Read the temperature, $t_2$ , to the nearest 0.5 °C of the water after exactly <b>two minutes</b> Record this temperature.	e e
		temperature of water, <b>t</b> <sub>2</sub> =°C [1	]
	(ii)	Remove the lagging and pour the water into a measuring cylinder. Record the volume.	e
		volume of water =cm <sup>3</sup> [1	]
(	(iii)	$1~\text{cm}^3$ of water has a mass of 1 g. Calculate the mass, $\mathbf{m_2}$ , of the volume of wate you recorded in <b>(c)(ii)</b> .	r
		mass of water, $\mathbf{m_2} = \phantom{aaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaa$	1
(d)	Cal	culate	
	(i)	$\mathbf{t_3}$ , the fall in temperature of the hot water, $\mathbf{t_3} = (70.0 - \mathbf{t_2})$ .	
		<b>t</b> <sub>3</sub> =°C	
	(ii)	$\mathbf{t_4}$ , the rise in temperature of the can, $\mathbf{t_4} = (\mathbf{t_2} - \mathbf{t_1})$ .	

www.xtrapapers.cor	n
* Odda	
al of the For iner's	
Tage Co.	1
The state of the s	

(e) Use the equation to calculate the specific heat capacity, shc, of the material of the

$$shc = \frac{\mathbf{m_2} \times \mathbf{t_3} \times 4.2}{\mathbf{t_4} \times \mathbf{m_1}}$$

specific heat capacity of the material of the can = 
$$_{\text{max}} J g^{-1} \circ C^{-1}$$
 [3]

Www.xtrapapers.com

3 You are going to investigate the rate of reaction between magnesium and hydrochlon

Read through the procedure before starting the experiment.

- (a) (i) Set up the apparatus as shown in Fig. 3.1.
  - Fill the 100 cm<sup>3</sup> measuring cylinder and trough with water.




Fig. 3.1

- (ii) Place 20 cm<sup>3</sup> of the hydrochloric acid in the large test-tube.
  - Cut 6 cm of magnesium ribbon from the length provided.
  - Loosely fold the piece of magnesium ribbon and place it in the acid contained in the test-tube. Immediately replace the stopper and delivery tube and start the timer.
  - Read the volume of gas in the measuring cylinder after 20, 40, 60 and 80 seconds.
  - Record the volumes in Table 3.1. [2]
- **(b) (i)** You will now repeat the procedure using the same length of magnesium but different volumes of acid and water.
  - Wash out the contents of the test-tube.
  - Refill the measuring cylinder with water.
  - Place 16 cm<sup>3</sup> of hydrochloric acid in the test-tube and 4 cm<sup>3</sup> of water.
  - Cut 6 cm of magnesium ribbon and place it in the acid. Replace the stopper and delivery tube.
  - Immediately start the timer.
  - Read the volume of gas in the measuring cylinder after 20, 40, 60 and 80 seconds.
  - Record the volumes in Table 3.1.

(ii) Repeat the experiment <b>two</b> more times using volumes of acid and water as in Table 3.1. Record the results in Table 3.1.  Table 3.1							apapers.com
volume of 2 mol/dm³							on on
hydrochloric acid/cm <sup>3</sup>		mixture/mol/ dm <sup>3</sup>	20 s	40 s	60 s	80 s	
20	0	2.0					]   I
16	4	1.6					]
12	8	1.2					]
4	16	0.4					]

11

(c) Draw a graph of volume of gas collected after 40 s (vertical axes) against concert of hydrochloric acid. Include the origin in your plots and draw a smooth curve.

For iner's

volume of gas collected after 40 s/cm<sup>3</sup>

concentration of acid/mol dm-3

(d)	How is the rate of reaction affected by concentration of acid? Explain how your enable you to decide this.	bric
	[2]	
(e)	Had any of the reactions finished by the time 80 s had been reached? Explain your answer.	
	[1]	

14

**BLANK PAGE** 

Www.xtrapapers.com

### **CHEMISTRY PRACTICAL NOTES**

### **Test for anions**

Test for anions	16 CHEMISTRY PRACTICAL NO	TES test result
anion	test	test result
carbonate (CO <sub>3</sub> <sup>2</sup> -)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ·) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
nitrate (NO <sub>3</sub> -) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate	white ppt.

# Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

## **Test for gases**

gas	test and test results
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	"pops" with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.