

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		



CHEMISTRY 0620/52

Paper 5 Practical Test May/June 2011

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use			
1			
2			
Total			

This document consists of 7 printed pages and 1 blank page.



1 You are going to investigate the reaction between excess magnesium and two different dilute acids, **X** and **Y**.

For Examiner's Use


Read all the instructions below carefully before starting the experiments.

Instructions

You are going to carry out two experiments.

(a) Experiment 1

Set up the apparatus as shown in the diagram below.



Remove the bung from the conical flask and move the measuring cylinder away from the delivery tube without letting any water run out. Twist one of the strips of magnesium provided to break it into four pieces and place all of them into the conical flask.

Using a different measuring cylinder, add $50\,\mathrm{cm^3}$ of aqueous acid **X** into the conical flask and replace the bung firmly. Place the measuring cylinder back over the delivery tube and start the timer. In the table, record the volume of gas collected in the measuring cylinder every thirty seconds for three minutes.

time/s	0	30	60	90	120	150	180
volume of gas/cm ³							

[3]

(b) Experiment 2

Repeat the whole of Experiment 1 using 50 cm³ of aqueous acid **Y**. In the table, record the volume of gas collected in the measuring cylinder every thirty seconds for three minutes.


time/s	0	30	60	90	120	150	180
volume of gas/cm ³							

[3]

© UCLES 2011 0620/52/M/J/11

(c) Plot the results you have obtained for both experiments on the grid below. For each set of results, draw a smooth line graph. Indicate clearly which line represents Experiment 1 and which line represents Experiment 2.

For Examiner's Use



(ii) Suggest why the speed was fastest in this experiment.

[1]

(iii) Suggest why the speed was fastest in this experiment.

[1]

(e) Why eventually will no more gas be produced?

[2]

4

(f)	From your graph , deduce the time required to collect 25 cm³ of gas in Experiment 1. Show clearly on the graph how you worked out your answer.
	[2]
(g)	Why was the measuring cylinder moved away from the delivery tube while the acid solution was added to the flask?
	[2]
(h)	Give one advantage and one disadvantage of using a measuring cylinder to add the acids to the flask.
	advantage
	disadvantage[2]
	[Total: 22]

For Examiner's Use

For Examiner's Use

2 You are provided with mixture **Z**.

Z consists of two solids, **W** and **V**. Solid **W** is water-soluble and solid **V** is insoluble. Carry out the following tests on **Z**, recording all of your observations in the table. Conclusions must **not** be written in the table.

	tests	observations
test	s on mixture Z	
(a) Describe the appearance of the mixture.		[1]
(b)	Place a little of mixture Z in a test-tube. Heat the mixture for about 1 minute. After 1 minute, test the gas with damp pH indicator paper. Leave the test-tube to cool.	[3]
disti Stor con con Kee follo	the rest of mixture Z to about 10 cm ³ of lled water in a boiling tube. Oper the boiling tube and shake the tents for about one minute. Filter the tents of the test-tube. In the residue and the filtrate for the wing tests.	
iesi	s on the filtrate	
(c) (i) To about 1 cm³ of the filtrate, add a few drops of dilute nitric acid followed by about 1 cm³ of silver nitrate solution and shake the test-tube.		[2]
(To about 1 cm³ of the filtrate, add about 1 cm³ of aqueous sodium hydroxide. Gently heat the mixture. Test the gas given off with damp pH indicator paper.	[1]
(i	ii) To the third portion of the filtrate, add about 1 cm³ of dilute hydrochloric acid followed by about 1 cm³ of barium chloride solution.	[1]

For Examiner's

tests	observations
tests on the residue	
(d) Using a spatula, transfer some of the residue from the filter paper to a test-tube. Using a teat pipette, add about 2 cm³ of dilute hydrochloric acid to the residue. Test the gas given off.	[3]
Add to the solution in the test-tube an equal volume of distilled water. Shake the contents and divide into two portions.	
(e) (i) To the first portion of the solution, add excess aqueous sodium hydroxide.	[2]
(ii) To the second portion, add excess aqueous ammonia solution.	[1]
(f) What conclusions can you draw about s	solid W ?
(g) What conclusions can you draw about s	[2] solid V ?
	[2]
	[Zotal: 18]

[Total: 18]

© UCLES 2011 0620/52/M/J/11 7

BLANK PAGE

© UCLES 2011 0620/52/M/J/11

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C1 ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻⁾ [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia	
aluminium (Al³+)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess	
ammonium (NH ₄ ⁺) ammonia produced on warming		_	
calcium (Ca ²⁺) white ppt., insoluble in excess		no ppt., or very slight white ppt.	
copper (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution	
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess	
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess	
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution	

Test for gases

gas	test and test results		
ammonia (NH ₃)	turns damp red litmus paper blue		
carbon dioxide (CO ₂)	turns limewater milky		
chlorine (Cl ₂)	bleaches damp litmus paper		
hydrogen (H ₂)	'pops' with a lighted splint		
oxygen (O ₂)	relights a glowing splint		

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2011 0620/52/M/J/11