

0

Paper 6 Alternative to Practical

October/November 2014

1 hour

Candidates answer on the Question Paper.

No Additional Materials are required.

### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 12 printed pages.



[Turn over







	www.xtrapapers.com
	3
(d) (i)	How would the student know when all of the dilute nitric acid had reacted in
(ii)	What method is used to separate the mixture in stage 3?
(iii)	What term is used to describe the unreacted lead(II) oxide? [1]
<b>(e)</b> De furt	scribe the effect of heating the solution of lead(II) nitrate until it boils and then heating for a ther ten minutes.
	[2]
	[Total: 10]

[Turn over



position for ten minut cm<sup>3</sup> of water and 4 2 Two experiments were carried out to show what factors affect the rate of decomposition peroxide,  $H_2O_2$ . In each experiment the volume of gas produced was measured every minute for ten minu

#### Experiment 1

The student used a mixture of 50 cm<sup>3</sup> of hydrogen peroxide, 50 cm<sup>3</sup> of water and 1 g of manganese(IV) oxide at a room temperature of 20 °C. The results were plotted to obtain the graph shown.

### Experiment 2

The student repeated Experiment 1 but did not record how much of each substance was used. The points were plotted on the grid.



	www.xtrapapers.com
	5
(a)	Complete the graph for Experiment 2.
(b)	Suggest the composition of the mixture used in Experiment 2. Explain your suggestion.
	composition
	explanation
	[4]
(c)	What is the function of the manganese(IV) oxide?
	[1]
(d)	Sketch on the grid the curve that you would expect if Experiment 1 was repeated at 10 $^{\circ}$ C. [2]

[Total: 8]

**3** A student investigated the colours present in a fruit drink. The fruit drink was tested to check that no artificial colours had been added. The apparatus below was used.



(a) (i) Name the method used.
[1]
(ii) Why is there a glass cover on the beaker?
[1]
(b) When should the paper be removed from the beaker?
[1]

[Turn over

#### © UCLES 2014



A student investigated the reaction between two different solutions of dilute hydrod 4 and **B**, and solution **C** which is alkaline.

Two experiments were carried out.

(a) Experiment 1

www.papaCambridge.com A burette was filled with solution A of dilute hydrochloric acid to the 0.0 cm<sup>3</sup> mark. Using a measuring cylinder, 20 cm<sup>3</sup> of solution **C** was poured into a conical flask. A few drops of methyl orange were added to the flask.

Solution **A** was added to the flask, with shaking, until the mixture just changed colour. Use the burette diagram to record the burette reading in the table and complete the table.



initial reading

final burette reading/cm <sup>3</sup>	
initial burette reading/cm <sup>3</sup>	
difference/cm <sup>3</sup>	

[2]

(b) Experiment 2

The burette was emptied and rinsed, first with distilled water, and then with a little of solution **B**. The burette was filled with solution **B** of dilute hydrochloric acid to the 0.0 cm<sup>3</sup> mark.

Experiment 1 was repeated using solution B.

Use the burette diagram to record the burette reading in the table and complete the table.



#### final reading

final burette reading/cm3	
initial burette reading/cm <sup>3</sup>	
difference/cm <sup>3</sup>	

[2]

[Turn over

	www.xtrapa	pers.c
	8	
(c) (i)	What type of chemical reaction takes place when hydrochloric acid reacts solutions?	brids
(ii)	Why is methyl orange added to the flask?	9¢.co.
( <b>d)</b> Wh Exp	y was the burette rinsed, first with distilled water and then with solution <b>B</b> , before star periment 2?	ting
		[2]
(e) (i)	In which experiment was the greater volume of dilute hydrochloric acid used?	[1]
(ii)	Compare the volumes of dilute hydrochloric acid used in Experiments 1 and 2.	[1]
(iii)	Suggest, in terms of the concentration of solutions <b>A</b> and <b>B</b> , an explanation for difference in volumes used.	the
(f) If E aci	experiment 2 was repeated using 10 cm <sup>3</sup> of solution <b>C</b> , what volume of dilute hydroch d would be used? Explain your answer.	[2] oric
		[2]
<b>(g)</b> Giv	e <b>one</b> advantage and <b>one</b> disadvantage of using a measuring cylinder for solution <b>C</b> .	
adv	vantage	[2]

### © UCLES 2014

	www.xtrapapers	.com
	9	
(h)	Describe a method other than titration, using a <b>different</b> reactant, that could be compare the concentrations of the two solutions of dilute hydrochloric acid, <b>A</b> and <b>b</b>	
		com
		-
	[4]	
	[Total: 20]	

[Turn over



		www.xtrapape	ers.com
5	A solid <b>D</b> , which is a soluble metal su The tests on <b>D</b> , and some of the obse Complete the observations in the tabl	Ifate, was analysed. ervations, are in the following table. le.	to la
	tests	observations	CO2.
tes	ts on solid <b>D</b>		1
(a)	(i) Appearance of solid <b>D</b> .	pale green crystals	
	(ii) Solid <b>D</b> was heated in a test-tube gently and then strongly.	condensation formed at the top of the test-tube	
tes	s on the aqueous solution		
Solid <b>D</b> was added to distilled water and shaken to dissolve. The solution was divided into four equal portions in separate test-tubes.			
(b)	(i) Several drops of aqueous sodium hydroxide were added to the first portion of the solution.	green precipitate	
	Excess aqueous sodium hydroxide was added to the mixture.	green precipitate remained	
	(ii) Excess aqueous ammonia was added to the second portion of the solution.	green precipitate	
(c)	Aqueous silver nitrate and dilute nitric acid were added to the third portion of the solution.	[1]	
(d)	Aqueous barium nitrate and dilute nitric acid were added to the fourth portion of the solution.		



### 6

### **Fizzy water**

Fizzy water contains carbon dioxide dissolved under pressure. When the water is heated, the gas is given off.

(a) (i) Complete the labelled diagram to show how you could collect and measure the volume of gas given off when fizzy water is heated.



(ii) State a test for carbon dioxide.

.....[2]

[Turn over

[2]





[Total: 8]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.