



*Rewarding Learning*

**ADVANCED SUBSIDIARY (AS)  
General Certificate of Education  
2019**

---

## **Chemistry**

**Assessment Unit AS 3**

*assessing*

**Module 3: Practical Examination**

**Practical Booklet B**

**[SCH32]**

**WEDNESDAY 29 MAY, AFTERNOON**

---

**MARK  
SCHEME**

## Annotation

1. Please do all marking in **red** ink.
2. All scripts should be checked for mathematical errors. Please adopt a system of one tick (✓) equals 1 mark, e.g. if you have awarded 4 marks for part of a question then 4 ticks (✓) should be on this candidate's answer.
3. The total mark for each question should be recorded in a circle placed opposite the question number in the teacher mark column.
4. As candidates have access to scripts please do not write any inappropriate comments on their scripts.

## General points

- All calculations are marked according to the number of errors made.
- Errors can be carried through. If the wrong calculation is carried out then the incorrect answer can be carried through. One mistake at the start of a question does not always mean that all marks are lost.
- Listing is when more than one answer is given for a question that only requires one answer, e.g. the precipitate from a chloride with silver nitrate is a white solid; if the candidate states a white or a cream solid, one answer is correct and one answer is wrong. Hence they cancel out.
- Although names might be in the mark scheme it is generally accepted that formulae can replace them. Formulae and names are often interchangeable in chemistry.
- The marking of colours is defined in the 'CCEA GCE Chemistry Acceptable Colours' document.

## MARKING GUIDELINES

### Interpretation of the Mark Scheme

- **Carry error through**  
This is where mistakes/wrong answers are penalised when made, but if carried into further steps of the question, then no further penalty is applied. This pertains to calculations and observational/ deduction exercises. Please annotate candidates' answers by writing the letters c.e.t. on the appropriate place in the candidates' answers.
- **Oblique/forward slash**  
This indicates an acceptable alternative answer(s).
- **Brackets**  
Where an answer is given in the mark scheme and is followed by a word/words in brackets, this indicates that the information within the brackets is non-essential for awarding the mark(s).

- 1 (a) (i) effervescence/bubbling/fizzing [1]
- (ii) oxidation number of C in charcoal increased from 0 to +2 in carbon monoxide/oxidation number of C in CO<sub>2</sub> reduced from +4 to +2 [1]  
reducing agent is oxidised [1] [2]
- (iii) more particles with energy greater/equal to E<sub>act</sub> [1]  
more frequent/successful collisions [1] [2]
- (iv) incomplete reduction of carbon dioxide/charcoal not heated enough/  
carbon dioxide passed through too quickly (soon) [1]
- (v) CO<sub>2</sub> + 2NaOH → Na<sub>2</sub>CO<sub>3</sub> + H<sub>2</sub>O [1]
- (vi) add dilute acid [1]  
bubble gas produced through limewater [1]  
turns milky [1] [3]
- (b) 100 kg of Fe<sub>2</sub>O<sub>3</sub> = 100,000 g  
RFM of Fe<sub>2</sub>O<sub>3</sub> = 160  
moles of Fe<sub>2</sub>O<sub>3</sub> = 625
- 56 kg of CO = 56,000 g  
RMM of CO = 28  
moles of CO = 2000
- Fe<sub>2</sub>O<sub>3</sub> is limiting factor
- moles of Fe = 1250  
mass of Fe = 1250 × 56 = 70,000 g = 70 kg  
(error [-1]) [3]
- (c) (i) (high temperature) decreases yield [1]
- (ii) (high pressure) increases yield [1]
- (d) (i) to ensure the even distribution of heat [1]
- (ii) surround apparatus with screen to reduce draughts [1]  
place a lid on the copper can [1] [2]
- (iii) % error =  $\frac{2 \times 0.5}{70.5} \times 100 = 1.4\% = 1\%$  [2]
- (iv) mass of methanol = 20.33 – 18.92 = 1.41 g  
moles of methanol = 1.41/32 = 141/3200 (0.0440625)
- q = (-)100 × 4.2 × 70.5 = (-)29610 J (29.61 kJ)
- ΔH = -29610/0.044 = -672954.5 J (mol<sup>-1</sup>)  
= -673 kJ (mol<sup>-1</sup>) = -700 kJ (mol<sup>-1</sup>)  
(error [-1]) [3]
- (v) carbon [1]  
incomplete combustion [1] [2]
- (vi) more (bonds) broken/formed [1]

AVAILABLE  
MARKS

26

- 2 (a) (i)  $\text{H}_2\text{SO}_4 + \text{NaCl} \rightarrow \text{NaHSO}_4 + \text{HCl}$  [1]
- (ii)  $\text{MnO}_2 + 4\text{HCl} \rightarrow \text{Cl}_2 + \text{MnCl}_2 + 2\text{H}_2\text{O}$  [2]
- (iii) dissolves in the water (to form hydrochloric acid) [1]
- (iv) heavier than air [1]
- (v) bleaches damp Universal/litmus Indicator paper (error [-1]) [2]
- (b) (i) a solution of chlorine water is green [1]
- (ii) add silver nitrate solution [1]  
white precipitate is formed [1] [2]
- (c) (i) two layers are formed [1]
- (ii) the bottom layer turns colourless  
or  
the upper layer turns (pale) green [1]
- (iii) pale green colour disappears [1]
- (iv) separating funnel [1]
- (d) (i)
- |   | C:   | H:   | Cl   |
|---|------|------|------|
| Mass  | 32.2 | 4.5  | 63.3 |
| R.A.M   | 12   | 1    | 35.5 |
| Moles   | 2.68 | 4.5  | 1.78 |
| Ratio   | 1.50 | 2.52 | 1.00 |
|   | 3    | 5    | 2    |
| Empirical formula = $\text{C}_3\text{H}_5\text{Cl}_2$ |      |      |      |
- (error [-1]) [3]
- (ii) mass of hexane = 9.75 g  
RMM = 86  
moles of hexane =  $9.75/86$   
= 0.1134  
moles of X obtained = 0.01134  
RFM of X =  $2.54/0.01134 = 223.986$   
= 224  
(error [-1]) [3]
- (iii)  $\text{C}_6\text{H}_{10}\text{Cl}_4$  [1]
- (iv) substitution can replace any H atom/multiple substitution can occur [1]

AVAILABLE  
MARKS

22

3 (a) repeated boiling and condensing of a reaction mixture	[1]	<b>AVAILABLE MARKS</b>
(b) add anhydrous magnesium sulfate/calcium sulfate/calcium chloride to the distillate until clear/no longer cloudy filter off drying agent/decant off liquid (error [-1])	[2]	
<p>(c) mass of cyclohexanol = <math>0.96 \times 20 = 19.2</math> g  moles of cyclohexanol = <math>19.2/100 = 0.192</math>  expected moles of cyclohexanone = <math>0.192</math></p> <p>expected mass of cyclohexanone = <math>0.192 \times 98 = 18.82</math> g  actual mass of cyclohexanone = <math>15 \times 0.95 = 14.25</math> g  actual moles of cyclohexanone = <math>14.25/98 = 0.145</math>  % yield = <math>14.25/18.82 \times 100 = 75.7\% = 76\%</math></p> <p><b>or</b></p> <p>% yield = <math>0.145/0.192 \times 100 = 75.5\% = 76\%</math>  (error [-1])</p> <p><b>or</b></p> <p>% yield = <math>\frac{15 \times 0.95}{0.192 \times 98} \times 100 = 75.733</math></p>	[3]	
(d) cyclohexanol is a secondary alcohol/not a primary alcohol/ cyclohexanone will not be oxidised to the carboxylic acid (with this oxidising agent)	[1]	7
<b>Total</b>		<b>55</b>