



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
2011**

Centre Number

71

Candidate Number

Biology

Assessment Unit AS 1

assessing

Molecules and Cells

[AB111]



MONDAY 13 JUNE, AFTERNOON

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Write your answers in the spaces provided in this question paper.
Answer **all nine** questions.

You are provided with **Photograph 1.4** for use with Question 4 in this paper.

Do not write your answers on this photograph.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75.

Section A carries 60 marks. Section B carries 15 marks.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You are reminded of the need for good English and clear

presentation in your answers.

Use accurate scientific terminology in all answers.

You should spend approximately **20 minutes** on Section B.

You are expected to answer Section B in continuous prose.

Quality of written communication will be assessed in **Section B**, and awarded a maximum of 2 marks.

For Examiner's use only

Question Number	Marks
1	
2	
3	
4	
5	
6	
7	
8	
9	

**Total
Marks**

Identify the stage in each case.

- Bivalents are formed when homologous chromosomes pair

- Chromatids separate and are pulled to opposite poles


- Four haploid nuclei are formed

- Chiasmata occur

[4]

Examiner Only	
Marks	Remark

- 2 Fungi are composed of eukaryotic cells, similar to both plant and animal cells. The cellular structure of a fungus is represented in the diagram.



(a) Identify which structures labelled in the diagram above are also found

- in both plant and animal cells

- in plant cells but not in animal cells

- in animal cells but not in plant cells

[3]

(b) Identify **one** feature which is unique to the cells of fungi.


[1]

Examiner Only

Marks Remark

- 3 Using a series of tests, a key was produced to identify the following carbohydrates:

cellulose, glucose, maltose, starch and sucrose.



Examiner Only	
Marks	Remark

Using the key identify each of the carbohydrates **A** to **E**.

A _____

B _____

C _____

D _____

E _____

[5]

Examiner Only	
Marks	Remark

(a) Identify the structures labelled **A** to **E**.

A _____

B _____

C _____

D _____

E _____

[5]


(b) Suggest an interpretation for the area labelled **F**.

[1]

(c) Describe the role of the lacteal within each villus of the ileum.

[1]

Examiner Only	
Marks	Remark



[1]

- (b)** In a classic experiment, Matthew Meselson and Frank Stahl grew bacteria in a medium where the nitrogen source contained the 'light' nitrogen isotope, ^{14}N . The bacteria were then placed in a medium with a nitrogen source containing the 'heavy' nitrogen isotope, ^{15}N , and allowed to reproduce. After many generations, all DNA in the bacteria contained this 'heavy' ^{15}N isotope. This DNA was termed 'heavy' DNA.

DNA extracted from bacterial cells was centrifuged and observed under ultra-violet light. The DNA appeared as a black band in the centrifuge tube. The band produced by 'heavy' DNA was much lower in the centrifuge tube than that produced by 'light' DNA. They are shown in the diagrams below.



The bacteria containing 'heavy' DNA were then transferred to a medium with a nitrogen source containing the 'light' nitrogen isotope, ^{14}N and allowed to reproduce. After one generation, samples of the bacteria were removed and their DNA was extracted and centrifuged. This process was repeated after a further generation.

- (i) Complete the diagrams below to show the position of the extracted DNA by **drawing appropriate bands**.

after one
generation



after two
generations



[2]

- (ii) Explain the result produced after one generation.

[2]


Examiner Only	
Marks	Remark

6 Enzymes can be immobilised for use in biotechnology.

(a) Outline **two** methods used to immobilise enzymes.

1. _____
- _____
2. _____
- _____ [2]

(b) The graph below shows the effect of pH on the enzyme sucrase when in solution and when immobilised.



(i) The graph shows that enzyme immobilisation reduces the activity of sucrase at the optimum pH. Calculate the percentage reduction in the rate of the activity for this immobilised sucrase.

Answer _____ [3]

Monosaccharide	Rate of absorption/arbitrary units	
	Living intestinal cells	Cyanide-treated intestinal cells
Glucose	1.00	0.33
Arabinose	0.29	0.29

(a) Describe the trends evident in the data above.

[3]

(b) Identify which of the two monosaccharides is entirely absorbed by diffusion. Explain your answer.

[3]

Examiner Only	
Marks	Remark


[4]

-
-
-
-
- [2]

Examiner Only	
Marks	Remark

[1]

[1]



Examiner Only	
Marks	Remark

[2]

- [13]

[illegible]

Examiner Only	
Marks	Remark

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Examiner Only	
Marks	Remark

THIS IS THE END OF THE QUESTION PAPER

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.

GCE Biology Advanced Subsidiary (AS)
Assessment Unit AS 1: Molecules and Cells
Summer 2011

Photograph 1.4
(For use with Question 4)



© School of Anatomy and Human Biology, The University of Western Australia