



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
January 2011**

Chemistry

Assessment Unit AS 2

assessing

**Module 2: Organic, Physical
and Inorganic Chemistry**

[AC121]

THURSDAY 20 JANUARY, AFTERNOON

**MARK
SCHEME**

Section A

- 1 B
- 2 B
- 3 B
- 4 D
- 5 C
- 6 C
- 7 D
- 8 A
- 9 B
- 10 B

- [2]
- [2]
- [2]
- [2]
- [2]
- [2]
- [2]
- [2]
- [2]
- [2]

AVAILABLE
MARKS


20

Section B

11 (a) (i) Endothermic

[1]

(ii)



Reaction path/reaction co-ordinate/extent of reaction/progress

([−1] for each mistake)

[3]

(iii) Moles of $N_2 = 50/24 = 2.083$ mole
 Moles of $NaN_3 = 2.083 \times 2/3 = 1.389$ mole
 Mass of $NaN_3 = 1.389 \times 65 = 90.28$ g
 ([−1] for each mistake)

[3]

(b) (i) $C_7H_{16} + 11O_2 \rightarrow 7CO_2 + 8H_2O$

[2]

(ii) $2CO + 2NO \rightarrow 2CO_2 + N_2$

[2]

11

		AVAILABLE MARKS
12 (a)	The outer electrons are in the s-subshell/orbital/shell	[1]
(b) (i)	$\text{Mg(g)} \rightarrow \text{Mg}^{\text{+}}(\text{g}) + \text{e}^{-}$ (equation [1], state symbols [1])	[2]
(ii)	1st ionisation energy decreases going down the group [1] Increased shielding (from inner electrons) [1] Outer electrons further from the nucleus [1]	[3]
(c) (i)	$\text{Ca(OH)}_2 \rightarrow \text{CaO} + \text{H}_2\text{O}$	[1]
(ii)	Mg(OH)_2 less stable than Ba(OH)_2 [1] Mg^{2+} smaller/high charge density [1] Polarisation of OH^{-} /Comparison in size to $\text{O}^{2-}/\text{OH}^{-}$ [1]	[3]
(iii)	pH of $\text{Ba(OH)}_2 > \text{Mg(OH)}_2$ [1] Solubility of the hydroxides increases going down the group [1]	[2]
13 (a)	Fractional distillation	[1]
(b) (i)	Hydrocarbon – compound containing carbon and hydrogen only [1] Saturated – contains no C=C bonds [1]	[2]
(ii)	2,2,3,3-tetramethylbutane	[1]
(c) (i)	Cracking	[1]
(ii)	High temperature/catalyst	[1]
(iii)	$\text{C}_8\text{H}_{18} \rightarrow \text{C}_5\text{H}_{12} + \text{C}_3\text{H}_6$ or $\text{C}_8\text{H}_{18} \rightarrow \text{C}_2\text{H}_6 + 2\text{C}_3\text{H}_6$	[1]
(iv)	Shake propene gas with bromine water Decolourises [1] bromine water [1]	[2]
(v)	$\text{C}_3\text{H}_6 + \text{Br}_2 \rightarrow \text{C}_3\text{H}_6\text{Br}_2$	[1]
(d)	High electron density around the double bond	[1]
(e) (i)	$\begin{array}{c} \diagup \quad \diagdown \\ \text{C}=\text{C} \\ \diagdown \quad \diagup \end{array} + \text{HBr} \rightarrow \begin{array}{c} \text{H} \quad + \\ \quad \\ -\text{C}-\text{C}- \\ \quad \end{array} + \text{Br}^{-}$ \downarrow $\begin{array}{c} \text{H} \quad \text{Br} \\ \quad \\ -\text{C}-\text{C}- \\ \quad \end{array}$	[3]
(ii)	Electrophilic [1] addition [1]	[2]

12

16

			AVAILABLE MARKS
14 (a) (i)	$\Delta H = mc\Delta T = 100 \times 4.2 \times 0.9$ [1] $= 378 \text{ J}$ [1]	[2]	
(ii)	$5/80 = 0.0625$	[1]	
(iii)	$378/0.0625 = 6048 \text{ (J mol}^{-1}\text{)}$	[1]	
(b)	5g produces 0.9°C change $\frac{5}{0.9} \times 25 \text{ g}$ produces 25°C change $\frac{5}{0.9} \times 25 \times \frac{120}{100}$ in 120g water = 166.7g	[2]	6
15 (a)	Rate of reaction increases as more molecules will have enough energy to react when they collide [1] (Equilibrium moves to the right) increasing the yield of NO [1] The forward reaction is endothermic/removes the extra heat [1]	[3]	
(b)	Rate of the reaction will increase due to increased concentration [1] Equilibrium moves to the right increasing the yield of NO [1] This removes the additional N_2 (restoring the equilibrium) [1]	[3]	
(c)	Rate of the reaction will increase as this increases the concentration of the gases [1] Equilibrium position is unaffected [1] Same volume/number of molecules on each side [1]	[3]	
(d)	Rate of the reaction increases as the catalyst provides an alternative pathway with a lower energy [1] The catalyst will not affect the position of the equilibrium and the yield will not change [1]	[2]	
16 (a) (i)	Molecular vibrations	[1]	
(ii)	Spectrum A: C—H bond [1] Spectrum C: C—O bond [1]	[2]	
(iii)	Spectrum B: ethanoic acid [1] Spectrum C: ethanal [1]	[2]	
(b) (i)	$\text{H}^+/\text{Cr}_2\text{O}_7^{2-}$ acidified dichromate	[1]	
(ii)	Cr^{3+}	[1]	
(iii)	Ethanal: Heat under distillation [1] Ethanoic acid: Heat under reflux [1]	[2]	
(c)	$\text{Na} \rightarrow \text{CH}_3\text{CH}_2\text{ONa}$ [1] $\text{CH}_3\text{COCl} \rightarrow \text{CH}_3\text{COOCH}_2\text{CH}_3$ [1] $\text{SOCl}_2 \rightarrow \text{CH}_3\text{CH}_2\text{Cl}$ [1] $\text{PCl}_5 \rightarrow \text{CH}_3\text{CH}_2\text{Cl}$ [1]	[4]	13

17 (a) (i) HBr	[1]	AVAILABLE MARKS
(ii) Nucleophilic [1] substitution [1]	[2]	
(iii) Heat [1] with (concentrated) ammonia [1]	[2]	
(b) Use of separating funnel [1] Use of NaHCO ₃ solution (to remove acid impurities) – release pressure [1] Use of anhydrous Na ₂ SO ₄ or CaCl ₂ (for drying) – filter [1] Distil (product for final purification) [1]	[4]	
Quality of written communication	[2]	
Section B		80
Total		100