



Rewarding Learning

ADVANCED  
General Certificate of Education  
2011

Centre Number

71

Candidate Number

## Chemistry

### Assessment Unit A2 2

*assessing*

Analytical, Transition Metals, Electrochemistry  
and Further Organic Chemistry

[AC222]



FRIDAY 27 MAY, AFTERNOON

#### TIME

2 hours.

#### INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Answer **all fifteen** questions.

Answer **all ten** questions in **Section A**. Record your answers by marking the appropriate letter on the answer sheet provided. Use only the spaces numbered 1 to 10. Keep in sequence when answering.

Answer **all five** questions in **Section B**. Write your answers in the spaces provided in this question paper.

#### INFORMATION FOR CANDIDATES

The total mark for this paper is 120.

Quality of written communication will be assessed in question **13(d)(iii)**.

In Section A all questions carry equal marks, i.e. **two** marks for each question.

In Section B the figures printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A Periodic Table of Elements (including some data) is provided.

For Examiner's use only	
Question Number	Marks
Section A	
1–10	
Section B	
11	
12	
13	
14	
15	
<b>Total Marks</b>	

## Section A

For each of the questions only **one** of the lettered responses (A–D) is correct.

Select the correct response in each case and mark its code letter by connecting the dots as illustrated on the answer sheet.

- 1 Chlorine has two isotopes, chlorine-35 and chlorine-37. Which one of the following is the number of peaks found in the mass spectrum of chlorine gas?
- A 2  
B 3  
C 4  
D 5
- 2 The mechanism for the nitration of benzene is described as
- A electrophilic addition.  
B electrophilic substitution.  
C nucleophilic addition.  
D nucleophilic substitution.
- 3 Copper(II) ions form a coloured complex with the ligand L. The following absorbances were recorded on mixing different volumes of 0.05 M copper(II) sulfate and 0.1 M ligand L.

volume of 0.05 M $\text{CuSO}_4(\text{aq})/\text{cm}^3$	volume of 0.1 M $\text{L}(\text{aq})/\text{cm}^3$	absorbance
3.0	7.0	0.412
4.0	6.0	0.457
5.0	5.0	0.406
6.0	4.0	0.335
7.0	3.0	0.251

Which one of the following is the cation to ligand ratio in this complex?

- A 1:2  
B 1:3  
C 2:3  
D 3:2

Examiner Only

Marks Remark

4 In which one of the following is a metal in the +2 state?

- A  $[\text{Ag}(\text{NH}_3)_2]^+$   
 B  $[\text{Co}(\text{NH}_3)_5\text{Br}]^{2+}$   
 C  $\text{K}_4\text{Fe}(\text{CN})_6$   
 D  $\text{K}_2\text{Cr}_2\text{O}_7$

5 An organic compound X contains carbon, hydrogen and oxygen only. When 1.29g of X is burnt completely, 3.30g of carbon dioxide and 1.35g of water are formed. Which one of the following is the empirical formula of X?

- A  $\text{CH}_2\text{O}$   
 B  $\text{C}_2\text{H}_6\text{O}$   
 C  $\text{C}_4\text{H}_8\text{O}$   
 D  $\text{C}_5\text{H}_{10}\text{O}$

6 Which one of the following is the relative molecular mass of 2,4-dichloro-3,5-dimethylphenol?

- A 190  
 B 191  
 C 192  
 D 196

7 Given the following standard electrode potentials:

	$E^\ominus/\text{V}$
$\text{V}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{V}^{2+}(\text{aq})$	-0.26
$\text{SO}_4^{2-}(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{SO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.17
$\text{VO}^{2+}(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{V}^{3+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.34
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00

Which one of the following reagents will convert  $\text{V}^{3+}(\text{aq})$  to  $\text{VO}^{2+}(\text{aq})$ ?

- A Aqueous iron(II) ions  
 B Aqueous iron(III) ions  
 C Aqueous sulfate ions in acidic solution  
 D Aqueous sulfite ions

Examiner Only	
Marks	Remark



## Section B

Answer **all five** questions in the spaces provided.

- 11 Lysine (2,6-diaminohexanoic acid) has the formula



The molecule is optically active and may undergo polymerisation.

- (a) (i) Explain the term **optically active**.

\_\_\_\_\_ [2]

- (ii) Draw the 3D structure of lysine labelling the asymmetric carbon with an asterisk (\*).

[2]

- (b) Explain why lysine has a relatively high melting point.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [2]

- (c) Write the formula of the organic ion present when lysine is dissolved in an alkaline solution.

\_\_\_\_\_ [1]

Examiner Only	
Marks	Remark

- (d) Draw the structure of a dimer formed when two molecules of lysine react.

[2]

- (e) A mixture of amino acids may be separated using paper chromatography.

- (i) Explain the term  **$R_f$  value** as applied to paper chromatography.

\_\_\_\_\_ [1]

- (ii) Explain, in terms of partition, what a low  $R_f$  value indicates about a particular amino acid.

\_\_\_\_\_  
 \_\_\_\_\_ [2]

- (f) An amino acid was found to have the following composition by mass:

element	% composition
N	10.5
C	36.1
H	5.3
O	48.1

Deduce the empirical formula for this amino acid.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [3]

Examiner Only


Marks Remark

12 Chromium is a transition metal which was discovered by Louis Vauquelin in 1797. He was able to detect traces of chromium in precious gems, such as ruby and emerald.

- (a) State the electronic structure of a chromium atom and explain why the arrangement is stable.

\_\_\_\_\_  
\_\_\_\_\_ [2]

- (b) Chromium(III) oxide,  $\text{Cr}_2\text{O}_3$ , can be reduced to the metal by heating with aluminium powder according to the equation:



Calculate the percentage yield when 42.5g of chromium are obtained from a reaction between 25g of aluminium and 100g of chromium(III) oxide.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_ [3]

- (c) Chromium trioxide,  $\text{CrO}_3$ , is formed by the reaction between excess concentrated sulfuric acid and a concentrated solution of potassium dichromate. Potassium sulfate and water are formed as by-products.


Write the equation for this reaction.

\_\_\_\_\_ [2]

Examiner Only

Marks Remark

(d) A data book lists the following standard electrode potentials:



(i)  $\text{Cr}^{2+}$  can reduce silver ions to silver atoms. Write the equation for this reduction.

\_\_\_\_\_ [1]

(ii) Deduce the electrode potential for this change.

\_\_\_\_\_ [1]

(e) Chrome alum is a double salt prepared by the reduction of potassium dichromate,  $\text{K}_2\text{Cr}_2\text{O}_7$ , using a suitable reducing agent.


(i) Name a suitable reducing agent.

\_\_\_\_\_ [1]

(ii) State the colour and formula of crystalline chrome alum.

\_\_\_\_\_  
\_\_\_\_\_ [2]

(f) Chrome alum may be used as a mordant in dyeing, binding the dye molecules to the fibre. The dye Prontosil is prepared by the following sequence:



(i) Deduce the empirical formula for 4-aminobenzenesulfonamide.

\_\_\_\_\_ [1]

Examiner Only	
Marks	Remark

(ii) Name the reagents required for step A.

\_\_\_\_\_ [1]

(iii) State and explain the condition necessary for the first step.

\_\_\_\_\_  
\_\_\_\_\_ [2]

(iv) Suggest a structure of the dye Prontosil which is formed by a coupling reaction.

[2]

(v) Explain why Prontosil is coloured.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_ [3]

Examiner Only


Marks Remark

13 The first person to extract ethanoic acid from vinegar was the alchemist Jabir ibn Hayyan Geber (c. 721–815 AD). However the pure compound was not produced for another ten centuries.

(a) The mass spectrum of ethanoic acid shows a distinct peak at  $m/e = 59$ . State the formula of the species giving rise to this peak.

\_\_\_\_\_ [1]

(b) Ethanoic acid reacts with ethanol to form the ester ethyl ethanoate,  $\text{CH}_3\text{COOC}_2\text{H}_5$ . The n.m.r. spectrum of ethyl ethanoate consists of three sets of peaks as shown below.



(i) Explain the peak integrations.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [2]

(ii) Explain the chemical shifts.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [2]

Examiner Only	
Marks	Remark

(iii) Explain the splitting pattern **a**.

\_\_\_\_\_ [1]  
\_\_\_\_\_

(iv) Explain why **b** is a singlet.

\_\_\_\_\_ [1]  
\_\_\_\_\_

(c) Ethanoic acid can be converted to ethanamide via the thermal decomposition of the ammonium salt.

(i) Write the equation for the reaction of ethanoic acid with ammonia.

\_\_\_\_\_ [1]

(ii) Write the equation for the thermal decomposition.

\_\_\_\_\_ [1]


(iii) Ethanamide is hydrolysed by either hydrochloric acid or sodium hydroxide to form different organic products. State the formula of the organic product in each case.

hydrochloric acid \_\_\_\_\_

sodium hydroxide \_\_\_\_\_ [2]

Examiner Only	
Marks	Remark

(d) Ethanamide can be converted to an amine in the following sequence.



(i) Give the formula of reagents A and B.

A \_\_\_\_\_

B \_\_\_\_\_ [2]

(ii) Write the equation for the reaction of an unknown amine,  $\text{RNH}_2$ , with ethanoyl chloride.

\_\_\_\_\_ [1]

(iii) Describe how you would identify the unknown amine using the pure N-substituted amide. Include relevant practical steps.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_ [4]

Quality of written communication. [2]

(e) 1,2-diaminoethane (en) is a bidentate ligand forming stable complex ions with transition metal ions.

(i) Explain the term **bidentate**.

\_\_\_\_\_

\_\_\_\_\_ [2]

Examiner Only	
Marks	Remark

- (ii) Hexaaquanickel(II) ions react with en in solution. Write the equation for this reaction in which all the water ligands are replaced.

\_\_\_\_\_ [2]

- (iii) Explain why this ligand replacement takes place.

\_\_\_\_\_  
\_\_\_\_\_ [2]

Examiner Only	
Marks	Remark

**14** Iron is a transition metal which fulfils vital biological, industrial and chemical roles.

- (a) The most common oxidation states of iron in its compounds are +2 and +3. State the electronic structures of the  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  ions.

$\text{Fe}^{2+}$  \_\_\_\_\_

$\text{Fe}^{3+}$  \_\_\_\_\_ [2]

- (b) Metallic iron is used as a heterogeneous catalyst in the Haber Process to make ammonia.



Explain in terms of chemisorption how the reaction takes place.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_ [3]

- (c) Iron filings are used to catalyse the monohalogenation of aromatic compounds such as methylbenzene,  $\text{C}_6\text{H}_5\text{CH}_3$ .

- (i) Draw a flow scheme for the mechanism of the monobromination of methylbenzene to form 4-bromomethylbenzene.

[3]

- (ii) Suggest the name of the product formed in the monobromination of 1,4-dimethylbenzene.

\_\_\_\_\_ [2]

Examiner Only	
Marks	Remark

(d) (i) Explain the role of iron(II) in haemoglobin.

\_\_\_\_\_  
 \_\_\_\_\_ [2]



(ii) Explain the effect of the inhalation of carbon monoxide on haemoglobin.

\_\_\_\_\_ [1]

(e) Aqueous iron(III) ions form a stable complex with the bidentate ligand ethanedioate,  $C_2O_4^{2-}$ .  
 The iron(III) ions combine with three ethanedioate ions. Deduce the formula of the complex formed.

\_\_\_\_\_ [1]

(f) Potassium manganate(VII) oxidises iron(II) ions and ethanedioate ions according to the equations:



(i) Describe, with observations, how you could confirm the presence of aqueous iron(III) ions, following the oxidation of iron(II) ions, without interference from manganese(II) ions.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [3]

(ii)  $25.0\text{ cm}^3$  of an acidified iron(II) ethanedioate solution required  $32.2\text{ cm}^3$  of  $0.025\text{ mol dm}^{-3}$  of potassium manganate(VII) solution for complete reaction. Calculate the concentration, in  $\text{mol dm}^{-3}$ , of the iron(II) ethanedioate solution.

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_ [4]

Examiner Only	
Marks	Remark

15 Polymers have become invaluable materials with many familiar names e.g. polythene, Perspex, and nylon.

(a) The manufacture of High Density (HD) polythene was first developed by Ziegler in the 1950s.

(i) State the conditions used to manufacture HD polythene.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_ [3]

(ii) State and explain the flexibility and softening temperature of HD polythene.


\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_ [3]

(b) Perspex is an addition polymer made by polymerisation of methyl 2-methylpropenoate monomer shown below.



Draw a section of the Perspex polymer showing at least two repeating units.

[2]

Examiner Only	
Marks	Remark



---

**THIS IS THE END OF THE QUESTION PAPER**

---



Permission to reproduce all copyright material has been applied for.  
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA  
will be happy to rectify any omissions of acknowledgement in future if notified.