



Rewarding Learning

ADVANCED
General Certificate of Education
2011

Chemistry

Assessment Unit A2 1

assessing

Periodic Trends and Further Organic,
Physical and Inorganic Chemistry

[AC212]


MONDAY 23 MAY, AFTERNOON

**MARK
SCHEME**

Section B

AVAILABLE
MARKS

11 (a) (i)



$$(ii) K_c = \frac{[\text{CH}_3\text{COOCH}_2\text{CH}_2\text{OOCCH}_3][\text{H}_2\text{O}]^2}{[\text{CH}_3\text{COOH}]^2[\text{HOCH}_2\text{CH}_2\text{OH}]}$$

expression [1]

No units [1]


[2]



[2]

6

12 (a) (i)





[1] per box (order can vary) [4]

$$(ii) 2(\text{E.A.}) = (-121 \times 2) + (-1450) + (-736) + (-150) + (-642) + (2493)$$

$$\text{E.A.} = -727 \div 2 = -363.5 \text{ kJ mol}^{-1}$$

[1]

[1]


		AVAILABLE MARKS
(b)	$1s^2 2s^2 2p^6$ $1s^2 2s^2 2p^6 3s^2 3p^6$	[1] [1]
(c)	The enthalpy change which occurs when one mole of an (ionic) compound dissolves in water	[2] 10
13 (a)		
(i)	$C_nH_{2n}O$	[1]
(ii)	3-methylbutanal	[1]
(iii)	$CH_3CH_2CH_2COCH_3$ or $CH_3CH_2COCH_2CH_3$ or $(CH_3)_2CHCOCH_3$ correct name for each structure pentan-2-one, pentan-3-one or (3-)methylbutan(-2-)one	[2] [2]
(iv)	suitable reagent heat observation for ketone observation for aldehyde Quality of written communication	[4] [2]
(v)	e.g. $CH_3CH = C(OH)CH_2CH_3$ and many others	[1]
(vi)	 e.g. and labels	[3]
(b) (i)	yellow/orange [1] solid [1]	[2]
(ii)		[3]
(iii)	melting point determination [1] match (with melting point of butanone-2,4-dinitrophenylhydrazone) using tables of data [1]	[2]
(iv)	$CH_3COCH_2CH_3 + 2[H] \rightarrow CH_3CH(OH)CH_2CH_3$ Butan-2-ol	[2] [1]
		26

- 14 (a) (i) fully dissociates [1]
- (ii) H^+ donor [1]
- (iii) $pH = -\log_{10}[H^+]$ [1]
- (iv) $[H^+] = 0.4 \text{ mol dm}^{-3}$
 $pH = 0.40$ [2]
- (v) $K_w = [OH^-][H^+]$ [1]
- (vi) $[OH^-] = 0.2 \text{ mol dm}^{-3}$ $[H^+] = 5.0 \times 10^{-14} [1]$
 $pH = 13.30 [1]$ [2]
- (vii) $2NaOH + H_2SO_4 \rightarrow Na_2SO_4 + 2H_2O$ [2]
- (b) (i) $K_a = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$ [1]
- (ii) $[H^+]^2 = 3.48 \times 10^{-6}$
 $[H^+] = 1.87 \times 10^{-3}$
 $pH = 2.73$ [3]
- (c) (i) $CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O$ [1]
- (ii) initial moles of ethanoic acid = 5.0×10^{-3}
moles of sodium hydroxide added = 3.0×10^{-3}
moles of sodium ethanoate formed = 3.0×10^{-3}
moles of ethanoic acid left = 2.0×10^{-3}
 $[H^+] = 1.74 \times 10^{-5} \times \frac{2.0 \times 10^{-3}}{3.0 \times 10^{-3}} = 1.16 \times 10^{-5} \text{ mol dm}^{-3}$
 $pH = 4.94$ [4]
- (iii) sodium ethanoate gives ethanoate ions
ethanoate ions combine with H^+ ions or equation * essential
 $[H^+]/pH$ remains approximately constant [2]
- (iv) alkaline [1] – the salt of a weak acid and strong base [1] [2]

AVAILABLE
MARKS

23

15 (a) (i)



[2]

(ii) propane-1,2,3-triol [1]

(iii) The mass of KOH [1] in mg [1] required to completely hydrolyse/saponify 1.0g of a fat. [1] [3]

(iv) RMM of fat = 890
 moles of fat = 0.00112
 moles of KOH = 0.00337
 mass of KOH (in g) = 0.189
 mass of KOH (in mg)/S.V. = 189 [4]


(b) (i) $\text{C}_{19}\text{H}_{38}\text{O}_2$ [1](ii) $2\text{C}_{19}\text{H}_{38}\text{O}_2 + 55\text{O}_2 \rightarrow 38\text{CO}_2 + 38\text{H}_2\text{O}$ [2]

(iii) photosynthesis/respiration/solubility in surface waters/decay [2]


AVAILABLE
MARKS

15

- 16 (a) (i) First [1] valid explanation [1] [2]
- (ii) Zero [1] valid explanation [1] [2]
- (iii) Rate = $k [C_4H_9Br]$ [1] [1]
- (iv) 2000 [1] s^{-1} [1] [2]
- (v) tertiary structure [1]
 correct RDS [1]
 first step producing correct carbocation and bromide ion [1]
 second step involving hydroxide attacking carbocation [1] [4]



- (b) (i) carbon with 4 different atoms/groups attached [1]
- (ii) non-superimposable [1] mirror images [1] [2]
- (iii) structure [1] butan-2-ol [1] [2]
- (iv)



- (v) plane-polarised light [1] rotated in opposite directions [1] [2]

Section B

100

Total

120