



Rewarding Learning

ADVANCED
General Certificate of Education
2015

Chemistry

Assessment Unit A2 1

assessing

Periodic Trends and Further Organic,
Physical and Inorganic Chemistry

[AC212]

FRIDAY 22 MAY, MORNING

**MARK
SCHEME**

General Marking Instructions

Introduction

Mark schemes are published to assist teachers and students in their preparation for examinations. Through the mark schemes teachers and students will be able to see what the examiners are looking for in response to questions and exactly where the marks have been awarded. The publishing of the mark schemes may help to show that examiners are not concerned about finding out what a student does not know but rather with rewarding students for what they do know.

The purpose of mark schemes

Examination papers are set and revised by teams of examiners and revisers appointed by the Council. The teams of examiners and revisers include experienced teachers who are familiar with the level and standards expected of students in schools and colleges.

The job of the examiners is to set the questions and the mark schemes; and the job of the revisers is to review the questions and mark schemes commenting on a large range of issues about which they must be satisfied before the question papers and mark schemes are finalised.

The questions and the mark schemes are developed in association with each other so that the issues of differentiation and positive achievement can be addressed right from the start. Mark schemes, therefore, are regarded as part of an integral process which begins with the setting of questions and ends with the marking of the examination.

The main purpose of the mark scheme is to provide a uniform basis for the marking process so that all the markers are following exactly the same instructions and making the same judgements in so far as this is possible. Before marking begins a standardising meeting is held where all the markers are briefed using the mark scheme and samples of the students' work in the form of scripts. Consideration is also given at this stage to any comments on the operational papers received from teachers and their organisations. During this meeting, and up to and including the end of the marking, there is provision for amendments to be made to the mark scheme. What is published represents the final form of the mark scheme.

It is important to recognise that in some cases there may well be other correct responses which are equally acceptable to those published: the mark scheme can only cover those responses which emerged in the examination. There may also be instances where certain judgements may have to be left to the experience of the examiner, for example where there is no absolute correct response – all teachers will be familiar with making such judgements.

Section A

- 1 B
- 2 D
- 3 B
- 4 C
- 5 D
- 6 C
- 7 B
- 8 D
- 9 D
- 10 D

[2] for each correct answer

[20]
Section A

**AVAILABLE
MARKS**

20

20

Section B

AVAILABLE
MARKS

- 11 (a) iodine
low bond energy [1]
- (b) (i) energy is needed to separate the positive ions from the
negative ions/(ionic) bonds being broken [1]
- (ii) the size of the halide ions increase/the bond energies of the
halogens get less [1]
- (c) (i) $+178 + 590 + 1145 + 4912 + 3 \times 112 - 3 \times 349 = \Delta H + 4650$
 $\Delta H = +1464 \text{ kJ mol}^{-1}$ [3]
- (ii) the reaction is (highly) endothermic [1]
- (iii) size of charge on Ca ion increases, (hence greater attractive forces) [1]
- (iv) $2\text{CaCl} \rightarrow \text{Cl}_2 + 2\text{Ca} \quad + 2 \times 178 = +356$
 $\text{Ca} + \text{Cl}_2 \rightarrow \text{CaCl}_2 \quad -796$
 $2\text{CaCl} \rightarrow \text{Ca} + \text{CaCl}_2 \quad -796 + 356 = -440 \text{ kJ/kJ mol}^{-1}$ [2]
- (v) all the products and reactants are solids/very small change
in entropy/no entropy change [1]
- 12 (a) 2-methylpropenoic acid [2]
- (b) (i) $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{O} \\ \diagup \\ \text{CH}_3 \end{array} + \text{CN}^- \rightarrow \begin{array}{c} \text{CH}_3 \quad \text{O}^- \\ \diagdown \quad \diagup \\ \text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CN} \end{array} \xrightarrow{\text{H}^+} \begin{array}{c} \text{CH}_3 \quad \text{OH} \\ \diagdown \quad \diagup \\ \text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CN} \end{array}$ [3]
- (ii) sodium hydroxide(aq)/potassium hydroxide(aq) [1]
(dil.) hydrochloric acid/(dil.) sulfuric acid [1] [2]
- (iii) methanol [1]
concentrated/conc. sulfuric acid [1] [2]
- (iv) esterification [1]
- (c) (i) petroleum/North Sea gas [1]
Haber process [1] [2]
- (ii) $3\text{N}-\text{H} + 4\text{C}-\text{H} = 3 \times 391 + 4 \times 413 = 1173 + 1652 = +2825$
 $1\text{C}-\text{H} + 1\text{C}\equiv\text{N} + 3\text{H}-\text{H} = 1 \times 413 + 1 \times 887 + 3 \times 436 = -2608$
 $= +217 \text{ kJ/kJ mol}^{-1}$ [3]
- (iii) increase in the number of molecules/moles [1]
 $\Delta G = \Delta H - T\Delta S = 217 \times 10^3 - T \times 125$
- $T = 217000/125 = 1736 \text{ K}$ [2] [3]

11

			AVAILABLE MARKS
(d) (i)	landfill sites use up empty quarries		
	or landfill sites once refilled are used	[1]	
	waste incineration creates heat/power		
	or incineration reduces waste	[1]	[2]
(ii)	landfill sites are unsightly		
	incineration produces carbon dioxide/global warming		[2]
(iii)	waste can be recycled		
	use biodegradable products		
	reduce the amount of packaging	[2]	24
13 (a)	H_2O_2 and HO_2^- acid and base	[1]	
	H_2O and H_3O^+ base and acid	[1]	[2]
(b)	Adding water moves the equilibrium to the RHS [1] increases H^+ (decreases pH) [1]		[2]
(c) (i)	measure the volume of oxygen produced		
	draw a graph of volume against time		
	measure the slope of the graph (to determine rate)/calculate gradient		
	graph of rate versus concentration (of hydrogen peroxide)	[4]	
(ii)	step 1 is slow/rate determining	[1]	
	step 2 is fast	[1]	[2]
	or step 1 is slower than step 2		
(iii)	OBr^-	[1]	
(iv)	second	[1]	
(v)	a catalyst participates in a reaction but is not used up	[1]	
(vi)	add silver nitrate solution	[1]	
	filter off the silver bromide	[1]	
	weigh the silver bromide, (relate mass of AgBr to Br^-)	[1]	[3]
(d)	$X = 0$ $Y = 1$ $Z = 2$ (or more)	[2]	18

- 14 (a) olive oil/vegetable oil [1]
- (b) dissolve the fat in a suitable solvent/shake [1]
with a solution of iodine/bromine [1]
the colour disappears [1] [3]
- (c) boil/heat the sample with hydrochloric acid [1]
separate the insoluble petroleum oil [1]
weigh/determine volume of the petroleum oil [1]
Any **two** [2]
- (d) 3 mol of NaOH = mol of fat hence 8.33×10^{-4} mol of fat
0.84 g of iodine I = $0.84/127 = 0.0066 = 0.0033$ mol I₂
 $1 \text{ I}_2 = 1 \text{ C}=\text{C}$ $0.0033/8.33 \times 10^{-4} = 3.96/4$ per molecule
 $\frac{3.96}{3} = 1.32$ per fatty acid [3]
- (e) (i) glycerol [1]
- (ii)
$$\begin{array}{c} \text{CH}_2\text{OCOR}_2 \\ | \\ \text{H}^*\text{C OCOR}_2 \\ | \\ \text{CH}_2\text{OCOR}_1 \end{array}$$
 [3]
- (iii) hydrogen (gas) [1]
passed over/through/reacted with oil/180 °C [1]
nickel catalyst [1] [3]
- (f) place the sample in a capillary tube sealed (at one end) [1]
heat slowly [1]
in oil bath/melting point apparatus [1]
note temperature at which sample starts to melt and finishes melting [1]
comment on determining substance as an oil/fat at 25 °C [1] [5]
Quality of written communication [2]

AVAILABLE
MARKS

23

