



Rewarding Learning

ADVANCED SUBSIDIARY (AS)
General Certificate of Education
2017

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Chemistry

Assessment Unit AS 1
assessing
Basic Concepts in Physical
and Inorganic Chemistry

MV18

[SCH12]

FRIDAY 26 MAY, MORNING

Time

1 hour 30 minutes, plus your additional time allowance.

Instructions to Candidates

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Answer **all fifteen** questions.

Answer **all ten** questions in **Section A**. Record your answers by marking the appropriate letter on the answer sheet provided. Use only the spaces numbered 1 to 10. Keep in sequence when answering.

Answer **all five** questions in **Section B**. **You must answer the questions in the spaces provided.**

Complete in black ink only.

Information for Candidates

The total mark for this paper is 90.

Quality of written communication will be assessed in Question **13(c)**.

In Section A all questions carry equal marks, i.e. **one** mark for each question.

In Section B the figures in brackets printed at the end of each question indicate the marks awarded to each question or part question.

A Periodic Table of Elements, containing some data, is included with this question paper.

Section A – Multiple Choice

Select the correct response in each case and mark its code letter by connecting the dots as illustrated on the answer sheet.

Each multiple choice question is worth 1 mark.

1 Bromine is formed in the reaction below.



Which statement about the reaction is correct?

- A Bromide ions lose electrons
- B Bromine is reduced by chlorine
- C Chloride ions are reduced
- D Chlorine is a weaker oxidising agent than bromide

2 Which trend in the Periodic Table is correct?

- A Boiling point decreases from fluorine to bromine
- B First ionisation energy decreases from lithium to caesium
- C First ionisation energy increases from nitrogen to oxygen
- D Melting point decreases from sodium to silicon

3 Which of the following is the structure of $^{55}\text{Mn}^{2+}$?

	protons	neutrons	electrons
A	25	30	23
B	25	30	27
C	27	30	25
D	30	25	28

4 Potassium iodide is formed when potassium is warmed in iodine vapour. Which of the following shows the bonding in the three species?

	potassium	iodine	potassium iodide
A	ionic	covalent	ionic
B	metallic	ionic	covalent
C	covalent	covalent	ionic
D	metallic	covalent	ionic

5 The element astatine lies below iodine in the Periodic Table and is likely to

- A be black.
- B be a volatile liquid at room temperature and pressure.
- C form an astatide ion, At^{2-} .
- D oxidise iodide ions to iodine.

6 Which molecule is non-polar?

A H_2S

B NH_3

C PF_3

D SF_6

7 The element boron has a relative atomic mass of 10.8. In this sample, boron exists as two isotopes, ^{10}B and ^{11}B . The percentage abundance of ^{10}B in this sample of boron is


A 10.8%.

B 20.0%.

C 80.0%.

D 89.2%.

8 When burned in oxygen magnesium forms magnesium oxide.



What is the number of molecules of oxygen required for the complete oxidation of 1.2 g of magnesium?

A 1.5×10^{22}

B 3.0×10^{22}

C 3.0×10^{23}

D 6.0×10^{23}

9 Which statement describes the trends in electronegativity values in the Periodic Table?

A Decrease across a Period and increase down a Group

B Decrease across a Period and decrease down a Group

C Increase across a Period and increase down a Group

D Increase across a Period and decrease down a Group

10 Which of the following would exactly neutralise 10.0 cm³ of 1.00 mol dm⁻³ NaOH(aq)?

- A 2.50 cm³ of 1.00 mol dm⁻³ CH₃COOH
- B 5.00 cm³ of 1.00 mol dm⁻³ HCl
- C 5.00 cm³ of 1.00 mol dm⁻³ H₂SO₄
- D 3.00 cm³ of 1.00 mol dm⁻³ H₃PO₄

Section B

Answer **all five** questions in the spaces provided.

11 Sulfate, hydrogensulfate and thiosulfate ions are formed when sulfuric and thiosulfuric acids ionise.

(a) (i) Write the equation for the complete ionisation of thiosulfuric acid. [2 marks]

(ii) Write the formula for the hydrogensulfate ion. [1 mark]

(b) (i) Write the formula for ammonium sulfate. [1 mark]

(ii) Describe the bonding in ammonium sulfate. [2 marks]

(c) Describe how you could use chemical tests on an aqueous solution of ammonium sulfate to prove that it contains ammonium ions and sulfate ions. [4 marks]

12 Some properties of the metals sodium and aluminium are shown in the table below.

metal	charge on metal ion	electronic structure of the atom	melting point /°C
sodium	1+	$1s^2 2s^2 2p^6 3s^1$	98
aluminium	3+	$1s^2 2s^2 2p^6 3s^2 3p^1$	660

(a) Describe, without using a diagram, the bonding in sodium metal. [2 marks]

(b) Explain why aluminium has a higher melting point than sodium. [2 marks]

(c) (i) Write the equation, including state symbols, for the first ionisation energy of sodium. [2 marks]

(ii) The first six ionisation energies, in kJ mol^{-1} , of sodium are 496, 4563, 6913, 9544, 13352 and 16611. Explain which of these values can be used to identify sodium as belonging to Group I of the Periodic Table. [2 marks]

(iii) The outer electron in the sodium atom is located in the 3s orbital.
Explain what is meant by the term **orbital**.
[2 marks]

(d) Aluminium forms covalent bonds with chlorine.

(i) Explain what is meant by the term **covalent bond**.
[2 marks]

(ii) Write the equation for the reaction of aluminium with chlorine to form aluminium chloride, AlCl_3 .
[1 mark]

(iii) State the octet rule and explain whether the atoms in aluminium chloride obey the rule. [3 marks]

BLANK PAGE

(Questions continue overleaf)

13 (a) Zinc reacts with chlorine to form the ionic compound zinc chloride. Draw a dot and cross diagram, using outer electrons only, to show how zinc chloride, ZnCl_2 , is formed from zinc and chlorine atoms. [2 marks]

(b) Zinc is an essential trace element. People who have a zinc deficiency can take hydrated zinc sulfate, $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$, as a dietary supplement.

The value of x can be determined by heating hydrated zinc sulfate to constant mass.

A student heated 5.65 g of hydrated zinc sulfate and obtained 3.85 g of anhydrous zinc sulfate.

(i) Calculate the number of moles of anhydrous zinc sulfate obtained. [1 mark]

(ii) Calculate the mass of water present in the hydrated zinc sulfate. [1 mark]

(iii) Calculate the number of moles of water present in the hydrated zinc sulfate. [1 mark]

(iv) Calculate the value of x in $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$ [1 mark]

(c) Describe how you would prepare 250.0 cm³ of a 28.7 g dm⁻³ zinc sulfate solution from the anhydrous solid. [6 marks]

In this question you will be assessed on using your written communication skills including the use of specialist scientific terms.

14 Nitrogen and phosphorus are Group V elements. They form the toxic hydrides ammonia and phosphine.

(a) Ammonia is formed by the reversible reaction of nitrogen with hydrogen.

Write the equation for this reaction. [2 marks]

(b) Phosphine is formed by the reaction of phosphorus with aqueous sodium hydroxide.

(i) Balance the equation for the formation of phosphine.
[1 mark]



(ii) Deduce the oxidation number of phosphorus in:




[3 marks]

(iii) Explain, using the oxidation numbers of phosphorus, why the reaction is described as disproportionation. [3 marks]

(c) The boiling point of ammonia is -33°C while that of phosphine is -88°C . Explain why the boiling point of ammonia is higher than that of phosphine. [3 marks]

(d) Both ammonia and phosphine molecules react with H^+ ions.



(i) Name the type of bond formed between a phosphine molecule and the H^+ ion. [1 mark]

(ii) Draw and name the shapes of the molecule PH_3 and the ion PH_4^+ . [4 marks]



Shape _____



Shape _____

(iii) Explain why the bond angle in PH_3 is different from the bond angle in PH_4^+ . [3 marks]

(e) Ammonia is very soluble in water. Draw diagrams to show the two ways in which a molecule of ammonia can be attracted to a molecule of water. Include all partial charges and lone pairs in your diagram. [4 marks]

15 Ammonia is used to make nitric acid by the Ostwald Process outlined below.



(a) (i) Calculate the number of moles of oxygen needed to react with 6.8 kg of ammonia. [3 marks]

(ii) Calculate the number of moles of nitrogen(IV) oxide which can be obtained from 6.8 kg of ammonia. [2 marks]

(iii) Calculate the concentration of nitric acid, in g dm^{-3} , produced on reacting the nitrogen(IV) oxide obtained in part **(ii)** with 50 dm^3 of water. [3 marks]

(b) Ammonia reacts with nitric acid according to the equation below.



The following results were obtained by diluting 25.0 cm³ of a concentrated ammonia solution to 250.0 cm³ in a volumetric flask and then titrating 25.0 cm³ portions of the diluted ammonia solution using 0.100 mol dm⁻³ nitric acid.

titration	initial burette reading/cm ³	final burette reading/cm ³	titre/cm ³
rough	0.00	22.00	22.00
first accurate	0.10	21.40	21.30
second accurate	0.20	21.60	21.40

(i) Name a suitable indicator for the titration and state the colour change at the end point. [3 marks]

(ii) Calculate the mean titre. [1 mark]

(iii) A burette has an uncertainty of $\pm 0.05 \text{ cm}^3$. Calculate the uncertainty when two burette readings are used to calculate a titre value. [1 mark]

(iv) Calculate the concentration of the concentrated ammonia solution in mol dm⁻³. [5 marks]

THIS IS THE END OF THE QUESTION PAPER

For Examiner's use only	
Question Number	Marks
Section A	
1–10	
Section B	
11	
12	
13	
14	
15	
Total Marks	

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.

General Information

1 tonne = 10^6 g

1 metre = 10^9 nm

One mole of any gas at 293 K and a pressure of 1 atmosphere (10^5 Pa) occupies a volume of 24 dm³

Avogadro Constant = 6.02×10^{23} mol⁻¹

Planck Constant = 6.63×10^{-34} Js

Specific Heat Capacity of water = 4.2 J g⁻¹ K⁻¹

Speed of Light = 3×10^8 m s⁻¹

New
Specification

Characteristic absorptions in IR spectroscopy

Wavenumber/cm ⁻¹	Bond	Compound
550–850	C–X (X = Cl, Br, I)	Haloalkanes
750–1100	C–C	Alkanes, alkyl groups
1000–1300	C–O	Alcohols, esters, carboxylic acids
1450–1650	C=C	Arenes
1600–1700	C=C	Alkenes
1650–1800	C=O	Carboxylic acids, esters, aldehydes, ketones, amides, acyl chlorides
2200–2300	C≡N	Nitriles
2500–3200	O–H	Carboxylic acids
2750–2850	C–H	Aldehydes
2850–3000	C–H	Alkanes, alkyl groups, alkenes, arenes
3200–3600	O–H	Alcohols
3300–3500	N–H	Amines, amides

Proton Chemical Shifts in Nuclear Magnetic Resonance Spectroscopy

(relative to TMS)

Chemical Shift	Structure	
0.5–2.0	–CH	Saturated alkanes
0.5–5.5	–OH	Alcohols
1.0–3.0	–NH	Amines
2.0–3.0	–CO–CH	Ketones
	–N–CH	Amines
	C ₆ H ₅ –CH	Arene (aliphatic on ring)
2.0–4.0	X–CH	X = Cl or Br (3.0–4.0) X = I (2.0–3.0)
	–C=CH	Alkenes
4.5–6.0	RCONH	Amides
5.5–8.5	–C ₆ H ₅	Arenes (on ring)
6.0–8.0	–CHO	Aldehydes
9.0–10.0	–COOH	Carboxylic acids
10.0–12.0		

These chemical shifts are concentration and temperature dependent and may be outside the ranges indicated above.

GCE

CHEMISTRY DATA SHEET GCE A/AS EXAMINATIONS CHEMISTRY

Including the Periodic Table of the Elements

For the use of candidates taking
Advanced Subsidiary and Advanced Level
Chemistry Examinations

Copies must be free from notes or additions of any kind.
No other type of data booklet or information sheet is
authorised for use in the examinations.

For first teaching from September 2016
For first award of AS Level in Summer 2017
For first award of A Level in Summer 2018
Subject Code: 1110

THE PERIODIC TABLE OF ELEMENTS

Group

I	II	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 H Hydrogen																	4 He Helium
7 Li Lithium	9 Be Beryllium																2 Ne Neon
23 Na Sodium	24 Mg Magnesium																20 Ar Argon
39 K Potassium	40 Ca Calcium	45 Sc Scandium	48 Ti Titanium	51 V Vanadium	52 Cr Chromium	55 Mn Manganese	56 Fe Iron	59 Co Cobalt	59 Ni Nickel	64 Cu Copper	65 Zn Zinc	70 Ga Gallium	73 Ge Germanium	75 As Arsenic	79 Se Selenium	80 Br Bromine	84 Kr Krypton
19 Rb Rubidium	88 Sr Strontium	89 Y Yttrium	91 Zr Zirconium	93 Nb Niobium	96 Mo Molybdenum	98 Tc Technetium	101 Ru Ruthenium	103 Rh Rhodium	106 Pd Palladium	108 Ag Silver	112 Cd Cadmium	115 In Indium	119 Tl Tin	122 Sb Antimony	128 Te Tellurium	127 I Iodine	131 Xe Xenon
37 Cs Caesium	137 Ba Barium	139 La [*] Lanthanum	178 Hf Hafnium	181 Ta Tantalum	184 W Tungsten	186 Re Rhenium	190 Os Osmium	192 Ir Iridium	195 Pt Platinum	197 Au Gold	201 Hg Mercury	204 Tl Thallium	207 Pb Lead	209 Bi Bismuth	210 Po Polonium	210 At Astatine	222 Rn Radon
55 Fr Francium	226 Ra Radium	227 Ac [†] Actinium	261 Rf Rutherfordium	262 Db Dubnium	266 Sg Seaborgium	264 Bh Bohrium	277 Hs Hassium	268 Mt Meitnerium	271 Ds Darmstadtium	272 Rg Roentgenium	285 Cn Copernicium						

* 58–71 Lanthanum series

† 90–103 Actinium series

$\frac{a}{b}^x$ a = relative atomic mass (approx)
x = atomic symbol
b = atomic number

140 Ce Cerium	141 Pr Praseodymium	144 Nd Neodymium	145 Pm Promethium	150 Sm Samarium	152 Eu Europium	157 Gd Gadolinium	159 Tb Terbium	162 Dy Dysprosium	165 Ho Holmium	167 Er Erbium	169 Tm Thulium	173 Yb Ytterbium	175 Lu Lutetium
58 Th Thorium	90 Pa Protactinium	238 U Uranium	237 Np Neptunium	242 Pu Plutonium	243 Am Americium	247 Cm Curium	245 Bk Berkelium	251 Cf Californium	254 Es Einsteinium	253 Fm Fermium	256 Md Mendelevium	254 No Nobelium	257 Lr Lawrencium