



ADVANCED SUBSIDIARY (AS)
General Certificate of Education
January 2013

Mathematics
Assessment Unit F1
assessing
Module FP1: Further Pure Mathematics 1
[AMF11]

FRIDAY 11 JANUARY, MORNING

**MARK
SCHEME**

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for correct working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidates's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

				AVAILABLE MARKS
1	Subtract to give Simplifying gives Substitute into 2nd equation Expanding	$8x - 4y - 40 = 0$ $y = 2x - 10$ $\Rightarrow x^2 + (2x - 10)^2 - 10x + 20 = 0$ $\Rightarrow x^2 + 4x^2 - 40x + 100 - 10x + 20 = 0$ $\Rightarrow 5x^2 - 50x + 120 = 0$ $\Rightarrow x^2 - 10x + 24 = 0$ $\Rightarrow (x - 6)(x - 4) = 0$ $\Rightarrow x = 6, x = 4$ $\Rightarrow y = 2, y = -2$	MW1 MW1 M1 W1 W1 W1 W1	
	Hence the points of intersection are (6, 2) and (4, -2)		W1	8
2	(i)	$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$	M1	
	and	$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ -2 \end{pmatrix}$	MW1	
	Hence	$a - b = 1$ ①		
		$a - 2b = 3$ ②	MW1	
	and	$c - d = -1$ ③		
		$c - 2d = -6$ ④	MW1	
		① - ② $\Rightarrow b = -2$		
	Hence	$a = -1$	W1	
		③ - ④ $\Rightarrow d = 5$		
	Hence	$c = 4$	W1	
	Therefore	$\mathbf{M} = \begin{pmatrix} -1 & -2 \\ 4 & 5 \end{pmatrix}$		
	(ii)	$\mathbf{D} = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$ or $\mathbf{D} = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}$	MW1	7
3	(i)	$\begin{vmatrix} p & 2 & 4 \\ 1 & 0 & -5 \\ 0 & 3 & 1 \end{vmatrix} = 0$	M1	
		$\Rightarrow p(15) - 2(1) + 4(3) = 0$	MW1	
		$\Rightarrow 15p + 10 = 0$	W1	
		$\Rightarrow p = -\frac{2}{3}$	W1	

(ii) $\det \mathbf{N} = 15p + 10$

If $p = 1$ then $\det \mathbf{N} = 25$

Matrix of minors =
$$\begin{pmatrix} 15 & 1 & 3 \\ -10 & 1 & 3 \\ -10 & -9 & -2 \end{pmatrix}$$

Matrix of cofactors =
$$\begin{pmatrix} 15 & -1 & 3 \\ 10 & 1 & -3 \\ -10 & 9 & -2 \end{pmatrix}$$

Transpose =
$$\begin{pmatrix} 15 & 10 & -10 \\ -1 & 1 & 9 \\ 3 & -3 & -2 \end{pmatrix}$$

Inverse =
$$\frac{1}{25} \begin{pmatrix} 15 & 10 & -10 \\ -1 & 1 & 9 \\ 3 & -3 & -2 \end{pmatrix}$$

(iii) $\mathbf{N} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 5 \\ 3 \\ 3 \end{pmatrix}$

Hence $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \mathbf{N}^{-1} \begin{pmatrix} 5 \\ 3 \\ 3 \end{pmatrix}$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{25} \begin{pmatrix} 15 & 10 & -10 \\ -1 & 1 & 9 \\ 3 & -3 & -2 \end{pmatrix} \begin{pmatrix} 5 \\ 3 \\ 3 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{25} \begin{pmatrix} 75 \\ 25 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$$

Hence $x = 3, y = 1, z = 0$

MW1

AVAILABLE MARKS

MW3

MW1

MW1

MW1


M1

W1

W2

15

		MW3	M1	W1	W1	AVAILABLE MARKS
4	(a) Shear, parallel to the x -axis and with scale factor of 2					
	(b) $\begin{pmatrix} -1 & 3 \\ 2 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 13 \\ 4 \end{pmatrix}$					
	$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{-6} \begin{pmatrix} 0 & -3 \\ -2 & -1 \end{pmatrix} \begin{pmatrix} 13 \\ 4 \end{pmatrix}$					
	$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{-6} \begin{pmatrix} -12 \\ -30 \end{pmatrix}$					
	$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$					
	Hence P is the point (2, 5)					
	(c) The line $y = mx$ is mapped to $y = -\frac{1}{m}x$					
	Hence $\begin{pmatrix} 2 & 3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ mx \end{pmatrix} = \begin{pmatrix} t \\ -\frac{1}{m}t \end{pmatrix}$					
	$\Rightarrow 2x + 3mx = t$					
	$mx = -\frac{1}{m}t$					
	Divide to give $\frac{2 + 3m}{m} = 1 \div -\frac{1}{m}$					
	Hence $\frac{2 + 3m}{m} = \frac{-m}{1}$					
	$\Rightarrow 2 + 3m = -m^2$					
	$\Rightarrow m^2 + 3m + 2 = 0$					
	$\Rightarrow (m + 1)(m + 2) = 0$					
	$\Rightarrow m = -1, -2$					
5	(i) $ab = ba$					
	$\Rightarrow bab = b^2a$					
	$\Rightarrow bab = ea$					
	$\Rightarrow bab = a$					
	(ii) $abab = a(bab) = a^2$					
	(iii) $a^2 ba^2 b = aabaab = aababa = a(abab)a = aa^2a = a^4 = a$					
	(iv) e and b					

		AVAILABLE MARKS
(v)	Not isomorphic since there are 4 elements in H which are self-inverse and only 2 in G are self-inverse.	MW2
6 (a) $z = \frac{-8 \pm \sqrt{64 - 4(5)(5)}}{10}$	M1	12
$\Rightarrow z = \frac{-8 \pm \sqrt{-36}}{10}$	W1	
$\Rightarrow z = \frac{-8 \pm 6i}{10}$	MW2	
$\Rightarrow z = -0.8 \pm 0.6i$	MW2	
(b) $\begin{aligned} 5 + 2ai &= 7 - ai \\ \Rightarrow 25 + 4a^2 &= 49 + a^2 \\ \Rightarrow 3a^2 &= 24 \\ \Rightarrow a^2 &= 8 \\ \Rightarrow a &= \pm 2\sqrt{2} \end{aligned}$	MW2	
	W1	
	W2	
(c) (i) Circle, centre (6, 4), radius 3	MW3	
		
(ii) Vertical half line through (1, -2)	MW3	
(iii) Radius of circle = 3	MW1	
Distance from centre of circle to line equals 5	M1	
Minimum value of $ z - w = 5 - 3 = 2$	W1	18
Total	75	