



**ADVANCED
General Certificate of Education
January 2014**

Mathematics
Assessment Unit F2
assessing
Module FP2: Further Pure Mathematics 2

[AMF21]

MONDAY 27 JANUARY, MORNING

**MARK
SCHEME**

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

		AVAILABLE MARKS
1	(i) $(\sqrt{r} + \sqrt{r-1})(\sqrt{r} - \sqrt{r-1}) = r - (r-1) = 1$	MW1
	(ii) $\begin{aligned} \sum_{r=1}^n \frac{1}{\sqrt{r} + \sqrt{r-1}} &= \sum_{r=1}^n \frac{1}{\sqrt{r} + \sqrt{r-1}} \cdot \left(\frac{\sqrt{r} - \sqrt{r-1}}{\sqrt{r} - \sqrt{r-1}} \right) \\ &= \sum (\sqrt{r} - \sqrt{r-1}) \\ &= (1-0) + (\sqrt{2}-1) + \dots + (\sqrt{n}-\sqrt{n-1}) \\ &= \sqrt{n} \end{aligned}$	MW1 W1 W1
		4
2	$3 \tan^2 \theta = 2 \sin \theta$ $3 \frac{\sin^2 \theta}{\cos^2 \theta} = 2 \sin \theta$ $3 \sin^2 \theta = 2 \sin \theta (1 - \sin^2 \theta)$ $3 \sin \theta = 2(1 - \sin^2 \theta)$ or $\sin \theta = 0$ $2 \sin^2 \theta + 3 \sin \theta - 2 = 0$ $(2 \sin \theta - 1)(\sin \theta + 2) = 0$ $\sin \theta = \frac{1}{2}$ or $\sin \theta = -2$ $\sin \theta = -2 \Rightarrow$ no real solution $\sin \theta = \frac{1}{2} \Rightarrow \theta = 2n\pi + \frac{\pi}{6}$ or $2n\pi + \frac{5\pi}{6}$ $\sin \theta = 0 \Rightarrow \theta = n\pi \quad n = 0, 1, 2, \dots$	MW1 M1 W1 W1 MW1 MW1 MW1 MW1 MW1
		9
3	(a) (i) $z = 1 - i$ reason – equation has real coefficients or equivalent	MW1 M1
	(ii) $\begin{aligned} [z - (1 - i)][z - (1 + i)] &= z^2 - 2z + 2 \\ &\quad \underline{z^2 - 4z + 13} \\ z^2 - 2z + 2 &\overline{z^4 - 6z^3 + 23z^2 - 34z + 26} \\ &\quad \underline{z^4 - 2z^3 + 2z^2} \\ &\quad \underline{-4z^3 + 21z^2 - 34z} \\ &\quad \underline{-4z^3 + 8z^2 - 8z} \\ &\quad \underline{13z^2 - 26z + 26} \\ &\quad \underline{13z^2 - 26z + 26} \\ &\quad \underline{0} \\ z^2 - 4z + 13 &= 0 \\ z &= \frac{4 \pm \sqrt{16 - 52}}{2} \\ &= 2 \pm 3i \end{aligned}$	MW1 M1 W1 M1 MW1
		12

			AVAILABLE MARKS
4	(i) $\frac{dy}{dx} = \sec^2 x = 1 + \tan^2 x = 1 + y^2$	MW1	W1
	$\frac{d^2y}{dx^2} = 2y \frac{dy}{dx} = 2y(1 + y^2)$	MW1	
(ii)	$\frac{d^3y}{dx^3} = 2y \cdot 2y \frac{dy}{dx} + (1 + y^2) 2 \frac{dy}{dx}$ $= 6y^2 \frac{dy}{dx} + 2 \frac{dy}{dx} = (6y^2 + 2)(1 + y^2) = 6y^4 + 8y^2 + 2$	M1	W1
(iii)	$y = y(0) + x \left(\frac{dy}{dx} \right)_{x=0} + \frac{x^2}{2!} \left(\frac{d^2y}{dx^2} \right)_{x=0} + \frac{x^3}{3!} \left(\frac{d^3y}{dx^3} \right)_{x=0}$	M1	
	$x = 0 \ y = 0 \quad y = 0 + x + 0 + \frac{x^3}{6} \cdot 2$ $= x + \frac{x^3}{3}$		W3
(iv)	For small x		
	$\frac{x \sin^2 x}{\tan x - x \cos x} \approx \frac{x^3}{x + \frac{x^3}{3} - x \left(1 - \frac{x^2}{2}\right)}$ $= \frac{x^3}{\frac{5}{6}x^3} = \frac{6}{5}$	MW2	
		W1	12
5	Let $n = 1$ L.H.S. = $\cos x$		
	R.H.S. = $\frac{\sin 2x}{2 \sin x} = \frac{2 \sin x \cos x}{2 \sin x} = \cos x$	M1	W1
	Assume true for $n = k$ i.e. $\cos x + \cos 3x + \cos 5x + \dots + \cos(2k-1)x$	M1	
	Let $n = k + 1$		
	$\cos x + \cos 3x + \dots + \cos(2k-1)x + \cos(2k+1)x$ $= \frac{\sin 2kx}{2 \sin x} + \cos(2k+1)x$	MW2	
	$= \frac{\sin 2kx + (\cos 2kx \cos x - \sin 2kx \sin x) 2 \sin x}{2 \sin x}$	M2	
	$= \frac{\cos 2kx \sin 2x + \sin 2kx (1 - 2 \sin^2 x)}{2 \sin x}$		
	$= \frac{\cos 2kx \sin 2x + \sin 2kx \cos 2x}{2 \sin x} = \frac{\sin 2(k+1)x}{2 \sin x}$	MW1	
	True $n = k \Rightarrow$ true $n = k + 1$		
	True $n = 1$		
	\therefore True for all n	MW1	9

			AVAILABLE MARKS
6	Auxiliary Equation	$m^2 + n^2 = 0$	MW1
		$m = \pm in$	W1
	Complementary Function	$x = A \cos nt + B \sin nt$	W1
	Let $x = C \cos \omega t + D \sin \omega t$		M1
	$\frac{dx}{dt} = -\omega C \sin \omega t + \omega D \cos \omega t$		MW1
	$\frac{d^2x}{dt^2} = -\omega^2 C \cos \omega t - \omega^2 D \sin \omega t$		MW1
	$-(\omega^2 - n^2)C = 1 \quad \therefore C = \frac{1}{n^2 - \omega^2} \quad \omega \neq n$		
	$(-\omega^2 + n^2)D = 0 \quad \therefore D = 0 \quad \omega \neq n$		MW1
	General Solution $x = A \cos nt + B \sin nt + \frac{1}{n^2 - \omega^2} \cos \omega t$		MW1
	$\frac{dx}{dt} = -nA \sin nt + nB \cos nt - \frac{\omega}{n^2 - \omega^2} \sin \omega t$		W1
	$x = 0 \text{ when } t = 0 \quad \Rightarrow \quad A = -\frac{1}{n^2 - \omega^2}$		W1
	$\frac{dx}{dt} = V \text{ when } t = 0 \quad \therefore V = 0 + nB \Rightarrow B = \frac{V}{n}$		W1
	Solution $x = \frac{1}{n^2 - \omega^2} (\cos \omega t - \cos nt) + \frac{V}{n} \sin nt$		W1
			12
7	(i) $b^2 = a^2(1 - e^2)$		MW1
	$4 = 9(1 - e^2) \quad \therefore e = \frac{\sqrt{5}}{3}$		W1
	(ii) foci $(\pm \sqrt{5}, 0)$ directrices $x = \pm \frac{a}{e} = \pm \frac{9}{\sqrt{5}}$		MW1 MW1
	(iii) $\frac{x^2}{9} + \frac{y^2}{4} = 1$		
	$\frac{2x}{9} + \frac{2y}{4} \cdot \frac{dy}{dx} = 0 \quad \frac{dy}{dx} = \frac{-4x}{9y}$		M1
	At P $\frac{dy}{dx} = \frac{-2 \cos \theta}{3 \sin \theta}$		W1
	$y - 2 \sin \theta = \frac{-2 \cos \theta}{3 \sin \theta} (x - 3 \cos \theta)$		MW1
	$3y \sin \theta - 6 \sin^2 \theta = -2x \cos \theta + 6 \cos^2 \theta$		
	$\frac{y \sin \theta}{2} + \frac{x \cos \theta}{3} = 1$		W1

		AVAILABLE MARKS
(iv) line \perp to tangent	$m = \frac{3 \sin \theta}{2 \cos \theta}$	MW1
Equation of line	$y = \frac{3 \sin \theta}{2 \cos \theta} x$	MW1
Meets tangent	$\frac{3 \sin^2 \theta}{4 \cos \theta} x + \frac{x \cos \theta}{3} = 1$	M1
	$x \frac{(9 \sin^2 \theta + 4 \cos^2 \theta)}{12 \cos \theta} = 1$	
	$x = \frac{12 \cos \theta}{(5 \sin^2 \theta + 4)}$	W1
	$y = \frac{18 \sin \theta}{(5 \sin^2 \theta + 4)}$	W1
(v)	$x^2 + y^2 = \frac{144 \cos^2 \theta + 324 \sin^2 \theta}{(5 \sin^2 \theta + 4)^2} = \frac{180 \sin^2 \theta + 144}{(5 \sin^2 \theta + 4)^2}$	M1
	$= \frac{36}{(5 \sin^2 \theta + 4)}$	
	$(x^2 + y^2)^2 = \frac{36^2}{(5 \sin^2 \theta + 4)^2}$	
	$9x^2 + 4y^2 = \frac{9 \times 144 \cos^2 \theta + 4 \times 324 \sin^2 \theta}{(5 \sin^2 \theta + 4)^2} = \frac{36^2}{(5 \sin^2 \theta + 4)^2}$	M1 W1
	$\therefore (x^2 + y^2)^2 = 9x^2 + 4y^2$	W1
		17
	Total	75