



ADVANCED
General Certificate of Education
2012

Mathematics

Assessment Unit C4

assessing

Module C4: Core Mathematics 4

[AMC41]



FRIDAY 1 JUNE, MORNING

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.

Answer **all eight** questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

You are permitted to use a graphic or scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that $\ln z \equiv \log_e z$.



Answer all eight questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

- 1 (a) For each relation below state if it is a one-one function, a many-one function or a mapping.

(i) $a: x \rightarrow \pm\sqrt{x} \quad x \in \mathbb{R} \quad x \geq 0$ [1]

(ii) $b: x \rightarrow \frac{1}{x} \quad x \in \mathbb{R} \quad x \neq 0$ [1]

(iii) $c: x \rightarrow x^2 \quad x \in \mathbb{R}$ [1]

- (b) The function f is defined by

$$f(x) = x^2 - 3 \quad x \in \mathbb{R}$$

and the function g is defined by

$$g(x) = 2x + 1 \quad x \in \mathbb{R}$$

- (i) Find the composite function $gf(x)$, stating its domain. [3]

- (ii) Find the values of x such that $gf(x) = 3x$ [3]

- 2 (i) Sketch the graph of

$$y = 4 - x^2$$

clearly showing where it crosses the x -axis. [2]

A paperweight can be modelled as the solid formed when the area bounded by the curve

$$y = 4 - x^2$$

and the x -axis and the y -axis is rotated through 360° about the x -axis.

- (ii) Find the volume of the paperweight. [7]

- 3 During a science experiment, students create two waves in a ripple tank. The first wave can be modelled by the equation

$$h_1 = 6 \cos t \quad 0 \leq t \leq 2\pi$$

The second wave can be modelled by the equation

$$h_2 = 5 \sin t \quad 0 \leq t \leq 2\pi$$

where h_1 and h_2 are the heights of the waves, measured in millimetres, at any time t seconds from the start of the experiment.

The two waves join together.

- (i) Express the height $h_1 + h_2$ of the resultant wave in the form

$$r \cos(t - \alpha) \quad 0 \leq \alpha \leq \frac{\pi}{2} \quad r \in \mathbb{R} \quad [4]$$

- (ii) Hence find the times when the resultant wave has a height of 3 mm. [4]

- 4 (i) Given that

$$x^2 + 6xy + y^2 + 32 = 0$$

use implicit differentiation to show that

$$\frac{dy}{dx} = \frac{-(x+3y)}{3x+y} \quad [5]$$

- (ii) Hence find the coordinates of the stationary points on the curve

$$x^2 + 6xy + y^2 + 32 = 0 \quad [6]$$

- 5 Solve the differential equation

$$\cos^2 4x \frac{dy}{dx} = y$$

given that $y = e^3$ when $x = \frac{\pi}{16}$ [8]

- 6 (i) Find the vector equation of the line l through the points $(2, 4, 3)$ and $(1, 2, 3)$. [4]

The point P lies on the line l .
 O is the point $(0, 0, 0)$.

- (ii) Hence, using (i), write down the position vector \overrightarrow{OP} [1]

OP is perpendicular to the line l .

- (iii) Find the coordinates of the point P . [6]

- 7 Find the **exact** value of

$$\int_1^3 \ln x \, dx \quad [7]$$

- 8 (i) Prove the identity

$$\tan 3\theta \equiv \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \quad [7]$$

- (ii) Hence solve the equation

$$\tan 3\theta = \tan \theta \quad 0^\circ \leq \theta \leq 360^\circ \quad [5]$$

THIS IS THE END OF THE QUESTION PAPER
