



ADVANCED SUBSIDIARY (AS)
General Certificate of Education
2014

Mathematics
Assessment Unit F1
assessing
Module FP1: Further Pure Mathematics 1
[AMF11]

TUESDAY 24 JUNE, MORNING

**MARK
SCHEME**

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for correct working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidates's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.


When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

		AVAILABLE MARKS
1	$\mathbf{ABC} = \mathbf{I}$ $\Rightarrow \mathbf{A}^{-1} \mathbf{ABC} = \mathbf{A}^{-1} \mathbf{I}$ $\Rightarrow \mathbf{BC} = \mathbf{A}^{-1}$ $\Rightarrow \mathbf{BCC}^{-1} = \mathbf{A}^{-1} \mathbf{C}^{-1}$ $\Rightarrow \mathbf{B} = \mathbf{A}^{-1} \mathbf{C}^{-1}$	MW1
	$\mathbf{A}^{-1} = \frac{1}{1} \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix}$	MW1
	$\mathbf{C}^{-1} = \frac{1}{2} \begin{pmatrix} 2 & 0 \\ -4 & 1 \end{pmatrix}$	MW1
	Hence $\mathbf{B} = \frac{1}{2} \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ -4 & 1 \end{pmatrix}$ $= \frac{1}{2} \begin{pmatrix} -14 & 5 \\ -6 & 2 \end{pmatrix}$ $= \begin{pmatrix} -7 & 2\frac{1}{2} \\ -3 & 1 \end{pmatrix}$	MW1
		5
2	(i) $x^2 + y^2 + 2x - 6y - 15 = 0$ ① $x^2 + y^2 + 2y - 3 = 0$ ② $\text{①} - \text{②} \Rightarrow 2x - 8y - 12 = 0$ $\Rightarrow x = 4y + 6$	M1 W1
	Substitute into ②	M1
	$\Rightarrow (4y + 6)^2 + y^2 + 2y - 3 = 0$	W1
	$\Rightarrow 16y^2 + 48y + 36 + y^2 + 2y - 3 = 0$	
	$\Rightarrow 17y^2 + 50y + 33 = 0$	W1
	$\Rightarrow (17y + 33)(y + 1) = 0$	
	$\Rightarrow y = -1, -\frac{33}{17}$	W1
	$\Rightarrow x = 2, -\frac{30}{17}$	W1
	Points of intersection are $(2, -1)$ and $(-\frac{30}{17}, -\frac{33}{17})$	W1

(ii) $C_2: x^2 + (y + 1)^2 = 4$
 Hence centre is $(0, -1)$ and radius is 2
 $C_3: x^2 + y^2 - x - 4y - 31\frac{3}{4} = 0$
 $\Rightarrow (x - \frac{1}{2})^2 + (y - 2)^2 = 36$
 Hence centre is $(\frac{1}{2}, 2)$ and radius is 6

MW2

MW2



Therefore distance between centres $P_2P_3 = \sqrt{(\frac{1}{2})^2 + 3^2}$

M1

$$= \sqrt{\frac{37}{4}}$$

$$= \frac{\sqrt{37}}{2}$$

W1

For C_2 to lie inside C_3 it is necessary that $P_2P_3 < r_3 - r_2$

M1

Substituting values we have $\frac{\sqrt{37}}{2} < 6 - 2$ QED

MW1

Hence C_2 lies entirely inside C_3

16

3 (i) $(a, b) * [(c, d) * (e, f)]$
 $= (a, b) * (ce, d + f + 2)$
 $= (ace, b + d + f + 2 + 2)$
 Also, $[(a, b) * (c, d)] * (e, f)$
 $= (ac, b + d + 2) * (e, f)$
 $= (ace, b + d + f + 2 + 2)$
 Hence the operation is associative.

M1

W1

M1

W1

(ii) $(a, b) * (c, d) = (a, b)$
 $\Rightarrow (ac, b + d + 2) = (a, b)$
 $\Rightarrow ac = a \quad \text{and} \quad b + d + 2 = b$
 $\Rightarrow c = 1 \quad \Rightarrow d = -2$
 Hence identity = $(1, -2)$
 Check also that $(1, -2) * (a, b) = (a, -2 + b + 2) = (a, b)$

M1

M1

W2

8

AVAILABLE MARKS

		AVAILABLE MARKS
4	(i)	
	$\begin{vmatrix} 5 & -2 & -a \\ 4 & 2 & -6 \\ 1 & a & -4 \end{vmatrix}$ $= 5(-8 + 6a) + 2(-16 + 6) - a(4a - 2)$ $= -40 + 30a - 32 + 12 - 4a^2 + 2a$ $= -4a^2 + 32a - 60$	M1 W1 W1
	(ii) When $a = 3$, determinant $= -4(9) + 32(3) - 60 = 0$ Therefore there is not a unique solution.	MW1 MW1
	$5x - 2y - 3z = 3 \quad ①$ $4x + 2y - 6z = 2 \quad ②$ $x + 3y - 4z = 0 \quad ③$ $① + ② \Rightarrow 9x - 9z = 5 \quad ④$ $3 \times ② - 2 \times ③ \Rightarrow 10x - 10z = 6 \quad ⑤$ $④ \Rightarrow x - z = \frac{5}{9}$ $⑤ \Rightarrow x - z = \frac{6}{10}$	M1 MW1
	Since these equations contradict, there is no solution.	MW1
	When $a = 4$, determinant $= -4(16) + 32(4) - 60 = 4$ Since determinant is non-zero, there is one unique solution.	MW1 MW1
	(iii) $5x - 2y - 5z = 3 \quad ①$ $4x + 2y - 6z = 2 \quad ②$ $x + 5y - 4z = 0 \quad ③$ $① + ② \Rightarrow 9x - 11z = 5 \quad ④$ $5 \times ② - 2 \times ③ \Rightarrow 18x - 22z = 10 \quad ⑤$ Since $⑤ = 2 \times ④$, these equations are consistent. Re-arranging $④ \Rightarrow x = \frac{5 + 11z}{9}$ Substituting into $③ \Rightarrow \frac{5 + 11z}{9} + 5y - 4z = 0$ $\Rightarrow 5 + 11z + 45y - 36z = 0$ $\Rightarrow 5 + 45y = 25z$ $\Rightarrow y = \frac{5z - 1}{9}$	M1 MW1 MW1 MW1
	Hence general solution is $\left(\frac{5 + 11z}{9}, \frac{5z - 1}{9}, z\right)$	14

		AVAILABLE MARKS
5	(a) $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 2 & 5 \\ 3 & -1 \end{pmatrix} = \begin{pmatrix} 7 & 9 \\ 2 & 5 \end{pmatrix}$ $\Rightarrow 2a + 3b = 7 \quad \text{and} \quad 2c + 3d = 2 \quad \text{M1}$ $5a - b = 9 \quad \text{and} \quad 5c - d = 5 \quad \text{M1}$ $\Rightarrow 2a + 3b = 7 \quad \text{and} \quad 2c + 3d = 2$ $15a - 3b = 27 \quad \text{and} \quad 15c - 3d = 15 \quad \text{M1}$ $\Rightarrow 17a = 34 \quad \text{and} \quad 17c = 17$ $\Rightarrow a = 2 \quad \text{and} \quad c = 1 \quad \text{W1}$ $\Rightarrow b = 1 \quad \text{and} \quad d = 0 \quad \text{W1}$ Hence $\mathbf{M} = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}$	
(b)	$\begin{pmatrix} 0 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} X \\ Y \end{pmatrix} \quad \text{M1}$ $\Rightarrow -y = X \quad \textcircled{1}$ $x + 2y = Y \quad \textcircled{2}$ $\textcircled{1} \Rightarrow y = -X \quad \text{MW1}$ $\textcircled{2} \Rightarrow x - 2X = Y$ $\Rightarrow x = 2X + Y \quad \text{MW1}$ Substitute into $x^2 + 5y^2 + 4xy - 6x - 8y = 0 \quad \text{M1}$ $\Rightarrow (2X + Y)^2 + 5(-X)^2 + 4(2X + Y)(-X) - 6(2X + Y) - 8(-X) = 0 \quad \text{W1}$ $\Rightarrow 4X^2 + 4XY + Y^2 + 5X^2 - 8X^2 - 4XY - 12X - 6Y + 8X = 0 \quad \text{W1}$ $\Rightarrow X^2 + Y^2 - 4X - 6Y = 0 \quad \text{W1}$ $\Rightarrow (X - 2)^2 + (Y - 3)^2 = 13$ This is the equation of a circle of radius $\sqrt{13} \quad \text{MW1}$	14

		AVAILABLE MARKS
6	(a) (i) $z_1 = \sqrt{3} + i$ $\Rightarrow r_1 \cos \theta_1 = \sqrt{3}$ and $r_1 \sin \theta_1 = 1$ $\Rightarrow \tan \theta_1 = \frac{1}{\sqrt{3}}$ $\Rightarrow \theta_1 = \frac{\pi}{6}$ and $r_1 = 2$	M1 M1 W1 W1
	$z_2 = 1 - i$ $\Rightarrow r_2 \cos \theta_2 = 1$ and $r_2 \sin \theta_2 = -1$ $\Rightarrow \tan \theta_2 = -1$ $\Rightarrow \theta_2 = -\frac{\pi}{4}$ and $r_2 = \sqrt{2}$	MW1 MW1
(ii)	$z_1 z_2 = (\sqrt{3} + i)(1 - i)$ $= \sqrt{3} + 1 + i - \sqrt{3}i$ $\Rightarrow z_1 z_2 = \sqrt{(\sqrt{3} + 1)^2 + (1 - \sqrt{3})^2}$ $= \sqrt{3 + 2\sqrt{3} + 1 + 1 + 1 - 2\sqrt{3} + 3}$ $= \sqrt{8}$ $= 2\sqrt{2}$ $= z_1 z_2 $	MW1 M1 W1 MW1
	QED	


(b) (i) Perpendicular bisector of line joining $(4, 15)$ and $(0, -1)$

(ii) Half-line through $(2, 7)$ at angle of $\frac{\pi}{6}$ with positive x -axis

MW3

MW3

AVAILABLE MARKS



(iii) Region as shown shaded in diagram above.

MW2

18

Total**75**