

Mathematics
Higher level
Paper 3 – sets, relations and groups

Wednesday 18 November 2015 (afternoon)

1 hour

Instructions to candidates

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A graphic display calculator is required for this paper.
- A clean copy of the **mathematics HL and further mathematics HL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[60 marks]**.

Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 5]

Given the sets A and B , use the properties of sets to prove that $A \cup (B' \cup A)' = A \cup B$, justifying each step of the proof.

2. [Maximum mark: 14]

The function $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined as $f: x \rightarrow \begin{cases} 1 & , x \geq 0 \\ -1 & , x < 0 \end{cases}$.

(a) Prove that f is

(i) not injective;

(ii) not surjective.

[4]

The relation R is defined for $a, b \in \mathbb{R}$ so that aRb if and only if $f(a) \times f(b) = 1$.

(b) Show that R is an equivalence relation.

[8]

(c) State the equivalence classes of R .

[2]

3. [Maximum mark: 10]

The set of all permutations of the elements $1, 2, \dots, 10$ is denoted by H and the binary operation \circ represents the composition of permutations.

The permutation $p = (1\ 2\ 3\ 4\ 5\ 6)(7\ 8\ 9\ 10)$ generates the subgroup $\{G, \circ\}$ of the group $\{H, \circ\}$.

(a) Find the order of $\{G, \circ\}$. [2]

(b) State the identity element in $\{G, \circ\}$. [1]

(c) Find

(i) $p \circ p$;

(ii) the inverse of $p \circ p$. [4]

(d) (i) Find the maximum possible order of an element in $\{H, \circ\}$.

(ii) Give an example of an element with this order. [3]

4. [Maximum mark: 18]

The binary operation $*$ is defined on the set $T = \{0, 2, 3, 4, 5, 6\}$ by $a * b = (a + b - ab)(\text{mod } 7)$, $a, b \in T$.

(a) Copy and complete the following Cayley table for $\{T, *\}$. [4]

*	0	2	3	4	5	6
0	0	2	3	4	5	6
2	2	0	6	5	4	3
3	3	6				
4	4	5				
5	5	4				
6	6	3				

(b) Prove that $\{T, *\}$ forms an Abelian group. [7]

(c) Find the order of each element in T . [4]

(d) Given that $\{H, *\}$ is the subgroup of $\{T, *\}$ of order 2, partition T into the left cosets with respect to H . [3]

5. [Maximum mark: 13]

A group $\{D, \times_3\}$ is defined so that $D = \{1, 2\}$ and \times_3 is multiplication modulo 3.

A function $f: \mathbb{Z} \rightarrow D$ is defined as $f: x \mapsto \begin{cases} 1, & x \text{ is even} \\ 2, & x \text{ is odd} \end{cases}$.

- (a) Prove that the function f is a homomorphism from the group $\{\mathbb{Z}, +\}$ to $\{D, \times_3\}$. [6]
- (b) Find the kernel of f . [3]
- (c) Prove that $\{\text{Ker}(f), +\}$ is a subgroup of $\{\mathbb{Z}, +\}$. [4]
-