

GCE

Chemistry A

H432/01: Periodic table, elements and physical chemistry

Advanced GCE

Mark Scheme for Autumn 2021

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021



Tuesday 5 October 2021 –Afternoon

A Level Chemistry A

H432/01 Periodic table, elements and physical chemistry

MARK SCHEME















Duration: 2 hours 15 minutes

MAXIMUM MARK 100

**Last updated: 17/10/2021
Post-standardisation**

This document consists of 27 pages


1. Annotations

Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore

2. Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

Annotation	Meaning
/	alternative and acceptable answers for the same marking point
✓	Separates marking points
DO NOT ALLOW	Answers which are not worthy of credit
IGNORE	Statements which are irrelevant
ALLOW	Answers that can be accepted
()	Words which are not essential to gain credit
<u>—</u>	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument

Question	Answer	Marks	AO element	Guidance
1	C	1	AO1.1	
2	B	1	AO1.2	
3	D	1	AO2.6	
4	B	1	AO2.2	
5	D	1	AO2.6	
6	C	1	AO2.6	
7	A	1	AO1.1	
8	B	1	AO2.2	
9	B	1	AO2.2	
10	A	1	AO2.6	
11	A	1	AO1.2	
12	C	1	AO1.2	
13	D	1	AO1.1	Accept 1
14	B	1	AO2.1	
15	C	1	AO2.3	
	Total	15		

Question		Answer	Marks	AO element	Guidance																									
16	(a)	 <p>(delocalised) electrons</p> <p>Diagram with regular arrangement of labelled 'Mg²⁺ ions' OR '2+ ions' AND attempt to show electrons ✓</p> <p>Labelled electrons between other species AND statement anywhere of delocalised electrons (can be in text or in diagram)</p> <p>Electrons move ✓</p>	3		<p>Regular arrangement must have at least two rows of correctly charged ions and a minimum of two ions per row</p> <p>ALLOW as label: +2 ions OR + 2 cations OR +2/2+ seen within circle</p> <p>ALLOW e⁻ or 'e' as a label for electron</p> <p>IGNORE "-" for electron label</p> <p>ALLOW mobile/flow for move</p> <p>IGNORE 'carry charge'</p>																									
	(b)	(i)	$\text{Mg}^{3+}(\text{g}) \rightarrow \text{Mg}^{4+}(\text{g}) + \text{e}^{-} \checkmark$	1	AO1.2	<p>State symbols required (ignore states on electrons)</p> <p>ALLOW $\text{Mg}^{3+}(\text{g}) - \text{e}^{-} \rightarrow \text{Mg}^{4+}(\text{g})$</p> <p>ALLOW $\text{Mg}^{+3}(\text{g})$</p> <p>ALLOW e for e⁻</p>																								
	(b)	(ii)	Big jump/larger difference between 2 and 3 ✓	1	AO1.2	<p>IGNORE big jump between 10 and 11</p> <p>DO NOT ALLOW other combinations.</p>																								
	(b)	(iii)	<p>1st AND 3rd AND 4th AND 5th AND 9th AND 11th ✓ i.e.</p> <table border="1" data-bbox="414 1228 1075 1348"> <tr> <td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td><td>11</td><td>12</td> </tr> <tr> <td>✓</td><td></td><td>✓</td><td>✓</td><td>✓</td><td></td><td></td><td></td><td>✓</td><td></td><td>✓</td><td></td> </tr> </table>	1	2	3	4	5	6	7	8	9	10	11	12	✓		✓	✓	✓				✓		✓		1	AO2.1	
1	2	3	4	5	6	7	8	9	10	11	12																			
✓		✓	✓	✓				✓		✓																				

Question		Answer	Marks	AO element	Guidance
17	(a)	<p>Transition element: Has an ion with an incomplete/partially-filled d sub-shell/d-orbital ✓</p> <p>d-block d sub-shell/d-orbital is being filled/has highest energy OR Electron configurations shown for Sc: $1s^22s^22p^63s^23p^63d^14s^2$ AND Zn: $1s^22s^22p^63s^23p^63d^{10}4s^2$ ✓</p> <p><i>Electron configurations of ions</i> Sc³⁺: $1s^22s^22p^63s^23p^6$ AND d sub-shell empty / d orbital(s) empty ✓</p> <p>Zn²⁺: $1s^22s^22p^63s^23p^63d^{10}$ AND d sub-shell full / d-orbitals full ✓</p>	4	AO1.1 x4	<p>FULL ANNOTATIONS MUST BE USED -----</p> <p>DO NOT ALLOW d shell</p> <p>IGNORE d block</p> <p>IGNORE outer electron</p> <p>electron configurations ALLOW 4s⁰ ALLOW 4s² before 3d, i.e.4s²3d¹; 4s²3d¹⁰ IGNORE other Sc and Zn ions</p> <p>ALLOW ECF for short hand notation.</p> <p>For Sc³⁺, ALLOW Sc⁺³ OR Sc forms a 3+ ion;</p> <p>For Zn²⁺, ALLOW Zn⁺² OR Zn forms a 2+ ion;</p>
	(b)	(i)	1	AO1.1 x1	<p>ALLOW lone pairs for electron pairs</p> <p>ALLOW dative (covalent) bonds for coordinate bonds</p> <p>TWO is only needed once if bonds are plural, e.g. Donates 2 electron pairs to form coordinate bonds Donates electron pairs to form 2 coordinate bonds</p>

Question	Answer	Marks	AO element	Guidance
				<ul style="list-style-type: none"> Consistent use of 2 'out wedges', 2 'in wedges', 2 lines in plane of paper OR 4 lines, 1 'out wedge' and 1 'in wedge' <p>ALLOW following orientations</p> <p>Detail</p> <ul style="list-style-type: none"> Most bonding shown from Cr to O of H₂O and O⁻ C₂O₄²⁻
	Total	11		

Question		Answer	Marks	AO element	Guidance
18	(a)	Formula: CuCO_3 ✓ $\text{CuCO}_3 + 2\text{HNO}_3 \rightarrow \text{Cu}(\text{NO}_3)_2 + \text{CO}_2 + \text{H}_2\text{O}$ ✓	2	AO1.2 AO2.6	IGNORE state symbols ALLOW formula within equation. ALLOW other copper(II) compounds which can react with nitric acid to form a gas e.g. CuS , CuSO_3 for mark 1, with correct equation for mark 2. e.g. $\text{CuSO}_3 + 2\text{HNO}_3 \rightarrow \text{Cu}(\text{NO}_3)_2 + \text{SO}_2 + \text{H}_2\text{O}$
	(b)	$2\text{Cu}^{2+}(\text{aq}) + 4\text{I}^{-}(\text{aq}) \rightarrow 2\text{CuI}(\text{s}) + \text{I}_2(\text{aq})$ ✓	1	AO2.6	ALLOW multiples State symbols are required
	(c)	starch (solution) AND blue-black to colourless ✓	1	AO1.2	ALLOW blue OR black OR purple for colour of mixture ALLOW blue colour disappears (to colourless) IGNORE 'clear' IGNORE 'colorimetry'
	(d)	FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 1.35 award 4 marks ----- $n(\text{S}_2\text{O}_3^{2-}) = 0.0200 \times \frac{26.55}{1000}$ $= 5.31 \times 10^{-4} \text{ (mol) } \checkmark$ $n(\text{I}_2) = 2.655 \times 10^{-4}$ OR $n(\text{Cu}^{2+}) = 5.31 \times 10^{-4} \text{ (mol) } \checkmark$ $m(\text{Cu}/\text{Cu}^{2+}) \text{ in ore} = 63.5 \times 5.31 \times 10^{-4}$ $= 0.0337 \dots \text{ (g) } \checkmark$ $\text{percentage} = \frac{0.0337 \dots}{2.50} \times 100$ $= 1.35 \text{ (\%)} \checkmark$ (3SF required)	4	AO2.8 ×5	FULL ANNOTATIONS MUST BE USED ----- ALLOW ECF throughout If 1:2 ratio for $\text{I}_2:\text{Cu}^{2+}$ not used check ratio in b) and allow ECF IGNORE rounding errors after 3 SF Calculator: 0.0337185 ALLOW 3 SF (0.0337) up to calculator value ECF dependent on the use of a calculated mass of Cu/Cu^{2+}

Question		Answer	Marks	AO element	Guidance	
	(e)	(i)	Lower AND smaller titre ✓	1	AO3.4	ALLOW less I ₂ produced / less Cu ²⁺ reacts
		(ii)	The same AND burette measures by difference ✓	1	AO3.4	ALLOW AW
	(f)		Any two of the following: Make up a (standard solution) from Step 2 to a stated volume (e.g. 250 cm ³) OR Repeat titrations AND Take mean of concordant/closest titres/ identify anomalies OR lower [S ₂ O ₃] ²⁻ to increase titre volume (to reduce the percentage error). OR higher [S ₂ O ₃] ²⁻ so not to refill the burette. OR Use a 3 dec place balance (to reduce the percentage error).	2	AO3.4 x 2	
			Total	12		

Question			Answer	Marks	AO element	Guidance
19	(a)	(i)	<p>Complete circuit with voltmeter AND labelled salt bridge linking two half-cells ✓</p> <p>Cr electrode in Cr³⁺ ✓ Pt electrode in MnO₄⁻ AND H⁺ AND Mn²⁺ ✓</p>	3	AO1.2 ×3	<p>Half cells can be drawn in either order Half cells must show electrodes dipping into solutions</p> <p>ALLOW small gaps in circuit</p> <p>IGNORE any stated concentrations</p> <p>IGNORE state symbols</p> <p>In salt bridge, ALLOW any stated ion that may be present, e.g. Cr³⁺, MnO₄⁻, Mn²⁺, H⁺</p>
	(a)	(ii)	$5\text{Cr} + 3\text{MnO}_4^- + 24\text{H}^+ \rightarrow 5\text{Cr}^{3+} + 3\text{Mn}^{2+} + 12\text{H}_2\text{O} \checkmark$	1	AO2.6	<p>IGNORE state symbols</p> <p>ALLOW multiples</p>
	(b)	(i)	<p>Mn is oxidised from +6 (in MnO₄²⁻) to +7 (in MnO₄⁻) ✓ Mn is reduced from +6 (in MnO₄²⁻) to +4 (in MnO₂) ✓</p>	2	AO2.1 ×2	<p>IGNORE '6' (signs required) ALLOW after number, e.g. 5+ ALLOW 1 mark for correct oxidation numbers but not linked to oxidation/reduction. IGNORE any reference to electron loss/gain (even if wrong)</p>

Question		Answer	Marks	AO element	Guidance
	(b)	(ii)			
	(b)	(ii)	2	AO3.1 ×2	<p>IGNORE 'lower/higher' ALLOW reverse argument: System 5 more positive than system 3, etc Must be comparative ALLOW response in terms of E_{cell} $E = (+)1.14 \text{ V}$ for system 5 – system 3</p> <p>Shift dependent on systems 3 and 5 correctly identified</p>
	(c)	(i)	1	AO2.6	<p>ALLOW multiples ALLOW $\text{H}_2 + 2\text{OH}^- - 2\text{e}^- \rightarrow 2\text{H}_2\text{O}$ ALLOW equation with equilibrium sign</p>
	(c)	(ii)	1	AO1.2	
	(c)	(iii)	1	AO1.1	<p>ALLOW named fuel. e.g. hydrogen/H_2; ethanol; methanol, etc</p> <p>ALLOW fuel cell requires <u>continuous</u> supply of fuel AND oxygen/an oxidant OR fuel cell operates <u>continuously</u> as long as a fuel AND oxygen/an oxidant are added</p> <p>IGNORE 'reactants' 'products' and comments about pollution and efficiency</p>
			Total	11	

Question		Answer	Marks	AO element	Guidance
20	(a)	rate of forwards reaction = rate of backwards reaction OR concentrations/pressure/temperature are constant /do not change ✓	1	AO1.1	DO NOT ALLOW “are the same”
	(b)	(i)	2	AO2.2 ×2	ALLOW -114000 – (298 × -147) ALLOW -70 up to calculator value of -70.194 correctly rounded, i.e. -70 OR -70.2 OR -70.19 ALLOW -70000 up to -70194 (J mol ⁻¹) ALLOW ECF for an incorrectly calculated negative value of ΔG linked to feasibility statement IGNORE rounding after 3 SF ORA for comment about – sign required for feasibility
	(b)	(ii)	1	AO2.2	
		776 (K) ✓ i.e. Maximum temperature = $\frac{\Delta H}{\Delta S} = \frac{-114}{-0.147} = 776$ (K) 3 SF required (appropriate from supplied data)			

Question		Answer	Marks	AO element	Guidance
	(c)	(i)	4	AO2.4 ×4	<p>FULL ANNOTATIONS MUST BE USED</p> <p>-----</p> <p>ALLOW ECF throughout</p> <p>ALLOW 20.6 from 3 SF partial pressures, 0.194, 0.436 and 0.581</p> <p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p> <p>-----</p> <p>Look for values to 3 SF here: 0.194, 0.436 and 0.581</p> <p>ALLOW 25.0 as ECF (from omission of partial pressures for 3 marks)</p>

FIRST, CHECK FOR VALUE OF K_p .
IF answer = 20.7 (MPa⁻¹), award 4 marks

Equilibrium amounts

$$n(\text{NO}) = 0.4 \text{ (mol)}$$

AND $n(\text{O}_2) = 0.9 \text{ (mol)}$

AND $n(\text{NO}_2) = 1.2 \text{ (mol)} \checkmark$

Total moles at equilibrium

$$n_{\text{tot}} = 2.5 \text{ (mol)} \checkmark$$

Partial pressures

$$p(\text{NO}) = \frac{0.4}{2.5} \times 1.21 = 0.1936 \text{ (MPa)}$$

AND $p(\text{O}_2) = \frac{0.9}{2.5} \times 1.21 = 0.4356 \text{ (MPa)}$

AND $p(\text{NO}_2) = \frac{1.2}{2.5} \times 1.21 = 0.5808 \text{ (MPa)} \checkmark$

K_p value

$$K_p = \frac{0.5808^2}{0.1936^2 \times 0.4356} = 20.7 \text{ to 3 SF (MPa}^{-1}\text{)} \checkmark$$

Question			Answer				Marks	AO element	Guidance																				
	(c)	(ii)	<table border="1"> <thead> <tr> <th>Change</th> <th>K_p</th> <th>Equilibrium amount of NO_2</th> <th>Initial rate</th> </tr> </thead> <tbody> <tr> <td>Temperature increased</td> <td>smaller</td> <td>smaller</td> <td>greater</td> </tr> <tr> <td>Pressure increase</td> <td>same</td> <td>greater</td> <td>greater</td> </tr> <tr> <td>Catalyst added</td> <td>same</td> <td>same</td> <td>greater</td> </tr> <tr> <td></td> <td>✓</td> <td>✓</td> <td>✓</td> </tr> </tbody> </table>				Change	K_p	Equilibrium amount of NO_2	Initial rate	Temperature increased	smaller	smaller	greater	Pressure increase	same	greater	greater	Catalyst added	same	same	greater		✓	✓	✓	3	AO1.2 ×3	<p>Mark by COLUMN</p> <p>ALLOW obvious alternatives for greater/smaller/same, e.g. increases/decreases/more/less</p>
Change	K_p	Equilibrium amount of NO_2	Initial rate																										
Temperature increased	smaller	smaller	greater																										
Pressure increase	same	greater	greater																										
Catalyst added	same	same	greater																										
	✓	✓	✓																										
			Total				11																						

Question			Answer	Marks	AO element	Guidance
21	(a)	(i)	<p>(Expt 1 and 2) $[S_2O_3^{2-}]$ halves, ($[H^+]$ constant), AND rate halves AND first order (with respect to $[S_2O_3^{2-}]$) ✓</p> <p>(Expt 2 and 3) $[S_2O_3^{2-}]$ quarter AND $[H^+]$ halves, AND rate quarters AND zero order (with respect to $[H^+]$) ✓</p>	2	AO3.1 ×2	<p>ALLOW ORA i.e. (Expt 2 and 1) $[S_2O_3^{2-}]$ doubles, ($[H^+]$ constant), AND rate doubles AND first order with respect to $[S_2O_3^{2-}]$</p> <p>ALLOW comparison of Expt 1 and 3: $[S_2O_3^{2-}] \times 1/8$ AND $[H^+]$ halves, AND rate $\times 1/8$ AND zero order with respect to $[H^+]$</p>
	(a)	(ii)	<p>$S_2O_3^{2-}$ as only reactant species in step 1 ✓</p> <p>Rest of mechanism correct ✓</p>	2	AO3.2 ×2	<p>Step 1: $S_2O_3^{2-} \rightarrow S + SO_3^{2-}$ Step 2: $SO_3^{2-} + 2H^+ \rightarrow SO_2 + H_2O$ OR Step 1: $S_2O_3^{2-} \rightarrow SO_2 + SO^{2-}$ Step 2: $SO^{2-} + 2H^+ \rightarrow S + H_2O$</p> <p>Check with Team Leader for other equations</p>
	(b)	(i)	<p>Gradient gradient in range of -5700 to -6100 ✓</p> <p>E_a calculation $E_a = (-)$ gradient $\times 8.314$ e.g. from -5900, $E_a = (+)$ 49052.6 ($J mol^{-1}$) ✓</p> <p>E_a to 3SF and in $kJ mol^{-1}$ ✓ e.g. 49.1 ($kJ mol^{-1}$)</p>	3	AO2.8 ×3	<p>FULL ANNOTATIONS MUST BE USED ----- Marks are for intermediate calculations</p> <p>ALLOW ECF from an incorrect gradient</p> <p>ALLOW ECF on missing $\times 10^{-3}$, e.g. ALLOW 2 marks for: gradient = -5.9, leading to $E_a = 49.0526$ ($J mol^{-1}$) AND 0.0491 ($kJ mol^{-1}$)</p> <p>DO NOT ALLOW a negative E_a</p>

Question			Answer	Marks	AO element	Guidance
	(b)	(ii)	ln A is intercept at 0 when 1/T OR x axis is 0 ✓	1	AO3.2	
		(iii)	<i>ln k</i> $\ln k = -2.59$ ✓ <i>Temperature</i> $1/T = 3.10 \times 10^{-3} \text{ (s}^{-1}\text{)}$ $T = 49.6 \text{ }^\circ\text{C}$ ✓	2	AO3.1 AO3.2	Correct T scores 2 marks ALLOW ECF for 1/T from incorrect lnK shown on the graph ALLOW in the range $1/T = 3.09 - 3.11 \text{ (} \times 10^{-3} \text{ s}^{-1}\text{)}$ $T = 48.5 \text{ to } 50.6 \text{ }^\circ\text{C}$ ALLOW $T = 50 \text{ }^\circ\text{C}$
			Total	10		

Question		Answer	Marks	AO element	Guidance
22	(a)	<p>FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 13.15 award 2 marks</p> <p>-----</p> $[\text{H}^+] = \frac{1.00 \times 10^{-14}}{0.140} = 7.14 \dots \times 10^{-14} \text{ (mol) } \checkmark$ $\text{pH} = -\log (7.14 \dots \times 10^{-14}) = 13.15 \checkmark$ <p style="text-align: center;">2 DP required</p>	2	AO2.2 ×2	<p>ALLOW ECF providing $\text{pH} > 7$</p> <p>Calculator: $7.142857143 \times 10^{-14}$</p> <p>ALLOW pOH method $\text{pOH} = -\log(0.14) = 0.85 \dots \dots \dots \checkmark$</p> $\text{pH} = 14.00 - (0.85 \dots \dots) = 13.15 \checkmark$
	(b)	(i)			
		$n(\text{H}_2\text{SO}_4) = 1.60 \times \frac{25.0}{1000} = 0.04(00) \text{ (mol)}$ <p>AND</p> $n(\text{NaOH}) = 1.50 \times \frac{55.0}{1000} = 0.0825 \text{ (mol) } \checkmark$ <p>0.04(00) mol H_2SO_4 reacts with 0.08(00) mol NaOH OR 1 mol H_2SO_4 reacts with 2 mol NaOH \checkmark</p>	2	AO2.2 ×2	<p>ALLOW $0.0825 > 0.08$</p>

Question		Answer	Marks	AO element	Guidance
	(b) (ii)	$q = mc\Delta T = 80.0 \times 4.18 \times 13.0$ $= 4347.2 \text{ (J) OR } 4.3472 \text{ (kJ) } \checkmark$ $\Delta H_1 = (-) \frac{4.3472}{0.0400} = (-)108.68 \text{ kJ mol}^{-1} \checkmark$ $\Delta_{\text{neut}}H = (-) \frac{108.68}{2} = (-)54.34 \text{ kJ mol}^{-1} \checkmark$ – sign for ΔH value(s) \checkmark	4	AO2.4 ×4	FULL ANNOTATIONS MUST BE USED ----- ALLOW 3 SF up to calculated answer throughout ALLOW ECF from q DO NOT ALLOW division by $n(\text{NaOH})$ ALLOW $\Delta_{\text{neut}}H$ from $\Delta H_1 / 2$ ALLOW alternative methods
	(b) (iii)	The same OR $13^\circ\text{C} \checkmark$ (Double the moles so) double the energy is spread over double the volume	2	AO3.1 ×2	ALLOW explanation that uses a calculation based on moles, volumes ALLOW mass for volume

Question		Answer	Marks	AO element	Guidance
	(c)*	<p><i>Please refer to the marking instructions on page 4 of this mark scheme for guidance on how to mark this question.</i></p> <p>Level 3 (5–6 marks) Reaches a comprehensive conclusion with most detail and few errors for the formation of the buffer AND Calculation of the correct buffer pH AND Correct mass of N₂O₃. <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p>Level 2 (3–4 marks) Reaches a sound conclusion with some detail and some errors for Formation of buffer AND Calculation of the buffer pH OR Formation of buffer AND Mass of N₂O₃. OR Calculation of the buffer pH AND Mass of N₂O₃. OR Partial explanations of formation of the buffer AND buffer pH AND Mass of N₂O₃. <i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p>Level 1 (1–2 marks) Attempts, with some success, to: Describe formation of buffer OR Calculate buffer pH OR Obtain mass of N₂O₃. <i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p>0 marks No response or no response worthy of credit.</p>	6	<p>AO1.2 ×2</p> <p>AO2.6 ×2</p> <p>AO3.1 ×2</p>	<p>Indicative scientific points may include:</p> <p>1. Formation of buffer</p> <ul style="list-style-type: none"> Acid / HNO₂ is in excess HNO₂ + NaOH → NaNO₂ + H₂O Partial neutralisation of HNO₂ → formation of NO₂⁻/ NaNO₂ Buffer contains HNO₂ AND NO₂⁻/NaNO₂ <p>2. Calculation of buffer pH</p> <ul style="list-style-type: none"> n(HNO₂) added = 0.0500 (mol) n(NaOH) added = 0.0150 (mol) n(NO₂⁻) formed = 0.0150 (mol) n(HNO₂) remaining = 0.0500 – 0.0150 = 0.0350 (mol) K_a = 10^{-3.34} = 4.57... × 10⁻⁴ (mol dm⁻³) Concentrations = mol (volume 1 dm³) [H⁺] = $\frac{4.57... \times 10^{-4} \times 0.0350}{0.0150}$ = 1.0665... × 10⁻³ (mol dm⁻³) pH = 2.97 pH to 2 dec places <p>3. Calculation of mass of N₂O₃</p> <ul style="list-style-type: none"> 1 mol N₂O₃ → 2 mol HNO₂ OR N₂O₃ + H₂O → 2HNO₂ n(HNO₂) = 0.0500 (mol) n(N₂O₃) = 0.0500/2 = 0.0250 (mol) m(N₂O₃) = 0.0250 × 76 = 1.9(0) g
			Total	16	

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored