

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS
A LEVEL**

H433/03

CHEMISTRY B (SALTERS)

Practical skills in chemistry

TUESDAY 27 JUNE 2017: Morning

**TIME ALLOWED: 1 hour 30 minutes
plus your additional time allowance**

MODIFIED ENLARGED

First name		Last name	
-----------------------	--	----------------------	--

Centre number						Candidate number				
--------------------------	--	--	--	--	--	-----------------------------	--	--	--	--

YOU MUST HAVE:

the Insert

the Data Sheet for Chemistry B (Salters)

YOU MAY USE:

a scientific or graphical calculator

READ INSTRUCTIONS OVERLEAF



INSTRUCTIONS

The Insert will be found with this document.

Use black ink. HB pencil may be used for graphs and diagrams only.

The practical insert is needed with this paper.

Complete the boxes on the front page with your name, centre number and candidate number.

Answer ALL the questions.

Write your answer to each question in the space provided. If additional space is required, you should use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.

INFORMATION

The total mark for this paper is 60.

The marks for each question are shown in brackets [].

Quality of extended responses will be assessed in questions marked with an asterisk (*).

BLANK PAGE

Answer ALL the questions.

- 1 A group of students decide to analyse a commercial sweetener.**

The major component is listed as the dipeptide aspartame.


- (a) The students crush a sweetener tablet and dissolve it in 1 cm^3 of water in a test tube. They add a similar volume of 6.0 mol dm^{-3} hydrochloric acid and some anti-bumping granules. They then cautiously boil the contents for about a minute.**

Suggest the purpose of adding the anti-bumping granules.

_____ **[1]**

(b) The students expect the aspartame to have been hydrolysed by the hot acid.

(i) The structure of aspartame is given below with three functional groups (1, 2, 3) ringed.



Name the homologous series for the functional groups 1, 2 and 3.

1 _____

2 _____


3 _____ [1]

- (ii) Functional group 2 AND functional group 3 are hydrolysed under the acid conditions to give three products. The products are methanol, and ions formed by the amino acids phenylalanine and aspartic acid.

The structure of the amino acid phenylalanine is given below.

In the box draw the structure of the ION formed by ASPARTIC ACID. [2]

PHENYLALANINE




structure of ION formed by aspartic acid

- (iii) Amino acids such as phenylalanine exist as 'zwitterions'.

Use the structure of phenylalanine to help explain how amino acids form zwitterions. Draw the structure of the zwitterion of phenylalanine. [2]

PHENYLALANINE



(c)* The students wanted to confirm that the aspartame had been hydrolysed to produce the two amino acids, aspartic acid and phenylalanine.

They decided the best way was to use paper chromatography on the solution formed after the hydrolysis reaction.

Describe how the students could carry out the chromatography experiment and explain how they could use their results to show hydrolysis had taken place.

You may include a diagram in your answer. [6]

[illegible]


- (d) The students decided to investigate carbonated drinks such as diet and regular cola. These drinks contain phosphoric acid (to increase the flavour) and aspartame. Regular cola has a lower concentration of acid and keeps for longer than diet cola.**

Suggest why regular cola keeps for longer.


[2]

- 2 Two students decide to study the reaction of magnesium with dilute hydrochloric acid to find the order of this reaction with respect to the acid.

The equation for the reaction is:



The students use the following apparatus:



- (a) The students run FIVE experiments each using a different concentration of hydrochloric acid.

They use the same length of magnesium ribbon each time.

- (i) Why was the length of the magnesium ribbon kept the same?

[1]

- (ii) 50 cm^3 acid is used each time.
The concentration ranged from 0.1 to 0.5 mol dm^{-3} .

Calculate the **MAXIMUM** mass of magnesium ribbon that would completely react with the acid in **ALL** five experiments.

mass = _____ g [3]

- (iii) Calculate the maximum length of magnesium ribbon needed to completely react with the acid in all the experiments.

The mass per unit length of Mg ribbon
 $= 1.3 \times 10^{-2} \text{ g cm}^{-1}$

length = _____ cm [1]

- (b) The students predict that the reaction will be second order with respect to acid.

Write the rate equation that would be appropriate for this prediction.

_____ [1]


(c) The students plot a graph of volume of hydrogen against time.

The graph of the results when the concentration of hydrochloric acid is 0.5 mol dm^3 is shown below.

Draw a line of best fit on the graph.

Use your line of best fit to calculate the initial rate of reaction at this concentration and give the units.

Show your working on the graph.




initial rate _____ units _____ [5]

- (d) The students decide to measure the activation enthalpy for the reaction. They repeat the experiment at different temperatures using a fixed concentration of acid.

The following table shows the data from their experiments.

$\ln k$	Temperature /°C	Temperature, T/K	$1/T/\text{K}^{-1}$ ($\times 10^{-3}$)
−5.46	15.6	288.6	3.47
−4.92	30.3	303.3	3.30
−4.34	45.0	318.0	3.14
−3.73	65.0	338.0	2.96

Use the data to plot a suitable graph opposite and work out the activation enthalpy for the reaction.




activation enthalpy = _____ kJ mol^{-1} [6]


- 3 A perfume laboratory is investigating the molecules responsible for the smell of some common flowers.

Their investigations suggest that the following two liquid compounds are important in the characteristic smell of hyacinth flowers.

ocimenol



cinnamyl alcohol



- (a) Acidified dichromate solution can be used to distinguish between these two compounds. Only cinnamyl alcohol can be oxidised using acid dichromate.

Explain why this is so, and describe any colour changes you would expect to see.

[2]

(b) The products of the oxidation of cinnamyl alcohol by acid dichromate differ depending on the conditions.

Two experiments are shown below.

Name, in the appropriate boxes below:

the technique used in each set up

the homologous series of the final product of oxidation [4]

Diagram 1: Reflux apparatus. A round-bottom flask containing cinnamyl alcohol and an oxidising agent is heated. A Liebig condenser is attached to the neck of the flask, with cold water flowing through it. The condensed liquid drips into a conical flask.

Labels: cinnamyl alcohol, oxidising agent, heat, cold water, technique used, homologous series of product

Diagram 2: Distillation apparatus. A round-bottom flask containing oxidising agent + cinnamyl alcohol is heated. A vertical thermometer is inserted into the flask. A vertical condenser is attached to the top of the flask, with cold water flowing through it. The condensed liquid drips out of the side of the condenser.

Labels: cold water, oxidising agent + cinnamyl alcohol, heat, technique used, homologous series of product


- (c) Ocimenol and cinnamyl alcohol are both unsaturated molecules.

Explain how you could use a solution of bromine water to show which of the molecules has the greater degree of unsaturation.


[4]

- (d) Both ocimenol and cinnamyl alcohol molecules show stereoisomerism.

ocimenol



cinnamyl alcohol




Name the TYPE of stereoisomerism shown by these molecules.

Explain how this isomerism arises.

The structure of one of the stereoisomers of cinnamyl alcohol is shown in the left hand box below; draw the structure of the other isomer in the right hand box. [3]

TYPE of stereoisomerism _____

Explanation



- (e) Mass spectrometry can also be used to distinguish between ocimenol and cinnamyl alcohol.

The molecular ion peak in a mass spectrum of ocimenol is at an m/z value of 154.

What m/z value would represent the molecular ion peak in a mass spectrum of cinnamyl alcohol?

m/z for cinnamyl alcohol molecular ion

= _____ [1]

This question refers to the Practical Insert that is provided as an insert to this paper.

- 4 (a) (i) Name the piece of apparatus which is most suitable for removing the 25.0 cm³ of the Cu²⁺ ion solution.**

_____ [1]

- (ii) Use the student's results in the Practical Insert to work out an average titre that the student should use in calculating the amount of Cu²⁺ in the 'coin solution'.**

average titre = _____ cm³ [1]

- (iii) Use your answer from (a)(ii) to calculate the percentage by mass of copper in the coin. Give your answer to an APPROPRIATE number of significant figures.**

percentage by mass of copper in coin

= _____ % [4]

- (iv) The student considers the uncertainties of measurement in the experiment.

The percentage measurement uncertainty marked on the apparatus used to transfer the 25 cm^3 of the 'copper' solution is $\pm 0.24\%$.
The volumetric flask is marked $\pm 0.08\%$.

Calculate the measurement uncertainties of the other pieces of apparatus used.

Which piece of apparatus contributes most to the measurement uncertainty in this experiment? Use the space below. [2]

- (v) Name another method that the student could use to find the concentration of Cu^{2+} ions in the 'coin solution'.

_____ [1]

(b)* The student's method requires a standard solution of sodium thiosulfate for the titration.

Describe the techniques and procedures used to prepare a standard solution of a specific concentration from a solid. [6]

[illegible]

END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

BLANK PAGE

BLANK PAGE

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.