



RECOGNISING ACHIEVEMENT

ADVANCED SUBSIDIARY GCE

MATHEMATICS (MEI)

Mechanics 1

4761

Candidates answer on the Answer Booklet

OCR Supplied Materials:

- 8 page Answer Booklet
- Graph paper
- MEI Examination Formulae and Tables (MF2)

Other Materials Required:

None

Thursday 11 June 2009

Morning

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- This document consists of **8** pages. Any blank pages are indicated.

Section A (36 marks)

1 The velocity-time graph shown in Fig. 1 represents the straight line motion of a toy car. All the lines on the graph are straight.




Fig. 1

The car starts at the point A at $t = 0$ and in the next 8 seconds moves to a point B.

(i) Find the distance from A to B. [2]

T seconds after leaving A, the car is at a point C which is a distance of 10 m from B.

(ii) Find the value of T . [3]

(iii) Find the displacement from A to C. [1]

2 A small box has weight \mathbf{W} N and is held in equilibrium by two strings with tensions \mathbf{T}_1 N and \mathbf{T}_2 N. This situation is shown in Fig. 2 which also shows the standard unit vectors \mathbf{i} and \mathbf{j} that are horizontal and vertically upwards, respectively.




Fig. 2

The tension \mathbf{T}_1 is $10\mathbf{i} + 24\mathbf{j}$.

(i) Calculate the magnitude of \mathbf{T}_1 and the angle between \mathbf{T}_1 and the vertical. [3]

The magnitude of the weight is w N.

(ii) Write down the vector \mathbf{W} in terms of w and \mathbf{j} . [1]

The tension \mathbf{T}_2 is $k\mathbf{i} + 10\mathbf{j}$, where k is a scalar.

(iii) Find the values of k and of w . [3]

3 Fig. 3 is a sketch of the velocity-time graph modelling the velocity of a sprinter at the start of a race.




Fig. 3

(i) How can you tell from the sketch that the acceleration is not modelled as being constant for $0 \leq t \leq 4$? [1]

The velocity of the sprinter, $v \text{ m s}^{-1}$, for the time interval $0 \leq t \leq 4$ is modelled by the expression

$$v = 3t - \frac{3}{8}t^2.$$

(ii) Find the acceleration that the model predicts for $t = 4$ and comment on what this suggests about the running of the sprinter. [3]

(iii) Calculate the distance run by the sprinter from $t = 1$ to $t = 4$. [4]

4 Fig. 4 shows a particle projected over horizontal ground from a point O at ground level. The particle initially has a speed of 32 m s^{-1} at an angle α to the horizontal. The particle is a horizontal distance of 44.8 m from O after 5 seconds. Air resistance should be neglected.




Fig. 4

(i) Write down an expression, in terms of α and t , for the horizontal distance of the particle from O at time t seconds after it is projected. [1]

(ii) Show that $\cos \alpha = 0.28$. [2]

(iii) Calculate the greatest height reached by the particle. [4]

5 The position vector of a toy boat of mass 1.5 kg is modelled as $\mathbf{r} = (2 + t)\mathbf{i} + (3t - t^2)\mathbf{j}$ where lengths are in metres, t is the time in seconds, \mathbf{i} and \mathbf{j} are horizontal, perpendicular unit vectors and the origin is O.

(i) Find the velocity of the boat when $t = 4$. [3]

(ii) Find the acceleration of the boat and the horizontal force acting on the boat. [3]

(iii) Find the cartesian equation of the path of the boat referred to x - and y -axes in the directions of \mathbf{i} and \mathbf{j} , respectively, with origin O. You are not required to simplify your answer. [2]

Section B (36 marks)

6 An empty open box of mass 4 kg is on a plane that is inclined at 25° to the horizontal.

In one model the plane is taken to be smooth.

The box is held in equilibrium by a string with tension T N parallel to the plane, as shown in Fig. 6.1.




Fig. 6.1

(i) Calculate T .

[2]

A rock of mass m kg is now put in the box. The system is in equilibrium when the tension in the string, still parallel to the plane, is 50 N.

(ii) Find m .

[3]

In a refined model the plane is rough.

The empty box, of mass 4 kg, is in equilibrium when a frictional force of 20 N acts down the plane and the string has a tension of P N inclined at 15° to the plane, as shown in Fig. 6.2.




Fig. 6.2

(iii) Draw a diagram showing all the forces acting on the box.

[2]

(iv) Calculate P .

[5]

(v) Calculate the normal reaction of the plane on the box.

[4]

7 A box of mass 8 kg slides on a horizontal table against a constant resistance of 11.2 N.

(i) What horizontal force is applied to the box if it is sliding with acceleration of magnitude 2 m s^{-2} ? [3]

Fig. 7 shows the box of mass 8 kg on a long, rough, horizontal table. A sphere of mass 6 kg is attached to the box by means of a light inextensible string that passes over a smooth pulley. The section of the string between the pulley and the box is parallel to the table. The constant frictional force of 11.2 N opposes the motion of the box. A force of 105 N parallel to the table acts on the box in the direction shown, and the acceleration of the system is in that direction.




Fig. 7

(ii) What information in the question indicates that while the string is taut the box and sphere have the same acceleration? [1]

(iii) Draw two separate diagrams, one showing all the horizontal forces acting on the box and the other showing all the forces acting on the sphere. [2]

(iv) Show that the magnitude of the acceleration of the system is 2.5 m s^{-2} and find the tension in the string. [7]

The system is stationary when the sphere is at point P. When the sphere is 1.8 m above P the string breaks, leaving the sphere moving upwards at a speed of 3 m s^{-1} .

(v) (A) Write down the value of the acceleration of the sphere after the string breaks. [1]

(B) The sphere passes through P again at time T seconds after the string breaks. Show that T is the positive root of the equation $4.9T^2 - 3T - 1.8 = 0$. [2]

(C) Using part (B), or otherwise, calculate the total time that elapses after the sphere moves from P before the sphere again passes through P. [4]

BLANK PAGE



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations, is given to all schools that receive assessment material and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity. For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1PB.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.