



ADVANCED GCE

MATHEMATICS (MEI)

Further Methods for Advanced Mathematics (FP2)

4756

Candidates answer on the Answer Booklet

OCR Supplied Materials:

- 8 page Answer Booklet
- MEI Examination Formulae and Tables (MF2)

Other Materials Required:

None

Monday 11 January 2010

Morning

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer **all** the questions in Section A and **one** question from Section B.
- Do **not** write in the bar codes.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- This document consists of **4** pages. Any blank pages are indicated.

Section A (54 marks)**Answer all the questions**

1 (a) Given that $y = \arctan \sqrt{x}$, find $\frac{dy}{dx}$, giving your answer in terms of x . Hence show that

$$\int_0^1 \frac{1}{\sqrt{x}(x+1)} dx = \frac{\pi}{2}. \quad [6]$$

(b) A curve has cartesian equation

$$x^2 + y^2 = xy + 1.$$

(i) Show that the polar equation of the curve is

$$r^2 = \frac{2}{2 - \sin 2\theta}. \quad [4]$$

(ii) Determine the greatest and least positive values of r and the values of θ between 0 and 2π for which they occur. [6]

(iii) Sketch the curve. [2]

2 (a) Use de Moivre's theorem to find the constants a, b, c in the identity

$$\cos 5\theta \equiv a \cos^5 \theta + b \cos^3 \theta + c \cos \theta. \quad [6]$$

(b) Let

$$C = \cos \theta + \cos \left(\theta + \frac{2\pi}{n} \right) + \cos \left(\theta + \frac{4\pi}{n} \right) + \dots + \cos \left(\theta + \frac{(2n-2)\pi}{n} \right),$$

$$\text{and } S = \sin \theta + \sin \left(\theta + \frac{2\pi}{n} \right) + \sin \left(\theta + \frac{4\pi}{n} \right) + \dots + \sin \left(\theta + \frac{(2n-2)\pi}{n} \right),$$

where n is an integer greater than 1.

By considering $C + jS$, show that $C = 0$ and $S = 0$. [7]

(c) Write down the Maclaurin series for e^t as far as the term in t^2 .

Hence show that, for t close to zero,

$$\frac{t}{e^t - 1} \approx 1 - \frac{1}{2}t. \quad [5]$$

3 (i) Find the inverse of the matrix

$$\begin{pmatrix} 1 & 1 & a \\ 2 & -1 & 2 \\ 3 & -2 & 2 \end{pmatrix}$$

where $a \neq 4$.

Show that when $a = -1$ the inverse is

$$\frac{1}{5} \begin{pmatrix} 2 & 0 & 1 \\ 2 & 5 & -4 \\ -1 & 5 & -3 \end{pmatrix}. \quad [6]$$

(ii) Solve, in terms of b , the following system of equations. [5]

$$\begin{aligned} x + y - z &= -2 \\ 2x - y + 2z &= b \\ 3x - 2y + 2z &= 1 \end{aligned}$$

(iii) Find the value of b for which the equations

$$\begin{aligned} x + y + 4z &= -2 \\ 2x - y + 2z &= b \\ 3x - 2y + 2z &= 1 \end{aligned}$$

have solutions. Give a geometrical interpretation of the solutions in this case. [7]

Section B (18 marks)

Answer one question

Option 1: Hyperbolic functions

4 (i) Prove, using exponential functions, that

$$\cosh 2x = 1 + 2 \sinh^2 x.$$

Differentiate this result to obtain a formula for $\sinh 2x$. [4]

(ii) Solve the equation

$$2 \cosh 2x + 3 \sinh x = 3,$$

expressing your answers in exact logarithmic form. [7]

(iii) Given that $\cosh t = \frac{5}{4}$, show by using exponential functions that $t = \pm \ln 2$.

Find the exact value of the integral

$$\int_4^5 \frac{1}{\sqrt{x^2 - 16}} dx. \quad [7]$$

Option 2: Investigation of curves

This question requires the use of a graphical calculator.

5 A line PQ is of length k (where $k > 1$) and it passes through the point $(1, 0)$. PQ is inclined at angle θ to the positive x -axis. The end Q moves along the y -axis. See Fig. 5. The end P traces out a locus.




Fig. 5

(i) Show that the locus of P may be expressed parametrically as follows. [3]

$$x = k \cos \theta \quad y = k \sin \theta - \tan \theta$$

You are now required to investigate curves with these parametric equations, where k may take any non-zero value and $-\frac{1}{2}\pi < \theta < \frac{1}{2}\pi$.

(ii) Use your calculator to sketch the curve in each of the cases $k = 2$, $k = 1$, $k = \frac{1}{2}$ and $k = -1$. [4]

(iii) For what value(s) of k does the curve have

- (A) an asymptote (you should state what the asymptote is),
- (B) a cusp,
- (C) a loop? [3]

(iv) For the case $k = 2$, find the angle at which the curve crosses itself. [2]

(v) For the case $k = 8$, find in an exact form the coordinates of the highest point on the loop. [3]

(vi) Verify that the cartesian equation of the curve is

$$y^2 = \frac{(x-1)^2}{x^2} (k^2 - x^2). \quad [3]$$