

## **Mathematics (MEI)**

Advanced Subsidiary GCE

**Unit 4752: Concepts for Advanced Mathematics**

## **Mark Scheme for January 2011**

---

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552610  
E-mail: publications@ocr.org.uk

4752

Mark Scheme

January 2011

SECTION A

1	11.4 o.e.	2	<b>M1</b> for $12/3 + 12/4 + 12/5 + 12/6$ o.e.	<b>M0</b> unless four terms summed
2	$\frac{1}{2}x^6 + 4x^{\frac{1}{2}} + c$	4	<b>B1</b> for $\frac{1}{2}x^6$ , <b>M1</b> for $kx^{\frac{1}{2}}$ , <b>A1</b> for $k = 4$ $\frac{4}{2}$ or $\frac{1}{2}$ , <b>B1</b> for $+ c$ dependent on at least one power increased	$\frac{3}{6}$ allow $\frac{1}{6}x^6$ isw,
3	$\frac{1}{2} \times 1.5 \times (0.6 + 0.7 + 2(2.3+3.1+2.8+1.8))$  $= 15.975$ rounded to 2 s.f. or more	M2  A1	<b>M1</b> if one error or <b>M2</b> for sum of 5 unsimplified individual trapezia: 2.175, 4.05, 4.425, 3.45, 1.875	basic shape of formula must be correct. Must be 5 strips. <b>M0</b> if pair of brackets omitted or $h = 7.5$ or 1. allow recovery of brackets omitted to obtain correct answer. <b>M0</b> for other than 5 trapezia isw only if 15.975 clearly identified as cross-sectional area
4	(i) (3, 15)	B2	<b>B1</b> for each coordinate	s.c. <b>B0</b> for (3, 5)
4	(ii) (1.5, 5)	B2	<b>B1</b> for each coordinate	s.c. <b>B0</b> for (3, 5)
5	$ar = 6$ and $ar^4 = -48$ $r = -2$ tenth term = 1536  $\frac{-3(1-(-2)^n)}{1-(-2)}$ o.e.  $(-2)^n - 1$	M1  M1  A1  M1  A1	<b>B2</b> for $r = -2$ www  <b>B3</b> for 1536 www  allow <b>M1</b> for $a = 6 \div$ their $r$ and substitution in GP formula with their $a$ and $r$  c.a.o.	ignore incorrect lettering such as $d = -2$  condone the omission of the brackets round “-2” in the numerator and / or the denominator

4752

Mark Scheme

January 2011

6	$a+2d=24$ and $a + 9d=3$ $d = -3; a = 30$ $S_{50} - S_{20}$ $-2205$ cao	<b>M1</b> <b>A1</b> <b>A1</b> <b>M1</b> <b>A1</b>	if <b>M0</b> , <b>B2</b> for either, <b>B3</b> for both ft their $a$ and $d$ ; <b>M1</b> for $S_{30} = \frac{30}{2}(u_{21} + u_{50})$ o.e. <b>B2</b> for -2205 www	do not award <b>B2</b> or <b>B3</b> if values clearly obtained fortuitously $S_{50} = -2175; S_{20} = 30$ $u_{21} = 30 - 20 \times 3 = -30$ $u_{50} = 30 - 49 \times 3 = -117$
7	(i) $17 \log_{10} x$ or $\log_{10} x^{17}$	<b>B2</b>	<b>M1</b> for $5\log_{10} x$ or $12 \log_{10} x$ or $\log_{10} x^{12}$ as part of the first step	condone omission of base
7	(ii) $-b$	<b>B2</b>	<b>M1</b> for $\log_a 1 = 0$ or $\log_a a = 1$ soi	allow $0 - b$
8	substitution of $\sin^2 \theta = 1 - \cos^2 \theta$ $-5\cos^2 \theta = \cos \theta$ $\theta = 90$ and $270$ , $102$ $258$ $101$ and $259$	<b>M1</b> <b>A1</b> <b>A1</b> <b>A1</b> <b>A1</b> <b>SC</b> <b>1</b>	soi or better accept 101.5(...) and 258.(46...) rounded to 3 or more sf; if <b>M0</b> , allow <b>B1</b> for both of 90 and 270 and <b>B1</b> for 102 and <b>B1</b> for 258 (to 3 or more sf)	if the 4 correct values are presented, ignore any extra values which are outside the required range, but apply a penalty of minus 1 for extra values in the range if given in radians deduct 1 mark from total awarded (1.57, 1.77, 4.51, 4.71)

4752

Mark Scheme

January 2011

9	area sector = $\frac{1}{2} \times r^2 \times \frac{\pi}{6} \left[ = \frac{\pi r^2}{12} \right]$	<b>M1</b>	soi	
	area triangle = $\frac{1}{2} \times a^2 \times \sin \frac{\pi}{6} \left[ = \frac{a^2}{4} \right]$	<b>M1</b>	soi	allow sin30
	$\frac{1}{2}a^2 \times \frac{1}{2} = \frac{1}{2} \times r^2 \times \frac{\pi}{6} \times \frac{1}{2}$	<b>M1</b>	soi	no follow through marks available
	$\frac{a^2}{4} = \frac{\pi r^2}{24}$ o.e. and completion to given answer	<b>A1</b>		at least one correct intermediate step required, and no wrong working to obtain given answer

Section A Total: 36

4752

Mark Scheme

January 2011

## SECTION B

10	(i) eqn of AB is $y = 3x + 1$ o.e. their “ $3x + 1$ ” = $4x^2$ $(4x + 1)(x - 1) = 0$ o.e. so $x = -1/4$ at C, $x = -1/4$ , $y = 4 \times (-1/4)^2$ or $3 \times (-1/4) + 1$ [=1/4 as required]	<b>M1</b> <b>M1</b> <b>M1</b> <b>A1</b>	or equiv in $y$ : $y = 4 \left( \frac{y-1}{3} \right)^2$ or rearranging and deriving roots $y = 4$ or $\frac{1}{4}$ condone verification by showing lhs = rhs o.e. or $y = \frac{1}{4}$ implies $x = \pm \frac{1}{4}$ so at C $x = -\frac{1}{4}$	<b>SC3</b> for verifying that A, B and C are collinear and that C also lies on the curve <b>SC2</b> for verifying that A, B and C are collinear by showing that gradient of AB = AC (for example) or showing C lies on AB solely verifying that C lies on the curve scores 0
10	(ii) $y' = 8x$ at A $y' = 8$ eqn of tgt at A $y - 4 = \text{their } "8"(x - 1)$ $y = 8x - 4$  at C $y' = 8 \times -1/4$ [=−2] $y - \frac{1}{4} = -2(x - (-\frac{1}{4}))$ or other unsimplified equivalent to obtain given result. allow correct verification that $(-\frac{1}{4}, \frac{1}{4})$ lies on given line	<b>M1</b> <b>A1</b> <b>M1</b> <b>A1</b> <b>M1</b> <b>A1</b>	ft their gradient  NB if $m = -2$ obtained from given answer or only showing that $(-\frac{1}{4}, \frac{1}{4})$ lies on given line $y = -2x - \frac{1}{4}$ then 0 marks.	 gradient must follow from evaluation of $\frac{dy}{dx}$ condone unsimplified versions of $y = 8x - 4$  dependent on award of first <b>M1</b> <b>SC2</b> if equation of tangent and curve solved simultaneously to correctly show repeated root
10	(iii) their “ $8x - 4$ ” = $-2x - \frac{1}{4}$ $y = -1$ www	<b>M1</b> <b>A1</b>	or $\frac{y+4}{8} = \frac{y+\frac{1}{4}}{-2}$	o.e. $[x = 3/8]$

4752

## Mark Scheme

January 2011

11	(i) $\frac{x^4}{4} - x^3 - \frac{x^2}{2} + 3x$ their integral at 3 – their integral at 1 [= $-2.25 - 1.75$ ]  $= -4$ isw represents area between curve and $x$ axis between $x = 1$ and 3  negative since below $x$ -axis	M2  <b>M1</b>  <b>A1</b>  <b>B1</b>  <b>B1</b>	<b>M1</b> if at least two terms correct  <b>M1</b> dependent on integration attempted  <b>A1</b>  <b>B1</b>  <b>B1</b>	ignore $+ c$  M0 for evaluation of $x^3 - 3x^2 - x + 3$ or of differentiated version  <b>B0</b> for area <i>under</i> or above curve between $x = 1$ and 3
11	(ii) $y' = 3x^2 - 6x - 1$ their $y' = 0$ soi  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ with $a = 3$ , $b = -6$ and $c = -1$ isw  $x = \frac{6 \pm \sqrt{48}}{6}$ or better as final answer  $\frac{6 - \sqrt{48}}{6} < x < \frac{6 + \sqrt{48}}{6}$ or ft their final answer	<b>M1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>  <b>B1</b>	  <b>M1</b> dependent on differentiation attempted  <b>M1</b> or $3(x - 1)^2 - 4 [= 0]$ or better  <b>A1</b> eg <b>A1</b> for $1 \pm \frac{2}{3}\sqrt{3}$  <b>B1</b> allow $\leq$ instead of $<$	       <b>A0</b> for incorrect simplification, eg $1 \pm \sqrt{48}$  allow <b>B1</b> if <i>both</i> inequalities are stated separately and it's clear that both apply allow <b>B1</b> if the terms and the signs are in reverse order
12	(i) 50% of 25 000 is 12 500 and the population [in 2005] is 12 000 [so consistent]	<b>B1</b>	or 12 000 is 48% of 25 000 so less than 50% [so consistent]	
12	(ii) $\log_{10} P = \log_{10} a - kt$ or  $\log_{10} \frac{P}{a} = -kt$ o.e. www	<b>B2</b>	condone omission of base; <b>M1</b> for $\log_{10} P = \log_{10} a + \log_{10} 10^{-kt}$ or better www	

4752

Mark Scheme

January 2011

12	(iii) 4.27, 4.21, 4.13, 4.08  plots ruled line of best fit drawn	<b>B1</b> <b>B1</b> <b>B1</b>	accept 4.273..., 4.2108..., 4.130..., 4.079... rounded to 2 or more dp 1 mm tolerance ft their values if at least 4 correct values are correctly plotted	f.t. if at least two calculated values correct must have at least one point on or above and at least one point on or below the line and must cover $0 \leq t \leq 25$
12	(iv) $a = 25000$ to $25400$  $0.01 \leq k \leq 0.014$  $P = a \times 10^{-kt}$ or $P = 10^{\log a - kt}$ with values in acceptable ranges	<b>B1</b>  <b>B2</b>  <b>B1</b>	allow $10^{4.4..}$  <b>M1</b> for $-k = \frac{y}{x}$ using values from table or graph; condone $+k$  <b>B0</b> if left in logarithmic form	<b>M1</b> for a correct first step in solving a pair of valid equations in either form <b>A1</b> for $k$ <b>A1</b> for $a$ <b>A1</b> for $P = a \times 10^{-kt}$
12	(v) $P = a \times 10^{-35k}$  8600 to 9000  comparing their value with 9375 o.e. and reaching the correct conclusion for their value	<b>M1</b>  <b>A1</b>  <b>A1</b>	T heir $a$ and $k$  f.t.	allow $\log P = \log a - 35k$

Section B Total: 36

**OCR (Oxford Cambridge and RSA Examinations)**  
1 Hills Road  
Cambridge  
CB1 2EU

**OCR Customer Contact Centre**

**14 – 19 Qualifications (General)**

Telephone: 01223 553998  
Facsimile: 01223 552627  
Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**www.ocr.org.uk**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
is a Company Limited by Guarantee  
Registered in England  
Registered Office: 1 Hills Road, Cambridge, CB1 2EU  
Registered Company Number: 3484466  
OCR is an exempt Charity

**OCR (Oxford Cambridge and RSA Examinations)**  
Head office  
Telephone: 01223 552552  
Facsimile: 01223 552553

