



GCE

Mathematics (MEI)

Advanced GCE

Unit 4768: Statistics 3

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Q1				
(i)	<p>t test might be used because</p> <ul style="list-style-type: none"> population variance is unknown background population is Normal 	E1 E1	Allow “sample is small” as an alternative.	2
(ii)	<p>$H_0: \mu = 15.3$ $H_1: \mu < 15.3$</p> <p>where μ is the mean of Gerry’s times.</p> $\bar{x} = 14.987 \quad s_{n-1} = 0.4567(5)$ <p>Test statistic is $\frac{14.987 - 15.3}{0.45675} \quad \sqrt{10}$</p> $= -2.167(0).$ <p>Refer to t_9. Single-tailed 5% point is -1.833.</p> <p>Significant. Seems that Gerry’s times have been reduced on average.</p>	B1 B1 B1 M1 A1 M1 A1 A1 A1	<p>Both hypotheses. Hypotheses in words only must include “population”. Do NOT allow “$\bar{X} = \dots$” or similar unless \bar{X} is clearly and explicitly stated to be a <u>population</u> mean.</p> <p>For adequate verbal definition. Allow absence of “population” if correct notation μ is used.</p> <p>$s_n = 0.4333$ but do <u>NOT</u> allow this here or in construction of test statistic, but FT from there.</p> <p>Allow c’s \bar{x} and/or s_{n-1}. Allow alternative: $15.3 + (c's - 1.833) \times \frac{0.45675}{\sqrt{10}}$ ($= 15.035$) for subsequent comparison with \bar{x}. (Or $\bar{x} - (c's - 1.833) \times \frac{0.45675}{\sqrt{10}}$ $= 15.252$) for comparison with 15.3.) c.a.o. but ft from here in any case if wrong. Use of $\mu - \bar{x}$ scores M1A0, but ft.</p> <p>No ft from here if wrong. Must be minus 1.833 unless absolute values are being compared. No ft from here if wrong. $P(t < -2.167(0)) = 0.0292$. ft only c’s test statistic. ft only c’s test statistic. Conclusion in context to include “average” o.e.</p>	9
(iii)	<p>A 5% significance level means that the probability of rejecting H_0 given that it is true is 0.05.</p> <p>Decreasing the significance level would make it less likely that a true H_0 would be rejected.</p> <p>Evidence for rejecting H_0 would need to be stronger.</p>	E1 E1 E1	Or equivalent. Allow answers that relate to the context of the question.	3
(iv)	<p>CI is given by $14.987 \pm$</p> $\times \frac{0.45675}{\sqrt{10}}$ $= 14.987 \pm 0.3267 = (14.66(0), 15.31(3))$	M1 B1 M1 A1	ZERO/4 if not same distribution as test. Same wrong distribution scores maximum M1B0M1A0. Recovery to t_9 is OK. c.a.o. Must be expressed as an interval.	4
				18

Q3				
(i) (A)		M1 A1 A1	Increasing curve, through (0, 0), in first quadrant only. Asymptotic behaviour. Asymptote labelled; condone absence of axis labels.	3
(B)	<p>For the UQ $G(u) = 0.75$ $\therefore \left(1 + \frac{u}{200}\right)^{-2} = \frac{1}{4} \quad \therefore u = 200$</p> <p>For the LQ $G(l) = 0.25$ $\therefore \left(1 + \frac{l}{200}\right)^{-2} = \frac{3}{4} \quad \therefore l = 200\left(\frac{2}{\sqrt{3}} - 1\right) = 30.94\dots$</p> <p>$\therefore \text{IQR} = 200 - 30.94 = 169.06$</p> <p>For an outlier $x > \text{UQ} + 1.5 \times \text{IQR} = 200 + 1.5 \times 169 = 453.58 \approx 454$ (nearest hour)</p>	M1 A1 A1 M1 M1 E1	<p>Use of $G(x)$ for either quartile. c.a.o.</p> <p>c.a.o.</p> <p>UQ – LQ UQ + 1.5 × IQR. Answer given; must be obtained genuinely.</p>	6
(ii) (A)	$F(x) = \int_0^x \frac{1}{200} e^{-\frac{t}{200}} dt$ $= \left[-e^{-\frac{t}{200}} \right]_0^x = \left(-e^{-\frac{x}{200}} \right) - \left(-e^0 \right) = 1 - e^{-\frac{x}{200}}$	M1 A1 E1	<p>Correct integral, including limits (which may be implied subsequently).</p> <p>Correctly integrated.</p> <p>Limits used. Answer given; must be shown convincingly. Condone the omission of $x < 0$ part.</p> <p>Allow use of “+ c” with $F(0) = 0$.</p>	3
(B)	$P(X > 50) = 1 - F(50)$ $= e^{-\frac{50}{200}} = e^{-0.25}$	M1 E1	<p>Use of $1 - F(x)$</p> <p>Answer given: must be convincing. (= 0.7788(0))</p>	2
(C)	$P(X > 400) = e^{-\frac{400}{200}} = 0.1353(35)$ $P(X > 450) = e^{-\frac{450}{200}} = 0.1053(99)$ $P(X > 450 X > 400) = \frac{P(X > 450)}{P(X > 400)}$ $= \frac{e^{-\frac{450}{200}}}{e^{-\frac{400}{200}}} = e^{-\frac{50}{200}} = e^{-0.25} (= 0.7788)$	B1 B1 M1 A1	<p>Accept any form.</p> <p>Accept any form.</p> <p>Conditional probability. Not $P(X > 50) \times P(X > 400)$ unless <u>clearly</u> justified.</p> <p>Accept division of decimals, 3dp or better. Accept a.w.r.t. 0.778 or 0.779.</p>	4
				18

Q4	$C \sim N(10, 0.4^2), \quad D \sim N(35, 3.5^2)$ When a candidate's answers suggest that (s)he appears to have neglected to use the difference columns of the Normal distribution tables penalise the first occurrence only.			
		M1 A1 A1	For standardising. Award once, here or elsewhere. c.a.o.	3
(i)	$P(C < 9.5) = P\left(Z < \frac{9.5 - 10}{0.4} = -1.25\right)$ $= 1 - 0.8944 = 0.1056$	B1 B1 M1 A1	Mean. Accept +5 for $S - D$. Variance. Accept sd ($= 3.590\dots$). Formulation of requirement. Accept $S - D < 0$. This mark could be awarded in (iii) if not earned here. c.a.o.	4
(ii)	$D - S = D - (C_1 + C_2 + C_3 + C_4) \sim N(-5, \sigma^2 = 3.5^2 + (0.4^2 + 0.4^2 + 0.4^2 + 0.4^2) = 12.89)$ Want $P(D > S) = P(D - S > 0)$ $= 1 - \Phi\left(\frac{0 - (-5)}{3.59} = 1.39(27)\right)$ $= 1 - 0.9182 = 0.0818$			
(iii)	$New (D - S) = (D \times 1.3) - (C_1 + \dots + C_5) \sim N(-4.5, \sigma^2 = (3.5^2 \times 1.3^2) + (0.4^2 + \dots + 0.4^2) = 21.5025)$ Again want $P(D > S) = P(D - S > 0)$ $= 1 - \Phi\left(\frac{0 - (-4.5)}{4.637} = 0.9704\right)$ $= 1 - 0.8341 = 0.1659$	B1 M1 A1 A1	Mean. Accept +4.5 for $S - D$. Correct use of $\times 1.3^2$ for variance. c.a.o. Accept sd ($= 4.637\dots$) Or $S - D < 0$. M1 for formulation in (ii) available here. c.a.o.	4
(iv)	CI is given by $9.73 \pm 1.96 \times \frac{0.4}{\sqrt{12}}$ $= 9.73 \pm 0.2263 = (9.50(37), 9.95(63))$ Since 10 lies above this interval, it seems that the cheeses are underweight. In repeated sampling, 95% of all confidence intervals constructed in this way will contain the true mean.	M1 B1 M1 A1 E1 E1	1.96 seen. c.a.o. Must be expressed as an interval. Ft c's interval.	7
				18

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office: 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

