

Mathematics (MEI)

Advanced GCE

Unit **4757**: Further Applications of Advanced Mathematics

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

GENERAL MARKING INSTRUCTIONS

- 1 Please mark in RED.
- 2 Errors in the scripts should be clearly indicated (e.g. by ringing them). Please tick correct work. Some indication must be given on every page that the work has been assessed.
- 3 Marks for sections of questions should be written in the right-hand margin; detailed marks (e.g. M1A0M1A1) need only be shown when the breakdown of marks is not obvious.
- 4 The total mark for each question should be written in the right-hand margin at the end of the answer, and ringed.
- 5 Work crossed out by the candidate, and not replaced, should be marked wherever possible.
- 6 M: method marks
 - A: accuracy marks which are dependent on the relevant previous method mark(s) having been awarded (e.g. marks M0A1 cannot be awarded; where appropriate, the correct answer obtained with no method shown is taken to imply the method marks).
 - B: accuracy marks which are not dependent on method marks.
- 7 Marks are indivisible; if the scheme says M2, give either 2 or 0, except where indicated in the scheme.
- 8 Where there are two or more method marks for a section of work, the method marks can be awarded independently, except where indicated in the scheme.
- 9 cao: correct answer only (i.e. no follow through).

All A and B marks are cao unless stated otherwise.

ft: award mark for correct working following through from a previous error; use \checkmark to indicate correct ft work, and \times to indicate that a further error has been made. Ft marks are intended to ensure that a small arithmetic error is not unduly penalised; when the previous error is one of principle, particularly if the nature of the work is changed, or made considerably easier, the ft might not be given.
- 10 'Correctly obtained' means that all the working leading to that result must be correct.
- 11 Where the candidate is asked to show a given result, we expect the explanation to be reasonably clear.
- 12 If the candidate misreads a question, but in such a way that the nature and difficulty of the work is not changed, transfer all marks (including cao) to the new equivalent figures. Deduct the first A or B mark so earned (M marks are not lost for misread).

| | | | |
|-------|--|--|---|
| 1 (i) | Distance is $\frac{2(-2) - (-7) + 2(7) - 11}{\sqrt{2^2 + 1^2 + 2^2}} = 2$ | M1 A1 A1 3 | Formula, or other complete method Numerical expression for distance |
| (ii) | $\begin{pmatrix} 1 \\ 4 \\ 7 \end{pmatrix} \times \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 15 \\ 12 \\ -9 \end{pmatrix}$ Equation of AD is $\mathbf{r} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix}$ | M1 A1 A1 ft 3 | Vector product of normals, or finding a point on AD, e.g. $(0, -2.6, 4.2)$, $(3.25, 0, 2.25)$, $(7, 3, 0)$ Correct direction Accept any form |
| (iii) | $\overrightarrow{CA} \times \mathbf{d} = \begin{pmatrix} -5 \\ -6 \\ 2 \end{pmatrix} \times \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} = \begin{pmatrix} 10 \\ -5 \\ 10 \end{pmatrix}$ Distance is $\frac{ \overrightarrow{CA} \times \mathbf{d} }{ \mathbf{d} } = \frac{\sqrt{10^2 + 5^2 + 10^2}}{\sqrt{5^2 + 4^2 + 3^2}} = \frac{\sqrt{225}}{\sqrt{50}}$ $= \sqrt{4.5} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2} \approx 2.12$ | M1 A2 ft M1 M1 A1 6 | Appropriate vector product Give A1 if just one error Formula for distance Finding magnitude <i>Both dependent on first M1</i> |
| (iv) | $\mathbf{d} \times \overrightarrow{BC} = \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} \times \begin{pmatrix} 9 \\ 12 \\ -6 \end{pmatrix} = \begin{pmatrix} 12 \\ 3 \\ 24 \end{pmatrix} = 3 \begin{pmatrix} 4 \\ 1 \\ 8 \end{pmatrix}$ Distance is $\frac{\begin{pmatrix} -4 \\ -6 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 1 \\ 8 \end{pmatrix}}{\sqrt{4^2 + 1^2 + 8^2}} = \frac{10}{9}$ | M1 A1 ft M1 A1 ft M1 A1 A1 6 | Vector product of directions Appropriate scalar product Evaluation of scalar product For denominator |
| (v) | $V = \frac{1}{6} (\overrightarrow{AB} \times \overrightarrow{AC}) \cdot \overrightarrow{AD} = \frac{1}{6} \left[\begin{pmatrix} -4 \\ -6 \\ 4 \end{pmatrix} \times \begin{pmatrix} 5 \\ 6 \\ -2 \end{pmatrix} \right] \cdot \lambda \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix}$ $= \frac{1}{6} \lambda \begin{pmatrix} -12 \\ 12 \\ 6 \end{pmatrix} \cdot \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} = -5\lambda$ $V = \pm 20 \Rightarrow \lambda = \pm 4$ D is $(22, 15, -9)$ or $(-18, -17, 15)$ | M1 M1 A1 ft M1 A1A1 6 | Appropriate scalar triple product Evaluation of scalar triple product or $-2a + 2b + c + 3$ (simplified) for D(a, b, c) Obtain a value of λ , or one of a, b, c |



4757

Mark Scheme

June 2011

Alternative methods for Question 1

| | | | |
|---------|---|--|---|
| 1 (ii) | Eliminating x $3y + 4z = 9$ $x = 7 - \frac{5}{3}t, y = 3 - \frac{4}{3}t, z = t$ | M1 A1A1 3 | Eliminating one of x, y, z or $3x + 5z = 21$ or $4x - 5y = 13$ |
| 1 (iii) | $\left[\begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} - \begin{pmatrix} 7 \\ 5 \\ 1 \end{pmatrix} \right] \cdot \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} = 0$ $25\lambda - 25 + 16\lambda - 24 + 9\lambda - 6 = 0$ $\lambda = 1.1, \text{ F is } (7.5, 3.4, -0.3)$ $\text{CF} = \sqrt{(0.5)^2 + (-1.6)^2 + (-1.3)^2}$ $= \sqrt{4.5}$ | M1 A1 ft A1 ft M1 M1 A1 6 | Appropriate scalar product Obtaining a value for λ Finding magnitude |
| 1 (iv) | $\left[\begin{pmatrix} -2 \\ -7 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 9 \\ 12 \\ -6 \end{pmatrix} - \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} - \lambda \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} \right] \cdot \begin{pmatrix} 5 \\ 4 \\ -3 \end{pmatrix} = 0$ $\text{and } \begin{pmatrix} -5\lambda + 9\mu - 4 \\ -4\lambda + 12\mu - 6 \\ 3\lambda - 6\mu + 4 \end{pmatrix} \cdot \begin{pmatrix} 9 \\ 12 \\ -6 \end{pmatrix} = 0$ $\lambda = \frac{4}{81}, \mu = \frac{128}{243}$ $\text{Distance is } \sqrt{\left(\frac{40}{81}\right)^2 + \left(\frac{10}{81}\right)^2 + \left(\frac{80}{81}\right)^2}$ $= \frac{10}{9}$ | M1 A1 ft A1 ft M1 M1 A1 6 | Two appropriate scalar products Obtaining values for λ and μ Obtaining distance |

| | | | |
|-------|---|------|--|
| 2 (i) | When $y = -3$, $z = 3x^2 + 36x - 216$ | B1 | |
| |  | B1 | Correct shape (parabola) and position For $(-6, -324)$ |
| | When $x = -6$, $z = 8y^3 - 216y - 756$ | B1 | |
| |  | B1 | Correct shape and position For $(-3, -324)$ For $(3, -1188)$ |
| (ii) | (-6, -3, -324) is a SP on both sections; hence it is a SP on S Saddle point | B1 | |
| | | B1 | 2 |
| (iii) | $\frac{\partial z}{\partial x} = -12xy - 30x + 36$, $\frac{\partial z}{\partial y} = 24y^2 - 6x^2$ | B1B1 | |
| | At a SP, $\frac{\partial z}{\partial x} = \frac{\partial z}{\partial y} = 0$ | M1 | |
| | $24y^2 - 6x^2 = 0 \Rightarrow y = \pm \frac{1}{2}x$ | M1 | |
| | $y = \frac{1}{2}x \Rightarrow -6x^2 - 30x + 36 = 0$ | M1 | |
| | $\Rightarrow x = -6, 1$; SPs are $(-6, -3, -324)$ $(1, 0.5, 19)$ | A1 | |
| | $y = -\frac{1}{2}x \Rightarrow 6x^2 - 30x + 36 = 0$ | M1 | |
| | $\Rightarrow x = 2, 3$; SPs are $(2, -1, 28)$ $(3, -1.5, 27)$ | A1 | |
| | | A1 | 8 |
| (iv) | $\frac{\partial z}{\partial x} = 120$ and $\frac{\partial z}{\partial y} = 0$ | M1 | (Allow M1 for $\frac{\partial z}{\partial x} = -120$) |
| | $y = \frac{1}{2}x \Rightarrow -6x^2 - 30x - 84 = 0$; $D = 30^2 - 4 \times 6 \times 84$ | M1 | |
| | $D = -1116 < 0$; so this has no roots | A1 | |
| | $y = -\frac{1}{2}x \Rightarrow 6x^2 - 30x - 84 = 0 \Rightarrow x = 7, -2$ | M1 | Obtaining at least one value of x |
| | When $x = 7$, $y = -3.5$, $z = 203$; so $k = 637$ | M1 | Obtaining a value of k |
| | When $x = -2$, $y = 1$, $z = -148$; so $k = -92$ | A1 | |
| | | A1 | 7 |

| | | | |
|--------------|---|--|--|
| 3(a)(i) | $\frac{dy}{dx} = \frac{1}{2}e^{\frac{1}{2}x} - \frac{1}{2}e^{-\frac{1}{2}x}$ $\left(\frac{dy}{dx}\right)^2 = \frac{1}{4}e^x - \frac{1}{2} + \frac{1}{4}e^{-x}$ $1 + \left(\frac{dy}{dx}\right)^2 = \frac{1}{4}e^x + \frac{1}{2} + \frac{1}{4}e^{-x} = \left(\frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x}\right)^2$ | B1 M1 A1 (ag) 3 | Correct completion |
| (ii) | Length is $\int_0^{\ln a} \left(\frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x}\right) dx$ $= \left[e^{\frac{1}{2}x} - e^{-\frac{1}{2}x} \right]_0^{\ln a}$ $= \left(\sqrt{a} - \frac{1}{\sqrt{a}} \right) - (1-1) = \frac{a-1}{\sqrt{a}}$ | M1 A1 A1 (ag) 3 | For $\int \left(\frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x}\right) dx$ For $e^{\frac{1}{2}x} - e^{-\frac{1}{2}x}$ Correctly shown |
| (iii) | Curved surface area is $\int 2\pi y \, ds$ $= \int_0^{\ln a} 2\pi (e^{\frac{1}{2}x} + e^{-\frac{1}{2}x}) \left(\frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x}\right) dx$ $= \pi \int_0^{\ln a} (e^x + 2 + e^{-x}) dx$ $= \pi \left[e^x + 2x - e^{-x} \right]_0^{\ln a}$ $= \pi \left(a + 2 \ln a - \frac{1}{a} \right)$ | M1 A1 M1 A1 A1 5 | For $\int y \, ds$ Correct integral form <i>including limits</i> Obtaining integrable expression For $e^x + 2x - e^{-x}$ |
| (b)(i) | $\frac{dy}{dx} = \frac{\cos \theta}{-2 \sin \theta}$ Gradient of normal is $\frac{2 \sin \theta}{\cos \theta} (= 2 \tan \theta)$ Normal is $y - \sin \theta = 2 \tan \theta (x - 2 \cos \theta)$ $y = 2x \tan \theta - 3 \sin \theta$ | B1 M1 A1 (ag) 3 | Correctly shown |
| (ii) | Differentiating partially w.r.t. θ $0 = 2x \sec^2 \theta - 3 \cos \theta$ $x = \frac{3}{2} \cos^3 \theta$ $y = 3 \cos^3 \theta \tan \theta - 3 \sin \theta$ $= 3 \sin \theta (\cos^2 \theta - 1) = -3 \sin^3 \theta$ $(2x)^{\frac{2}{3}} + y^{\frac{2}{3}} = (3 \cos^3 \theta)^{\frac{2}{3}} + (-3 \sin^3 \theta)^{\frac{2}{3}}$ $= 3^{\frac{2}{3}} (\cos^2 \theta + \sin^2 \theta) = 3^{\frac{2}{3}}$ | M1 A1 M1 A1 M1 A1 (ag) 6 | Obtaining an expression for y Any correct form Using $1 - \cos^2 \theta = \sin^2 \theta$ Correctly shown |
| (iii) (A) | $(2, 0)$ has $\theta = 0$ Centre of curvature is $(\frac{3}{2}, 0)$ $\rho = \frac{1}{2}$ | M1 A1 | Using param eqn with $\theta = 0$ (or other method for ρ or cc) |
| (B) | $(0, 1)$ has $\theta = \frac{1}{2}\pi$ Centre of curvature is $(0, -3)$ $\rho = 4$ | M1 A1 | Using param eqn with $\theta = \frac{1}{2}\pi$ (or other method for ρ or cc) |

4757

Mark Scheme

June 2011

| | | | | |
|--|--|---|----------|---|
| 4 (i) | | 1 3 4 5 9 | | |
| | 1 | 1 3 4 5 9 | | |
| | 3 | 3 9 1 4 5 | B2 | Give B1 if not more than 4 errors |
| | 4 | 4 1 5 9 3 | | |
| | 5 | 5 4 9 3 1 | | |
| | 9 | 9 5 3 1 4 | | |
| Composition table shows closure Identity is 1 | | | B1 B1 | Dependent on B2 for table |
| Element 1 3 4 5 9 | | | B2 | Give B1 for 3 correct |
| Inverse 1 4 3 9 5 | | | 6 | |
| So every element has an inverse | | | | |
| (ii) | Since 5 is prime, a group of order 5 must be cyclic Two cyclic groups of the same order must be isomorphic | | | B1 B1 B1 3 |
| (iii) | H | 1 $e^{\frac{2\pi j}{5}}$ $e^{\frac{4\pi j}{5}}$ $e^{\frac{6\pi j}{5}}$ $e^{\frac{8\pi j}{5}}$ | | |
| | G | 1 3 9 5 4 | B1 | For $1 \leftrightarrow 1$ |
| | or | 1 4 5 9 3 | B2 | For non-identity elements |
| | or | 1 5 3 4 9 | | |
| | or | 1 9 4 3 5 | | |
| (iv) | Identity is (1, 1) Inverse of (9, 3) is (5, 4) | | | B1 B1 2 |
| (v) | $(x, y)^5 = (x^5, y^5)$ Since G has order 5, $x^5 = 1$ and $y^5 = 1$ Hence $(x, y)^5 = (1, 1)$ | | | M1 M1 A1 (ag) 3 |
| (vi) | Order of (x, y) is a factor of 5 (so must be 1 or 5) Only identity (1, 1) can have order 1 Hence all other elements have order 5 | | | M1 B1 A1 (ag) 3 |
| (vii) | $\{(1, 1), (9, 3), (4, 9), (3, 5), (5, 4)\}$ | | | B2 2 |
| | | | | Give B1 ft for 5 elements including (1, 1), (9, 3), (5, 4) |
| (viii) | An element of order 5 generates a subgroup, and so can be in only one subgroup of order 5 Number is $24 \div 4 = 6$ | | | M1 A1 2 |
| | | | | Or for $24 \div 4$ Or listing at least 2 other subgroups Give B1 for unsupported answer 6 |

Pre-multiplication by transition matrix

| | | | |
|--------|--|--------------------------------|--|
| 5 (i) | $P = \begin{pmatrix} 1 & 0.07 & 0 & 0 \\ 0 & 0.8 & 0.15 & 0 \\ 0 & 0.13 & 0.75 & 0 \\ 0 & 0 & 0.1 & 1 \end{pmatrix}$ | B2 | Allow tolerance of ± 0.0001 in probabilities throughout this question Give B1 for two columns correct |
| (ii) | If system enters an absorbing state, it remains in that state A and D are absorbing states | B1 B1 2 | |
| (iii) | $P^9 = \begin{pmatrix} 0 \\ 0.4 \\ 0.6 \\ 0 \end{pmatrix} = \begin{pmatrix} 0.2236 \\ 0.2505 \\ 0.1998 \\ 0.3261 \end{pmatrix}$ Prob(owned by B) = 0.2505 | M1M1 A1 3 | For P^9 (or P^{10}) and $\begin{pmatrix} 0 \\ 0.4 \\ 0.6 \\ 0 \end{pmatrix}$ |
| (iv) | $P^4 = \begin{pmatrix} 1 & \dots & \dots & \dots \\ \dots & 0.4818 & \dots & \dots \\ \dots & \dots & 0.3856 & \dots \\ \dots & \dots & \dots & 1 \end{pmatrix}$ $0.2236 + 0.2505 \times 0.4818 + 0.1998 \times 0.3856 + 0.3261 = 0.7474$ | M1 M1 A1 3 | Using diagonal elements from P^4 Using probabilities for 10 th day |
| (v) | $(1 \ 0 \ 0 \ 1) P^n = \begin{pmatrix} 0 \\ 0.4 \\ 0.6 \\ 0 \end{pmatrix}$ $= (0.8971) \text{ when } n = 26$ $= (0.9057) \text{ when } n = 27$ i.e. on the 28 th day | M1 M1 A1 ft A1 4 | Considering P^n for some $n > 20$ Evaluating Prob(A or D) for some values of n Identifying $n = 26$ or $n = 27$ (Implies M1M1) |
| (vi) | $P^n \rightarrow \begin{pmatrix} 1 & \mathbf{0.5738} & \mathbf{0.3443} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & \mathbf{0.4262} & \mathbf{0.6557} & 1 \end{pmatrix} = Q$ | B2 2 | Give B1 for two bold elements correct (to 3 dp) |
| (vii) | $Q = \begin{pmatrix} 0 \\ 0.4 \\ 0.6 \\ 0 \end{pmatrix} = \begin{pmatrix} 0.4361 \\ 0 \\ 0 \\ 0.5639 \end{pmatrix}$ Prob(eventually owned by A) = 0.4361 | M1M1 A1 3 | Using Q and $\begin{pmatrix} 0 \\ 0.4 \\ 0.6 \\ 0 \end{pmatrix}$ |
| (viii) | $Q = \begin{pmatrix} 0 \\ p \\ q \\ 0 \end{pmatrix} = \begin{pmatrix} 0.5 \\ 0 \\ 0 \\ 0.5 \end{pmatrix}$ (where $q = 1 - p$) $0.5738p + 0.3443q = 0.5$ $p = 0.6786, q = 0.3214$ | M1M1 A1 ft M1 A1 5 | For LHS and RHS Or $0.4262p + 0.6557q = 0.5$ Solving to obtain a value of p Allow 0.678 - 0.679, 0.321 - 0.322 |

Post-multiplication by transition matrix

| | | | |
|--------|--|---------------------------|--|
| 5 (i) | $\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0.07 & 0.8 & 0.13 & 0 \\ 0 & 0.15 & 0.75 & 0.1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ | B2 | Allow tolerance of ± 0.0001 in probabilities throughout this question Give B1 for two rows correct |
| (ii) | If system enters an absorbing state, it remains in that state A and D are absorbing states | B1 B1 | |
| (iii) | $(0 \ 0.4 \ 0.6 \ 0) \mathbf{P}^9$ $= (0.2236 \ 0.2505 \ 0.1998 \ 0.3261)$ Prob(owned by B) = 0.2505 | M1M1 A1 | (0 \ 0.4 \ 0.6 \ 0) and \mathbf{P}^9 (or \mathbf{P}^{10}) |
| (iv) | $\mathbf{P}^4 = \begin{pmatrix} 1 & \dots & \dots & \dots \\ \dots & 0.4818 & \dots & \dots \\ \dots & \dots & 0.3856 & \dots \\ \dots & \dots & \dots & 1 \end{pmatrix}$ $0.2236 + 0.2505 \times 0.4818 + 0.1998 \times 0.3856 + 0.3261$ $= 0.7474$ | M1 M1 A1 | Using diagonal elements from \mathbf{P}^4 Using probabilities for 10 th day |
| (v) | $(0 \ 0.4 \ 0.6 \ 0) \mathbf{P}^n \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}$ $= (0.8971) \text{ when } n = 26$ $= (0.9057) \text{ when } n = 27$ i.e. on the 28 th day | M1 M1 A1 ft A1 | Considering \mathbf{P}^n for some $n > 20$ Evaluating Prob(A or D) for some values of n Identifying $n = 26$ or $n = 27$ (Implies M1M1) |
| (vi) | $\mathbf{P}^n \rightarrow \begin{pmatrix} 1 & 0 & 0 & 0 \\ \mathbf{0.5738} & 0 & 0 & \mathbf{0.4262} \\ \mathbf{0.3443} & 0 & 0 & \mathbf{0.6557} \\ 0 & 0 & 0 & 1 \end{pmatrix} = \mathbf{Q}$ | B2 | Give B1 for two bold elements correct (to 3 dp) |
| (vii) | $(0 \ 0.4 \ 0.6 \ 0) \mathbf{Q} = (0.4361 \ 0 \ 0 \ 0.5639)$ Prob(eventually owned by A) = 0.4361 | M1M1 A1 | Using (0 \ 0.4 \ 0.6 \ 0) and \mathbf{Q} |
| (viii) | $(0 \ p \ q \ 0) \mathbf{Q} = (0.5 \ 0 \ 0 \ 0.5)$ (where $q = 1 - p$) $0.5738p + 0.3443q = 0.5$ $p = 0.6786, \ q = 0.3214$ | M1M1 A1 ft M1 A1 | For LHS and RHS Or $0.4262p + 0.6557q = 0.5$ Solving to obtain a value of p Allow 0.678 - 0.679, 0.321 - 0.322 |

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office: 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

