

Wednesday 25 January 2012 – Afternoon

A2 GCE MATHEMATICS (MEI)

4763 Mechanics 3

QUESTION PAPER

Candidates answer on the Printed Answer Book.

OCR supplied materials:

- Printed Answer Book 4763
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found in the centre of the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

- 1 The surface tension of a liquid enables a metal needle to be at rest on the surface of the liquid. The greatest mass m of a needle of length a which can be supported in this way by a liquid of surface tension S is given by

$$m = \frac{2Sa}{g}$$

where g is the acceleration due to gravity.

- (i) Show that the dimensions of surface tension are MT^{-2} . [3]

The surface tension of water is 0.073 when expressed in SI units (based on kilograms, metres and seconds).

- (ii) Find the surface tension of water when expressed in a system of units based on grams, centimetres and minutes. [3]

Liquid will rise up a capillary tube to a height h given by $h = \frac{2S}{\rho gr}$, where ρ is the density of the liquid and r is the radius of the capillary tube.

- (iii) Show that the equation $h = \frac{2S}{\rho gr}$ is dimensionally consistent. [3]

- (iv) Find the radius of a capillary tube in which water will rise to a height of 25 cm. (The density of water is 1000 in SI units.) [2]

When liquid is poured onto a horizontal surface, it forms puddles of depth d . You are given that $d = kS^\alpha \rho^\beta g^\gamma$ where k is a dimensionless constant.

- (v) Use dimensional analysis to find α , β and γ . [4]

Water forms puddles of depth 0.44 cm. Mercury has surface tension 0.487 and density 13 500 in SI units.

- (vi) Find the depth of puddles formed by mercury on a horizontal surface. [3]

- 2 A light inextensible string of length 5 m has one end attached to a fixed point A and the other end attached to a particle P of mass 0.72 kg.

At first, P is moving in a vertical circle with centre A and radius 5 m. When P is at the highest point of the circle it has speed 10 m s^{-1} .

- (i) Find the tension in the string when the speed of P is 15 m s^{-1} . [5]

The particle P now moves at constant speed in a horizontal circle with radius 1.4 m and centre at the point C which is 4.8 m vertically below A.

- (ii) Find the tension in the string. [3]

- (iii) Find the time taken for P to make one complete revolution. [4]

Another light inextensible string, also of length 5 m, now has one end attached to P and the other end attached to the fixed point B which is 9.6 m vertically below A. The particle P then moves with constant speed 7 m s^{-1} in the circle with centre C and radius 1.4 m, as shown in Fig. 2.




Fig. 2

- (iv) Find the tension in the string PA and the tension in the string PB. [6]

- 3 A bungee jumper of mass 75 kg is connected to a fixed point A by a light elastic rope with stiffness 300 N m^{-1} . The jumper starts at rest at A and falls vertically. The lowest point reached by the jumper is 40 m vertically below A. Air resistance may be neglected.

(i) Find the natural length of the rope. [4]

(ii) Show that, when the rope is stretched and its extension is x metres, $\ddot{x} + 4x = 9.8$. [3]

The substitution $y = x - c$, where c is a constant, transforms this equation to $\ddot{y} = -4y$.

(iii) Find c , and state the maximum value of y . [3]

(iv) Using standard simple harmonic motion formulae, or otherwise, find

(A) the maximum speed of the jumper,

(B) the maximum deceleration of the jumper. [3]

(v) Find the time taken for the jumper to fall from A to the lowest point. [5]

- 4 (a) The region T is bounded by the x -axis, the line $y = kx$ for $a \leq x \leq 3a$, the line $x = a$ and the line $x = 3a$, where k and a are positive constants. A uniform frustum of a cone is formed by rotating T about the x -axis. Find the x -coordinate of the centre of mass of this frustum. [6]

- (b) A uniform lamina occupies the region (shown in Fig. 4) bounded by the x -axis, the curve $y = 16(1 - x^{-\frac{1}{3}})$ for $1 \leq x \leq 8$ and the line $x = 8$.




Fig. 4

(i) Find the coordinates of the centre of mass of this lamina. [8]

A hole is made in the lamina by cutting out a circular disc of area 5 square units. This causes the centre of mass of the lamina to move to the point (5, 3).

(ii) Find the coordinates of the centre of the hole. [4]

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.