



Oxford Cambridge and RSA

Monday 27 June 2016 – Morning

A2 GCE MATHEMATICS (MEI)

4764/01 Mechanics 4

QUESTION PAPER



Candidates answer on the Printed Answer Book.

OCR supplied materials:

- Printed Answer Book 4764/01
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** If additional space is required, you should use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ ms}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **16** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

Section A (24 marks)

1 A car of mass m moves horizontally in a straight line. At time t the car is a distance x from a point O and is moving away from O with speed v . There is a force of magnitude kv^2 , where k is a constant, resisting the motion of the car. The car's engine has a constant power P . The terminal speed of the car is U .

(i) Show that

$$mv^2 \frac{dv}{dx} = P \left(1 - \frac{v^3}{U^3}\right). \quad [3]$$

(ii) Show that the distance moved while the car accelerates from a speed of $\frac{1}{4}U$ to a speed of $\frac{1}{2}U$ is

$$\frac{mU^3}{3P} \ln A,$$

stating the exact value of the constant A . [6]

Once the car attains a speed of $\frac{1}{2}U$, no further power is supplied by the car's engine.

(iii) Find, in terms of m , P and U , the time taken for the speed of the car to reduce from $\frac{1}{2}U$ to $\frac{1}{4}U$. [3]

2 A thin rigid rod PQ has length $2a$. Its mass per unit length, ρ , is given by $\rho = k\left(1 + \frac{x}{2a}\right)$ where x is the distance from P and k is a positive constant. The mass of the rod is M and the moment of inertia of the rod about an axis through P perpendicular to PQ is I .

(i) Show that $I = \frac{14}{9}Ma^2$. [5]

The rod is initially at rest with Q vertically below P. It is free to rotate in a vertical plane about a smooth fixed horizontal axis passing through P. The rod is struck a horizontal blow perpendicular to the fixed axis at the point where $x = \frac{3}{2}a$. The magnitude of the impulse of this blow is J .

(ii) Find, in terms of a , J and M , the initial angular speed of the rod. [2]

(iii) Find, in terms of a , g and M , the set of values of J for which the rod makes complete revolutions. [5]

Section B (48 marks)

3 Fig. 3 shows a uniform rigid rod AB of length $2a$ and mass $2m$. The rod is freely hinged at A so that it can rotate in a vertical plane. One end of a light inextensible string of length l is attached to B. The string passes over a small smooth fixed pulley at C, where C is vertically above A and $AC = 6a$. A particle of mass λm , where λ is a positive constant, is attached to the other end of the string and hangs freely, vertically below C. The rod makes an angle θ with the upward vertical, where $0 \leq \theta \leq \pi$. You may assume that the particle does not interfere with the rod AB or the section of the string BC.




Fig. 3

(i) Find the potential energy, V , of the system relative to a situation in which the rod AB is horizontal, and hence show that

$$\frac{dV}{d\theta} = 2mga \sin \theta \left(\frac{3\lambda}{\sqrt{10-6\cos\theta}} - 1 \right). \quad [6]$$

(ii) Show that $\theta = 0$ and $\theta = \pi$ are the only values of θ for which the system is in equilibrium whatever the value of λ . [2]

(iii) Show that, if there is a third value of θ for which the system is in equilibrium, then $\frac{2}{3} < \lambda < \frac{4}{3}$. [4]

(iv) Given that there are three positions of equilibrium, establish whether each of these positions is stable or unstable. [10]

It is given that, for small values of θ ,

$$\frac{dV}{d\theta} \approx 2mga \left[\left(\frac{3}{2} \lambda - 1 \right) \theta - \left(\frac{13}{16} \lambda - \frac{1}{6} \right) \theta^3 \right].$$

(v) Investigate the stability of the equilibrium position given by $\theta = 0$ in the case when $\lambda = \frac{2}{3}$. [2]

4 A raindrop falls from rest through a stationary cloud. The raindrop has mass m and speed v when it has fallen a distance x . You may assume that resistances to motion are negligible.

(i) Derive the equation of motion

$$mv \frac{dv}{dx} + v^2 \frac{dm}{dx} = mg.$$

[4]

Initially the mass of the raindrop is m_0 . Two different models for the mass of the raindrop are suggested.

In the first model $m = m_0 e^{k_1 x}$, where k_1 is a positive constant.

(ii) Show that

$$v^2 = \frac{g}{k_1} (1 - e^{-2k_1 x}),$$

and hence state, in terms of g and k_1 , the terminal velocity of the raindrop according to this first model.

[7]

In the second model $m = m_0 (1 + k_2 x)$, where k_2 is a positive constant.

(iii) By considering the expression obtained from differentiating $v^2 (1 + k_2 x)^2$ with respect to x , show that, for the second model, the equation of motion in part (i) may be written as

$$\frac{d}{dx} [v^2 (1 + k_2 x)^2] = 2g(1 + k_2 x)^2.$$

Hence find an expression for v^2 in terms of g , k_2 and x .

[9]

(iv) Suppose that the models give the same value for the speed of the raindrop at the instant when it has doubled its initial mass. Find the exact value of $\frac{k_1}{k_2}$, giving your answer in the form $\frac{a}{b}$ where a and b are integers.

[4]

END OF QUESTION PAPER



Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.