



Level 3 Certificate

Quantitative Problem Solving (MEI)

Unit **H867/02** Statistical Problem Solving

OCR Level 3 Certificate in Quantitative Problem Solving

Mark Schemes for June 2016

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2016

Annotations and abbreviations

Annotation	Meaning
 and 	
	Benefit of doubt
	Follow through
	Ignore subsequent working
 	Method mark awarded 0, 1
 	Accuracy mark awarded 0, 1
 	Independent mark awarded 0, 1
	Special case
	Omission sign
	Misread
	Highlighting

Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep *’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be ‘follow through’. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g Rules for replaced work
- If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

- i Anything in the mark scheme which is in square brackets [...] is not required for the mark to be earned, but if present it must be correct.

Question		Answer	Marks	Guidance
1	(i)	Her time is very predictable when she walks. (It is about 25 minutes.)	B1	Allow any two different sensible comments relating to the times
		On a good day driving is quicker, (taking about 8 minutes,) but on a bad day it can take her much longer.	B1	
			[2]	
1	(ii)(A)	The day of the week of each journey	B1	Allow any sensible answer (eg the weather conditions that day, traffic conditions, whether there was a detour) Do not allow: distance to work, date
		The starting time of each journey	B1	
	(ii)(B)	She should time herself on more journeys and record this additional information	B1	
		Alternative for (B): Description of how to make a decision based on data from (A)	B1	Must be consistent with (A)
			[3]	

Question	Answer	Marks	Guidance
2 (i)	Self-selected	B1	Accept Opportunity
		[1]	
2 (ii)	<p>It is bell shaped</p> <p>Mean = 7, sd = 3, so Mean \pm 1 sd is from 4 to 10</p> <p>68% of a Normal distribution lies within this range</p> <p>Total frequency from 4 to 10 is 4 + 5 + 5 = 14 or 70% of the data (\approx 68%)</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p>	<p>Accept 'It decreases from the centre symmetrically'.</p> <p>Use of given Normal distribution</p> <p>Allow 67 – 69%</p> <p>70% must be seen and compared to their previous answer</p>
	<p>Alternative</p> <p>Total frequency is 20 and 68% of 20 is 13.6</p> <p>Observed frequency from 4 to 10 is 4 + 5 + 5 = 14 (\approx 13.6)</p>	A1	13.6 must be seen
	<p>30 hours is $\frac{30-7}{3} = 7.7$ standard deviations from the mean.</p> <p>However virtually all of a Normal distribution is included within 3 standard deviations of the mean.</p>	<p>M1</p> <p>A1</p>	
	<p>Alternative</p> <p>Mean + 3sd</p> <p>= 16 and compare to 30</p>	<p>M1</p> <p>A1</p>	Allow mean + 2sd (= 13)
	<p>Special Case</p> <p>Argument based on the range of the sample</p>	SC B1	
		[6]	

Question	Answer	Marks	Guidance
2 (iii)	<p>Total number of hours for all 20 boys = $20 \times 7.5 = 150$ (9000 minutes)</p> <p>Total for those playing less than 10 hours is given by $18 \times 4\frac{2}{3} = 84$ hours (5040 minutes)</p> <p>So the other two spent $150 - 84 = 66$ hours (3960 minutes)</p> <p>Alternative</p> <p>If the other two spent 30 hours the mean would be 7.2 hours (432 minutes)</p> <p>So at least one of them one spent more than 30 hours.</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Valid strategy seen.</p> <p>Or average of art 33</p> <p>FT</p>
		[3]	
2 (iv)	<p>Only a small minority of children spend anything like 30 hours playing computer games.</p> <p>They spend more than 30 hours on school work</p> <p>The newspaper should give its sources of information.</p> <p>The newspaper gives no evidence that playing computer games is damaging children.</p> <p>Etc.</p>	<p>B1</p> <p>B1</p>	<p>Two sensible but different points</p>
		[2]	

Question	Answer	Marks	Guidance																								
3 (i)	<p>H_0 The proportions of the different species of bat are independent of the site</p> <p>H_1 The proportions are not independent of the site</p>	<p>B1</p> <p>[1]</p>	<p>Allow 'association' but not 'correlation'.</p> <p>Allow 'numbers of <i>different</i> bats'</p> <p>Allow missing H_0, H_1 if right way round.</p>																								
3 (ii)	<table border="1" data-bbox="295 555 1281 960"> <thead> <tr> <th data-bbox="295 555 629 619">Observed frequency, f_o</th> <th data-bbox="629 555 889 619">A</th> <th data-bbox="889 555 1140 619">B</th> <th data-bbox="1140 555 1281 619">Total</th> </tr> </thead> <tbody> <tr> <td data-bbox="295 619 629 687">Pipistrelles</td> <td data-bbox="629 619 889 687">135</td> <td data-bbox="889 619 1140 687">91</td> <td data-bbox="1140 619 1281 687">226</td> </tr> <tr> <td data-bbox="295 687 629 756">Noctule</td> <td data-bbox="629 687 889 756">10</td> <td data-bbox="889 687 1140 756">5</td> <td data-bbox="1140 687 1281 756">15</td> </tr> <tr> <td data-bbox="295 756 629 825">Serotine</td> <td data-bbox="629 756 889 825">10</td> <td data-bbox="889 756 1140 825">5</td> <td data-bbox="1140 756 1281 825">15</td> </tr> <tr> <td data-bbox="295 825 629 893">Horseshoe Bat</td> <td data-bbox="629 825 889 893">20</td> <td data-bbox="889 825 1140 893">2</td> <td data-bbox="1140 825 1281 893">22</td> </tr> <tr> <td data-bbox="295 893 629 960">Total</td> <td data-bbox="629 893 889 960">175</td> <td data-bbox="889 893 1140 960">103</td> <td data-bbox="1140 893 1281 960">278</td> </tr> </tbody> </table> <p data-bbox="730 1043 848 1075" style="text-align: center;">Table 3.1</p> <p data-bbox="295 1110 1059 1142">The numbers of other bats are too small to be included in the test</p>	Observed frequency, f_o	A	B	Total	Pipistrelles	135	91	226	Noctule	10	5	15	Serotine	10	5	15	Horseshoe Bat	20	2	22	Total	175	103	278	<p>B1</p> <p>B1</p>	<p>Accept They are not a coherent group oe</p>
Observed frequency, f_o	A	B	Total																								
Pipistrelles	135	91	226																								
Noctule	10	5	15																								
Serotine	10	5	15																								
Horseshoe Bat	20	2	22																								
Total	175	103	278																								
		[2]																									

Question	Answer	Marks	Guidance																								
3 (iii)	<table border="1" data-bbox="309 288 1258 715"> <thead> <tr> <th data-bbox="309 288 526 391">Expected frequency, f_e</th> <th data-bbox="526 288 831 391">A</th> <th data-bbox="831 288 1135 391">B</th> <th data-bbox="1135 288 1258 391">Total</th> </tr> </thead> <tbody> <tr> <td data-bbox="309 391 526 509">Pipistrelle</td> <td data-bbox="526 391 831 509">$(\frac{175 \times 226}{278} =)$142.266</td> <td data-bbox="831 391 1135 509">$(\frac{103 \times 226}{278} =)$83.734</td> <td data-bbox="1135 391 1258 509">226</td> </tr> <tr> <td data-bbox="309 509 526 560">Noctule</td> <td data-bbox="526 509 831 560">9.442</td> <td data-bbox="831 509 1135 560">5.558</td> <td data-bbox="1135 509 1258 560">15</td> </tr> <tr> <td data-bbox="309 560 526 611">Serotine</td> <td data-bbox="526 560 831 611">9.442</td> <td data-bbox="831 560 1135 611">5.558</td> <td data-bbox="1135 560 1258 611">15</td> </tr> <tr> <td data-bbox="309 611 526 662">G horseshoe</td> <td data-bbox="526 611 831 662">13.849</td> <td data-bbox="831 611 1135 662">8.151</td> <td data-bbox="1135 611 1258 662">22</td> </tr> <tr> <td data-bbox="309 662 526 715">Total</td> <td data-bbox="526 662 831 715">175</td> <td data-bbox="831 662 1135 715">103</td> <td data-bbox="1135 662 1258 715">278</td> </tr> </tbody> </table> <p data-bbox="725 730 842 762" style="text-align: center;">Table 3.2</p> $X^2 = \frac{(142.266 - 135)^2}{142.266} + \dots$ $= 0.3711 + 0.6305 + 0.0321 + 0.0560 + 0.0321 + 0.0560 + 2.7320 + 4.6417$ $= 8.55153$ <p data-bbox="297 1086 779 1118">Correct method for drawing a conclusion</p> $\nu = (4 - 1) \times (2 - 1) = 3 and Critical value at the 5% significance level is 7.815$ <p data-bbox="297 1219 1267 1283">(Since $8.55 > 7.815$) the alternative hypothesis is accepted. (The evidence suggests that the proportions of the different species of bats are dependent on the site.)</p> <p data-bbox="297 1315 1021 1347">The greatest influence comes from the Greater horseshoe bats</p>	Expected frequency, f_e	A	B	Total	Pipistrelle	$(\frac{175 \times 226}{278} =)$ 142.266	$(\frac{103 \times 226}{278} =)$ 83.734	226	Noctule	9.442	5.558	15	Serotine	9.442	5.558	15	G horseshoe	13.849	8.151	22	Total	175	103	278	<p data-bbox="1317 443 1368 475">M1</p> <p data-bbox="1317 639 1368 671">A1</p> <p data-bbox="1317 863 1368 895">M1</p> <p data-bbox="1317 995 1368 1027">A1</p> <p data-bbox="1317 1086 1368 1118">M1</p> <p data-bbox="1317 1145 1368 1177">B1</p> <p data-bbox="1317 1219 1368 1251">A1</p> <p data-bbox="1317 1315 1368 1347">B1</p>	<p data-bbox="1413 443 1765 475">Attempt at correct calculation</p> <p data-bbox="1413 576 1899 608">Accept rounding to 1 or 2 decimal places</p> <p data-bbox="1413 639 1989 671">Accept truncation after 3 or more decimal places</p> <p data-bbox="1413 863 2040 895">Attempt at correct calculation (at least one term seen)</p> <p data-bbox="1413 948 1733 979">No FT except for rounding</p> <p data-bbox="1413 1086 2007 1118">Attempt to compare their “8.55” to a critical value</p> <p data-bbox="1413 1145 1532 1177">Both seen</p> <p data-bbox="1413 1219 1693 1251">FT their “8.55” and CV</p>
Expected frequency, f_e	A	B	Total																								
Pipistrelle	$(\frac{175 \times 226}{278} =)$ 142.266	$(\frac{103 \times 226}{278} =)$ 83.734	226																								
Noctule	9.442	5.558	15																								
Serotine	9.442	5.558	15																								
G horseshoe	13.849	8.151	22																								
Total	175	103	278																								
		[8]																									


Question	Answer	Marks	Guidance
3 (iv)	The data show a major difference between the sites for the Greater horseshoe bats. This is confirmed by the test result.	B1 B1	A comment about the <i>data</i> in the table, can be about overall number of bats A comment about the test, must relate to proportions of different species
		[2]	

Question	Answer	Marks	Guidance
4 (i)	Japan: Population 127 103 388, Birth rate 8.07 Number of babies = $\frac{127\,103\,388 \times 8.07}{1000}$ = 1 025 724 or about 1 026 000	M1 A1	Can use rounded values Accept without rounding Must round to 1 026 000
		[2]	
	(ii) Number of deaths = $\frac{127\,103\,388 \times 9.38}{1000} = 1\,192\,230$ New population = $127\,103\,388 + 1\,025\,724 - 1\,192\,230 = 126\,936\,882$ Japan's population is decreasing	M1 A1 B1	Needs complete strategy to find change in population Accept without rounding Must round to 126 937 000
		[3]	

Question	Answer	Marks	Guidance
5 (i)	<p>239</p> <p>Either use the Σ command on cells C2 to C240, or enter =SUM(C2:C240) into C241</p> <p>Enter =C241/239 into C242</p> <p>30019474.02... rounding to 30 019 500</p> <p>Malaysia = 30 073 353, Uzbekistan = 28 929 716</p> <p>One is above and the other one below 30 019 500</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p>	<p>Can be seen in C241/239</p> <p>Or equivalent</p> <p>Or equivalent</p> <p>FT 29 894 393 from n = 240 30 145 606 from n = 238</p> <p>Both seen</p> <p>Explicit comparison</p>
		[6]	
5 (ii)	<p>Malaysia is ranked 43 so 43 countries have above the mean population</p> <p>Uzbekistan is ranked 44 so $239 - 43 = 196$ countries have below the mean population</p> <p>There are a lot of small countries and a few large ones, (notably China and India which have a large effect on the mean).</p>	<p>B1</p> <p>B1</p> <p>B1</p>	<p>FT: (239 – above answer)</p> <p>FT their n from (i)</p>
		[3]	

Question		Answer	Marks	Guidance
6	(i)	It is not representative, being from tail of the GDP per capita distribution It is a small sample.	B1	Or other valid statements
			[1]	
6	(ii)	They are numbers 7, 37, 67, ... 217 in the spreadsheet. 7 is a (random) starting point and then every 30. $\left(\frac{239}{8} \approx 30\right)$	B1 B1	Accept systematic sample
			[2]	

Question	Answer	Marks	Guidance																																																																					
6 (iii)	<p>H_0 There is no association between Population and GDP per capita</p> <p>H_1 There is negative association between Population and GDP per capita</p>	B1																																																																						
	<table border="1"> <thead> <tr> <th>Country</th> <th>Population</th> <th>GDP</th> <th>Pop rank</th> <th>GDP rank</th> <th>d</th> <th>d^2</th> </tr> </thead> <tbody> <tr> <td>Angola</td> <td>19 088 106</td> <td>6300</td> <td>3</td> <td>6</td> <td>-3</td> <td>9</td> </tr> <tr> <td>Mozambique</td> <td>24 692 144</td> <td>1200</td> <td>2</td> <td>8</td> <td>-6</td> <td>36</td> </tr> <tr> <td>Kiribati</td> <td>104 488</td> <td>6400</td> <td>8</td> <td>5</td> <td>3</td> <td>9</td> </tr> <tr> <td>Haiti</td> <td>9 996 731</td> <td>1300</td> <td>4</td> <td>7</td> <td>-3</td> <td>9</td> </tr> <tr> <td>Hong Kong</td> <td>7 112 688</td> <td>52 700</td> <td>5</td> <td>2</td> <td>3</td> <td>9</td> </tr> <tr> <td>Slovenia</td> <td>1 998 292</td> <td>27 400</td> <td>7</td> <td>3</td> <td>4</td> <td>16</td> </tr> <tr> <td>Norway</td> <td>5 147 792</td> <td>55 400</td> <td>6</td> <td>1</td> <td>5</td> <td>25</td> </tr> <tr> <td>Mexico</td> <td>120 286 655</td> <td>15 600</td> <td>1</td> <td>4</td> <td>-3</td> <td>9</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> <td>Σ</td> <td>0</td> <td>122</td> </tr> </tbody> </table> <p> $r_s = 1 - \frac{6 \sum d^2}{n(n^2 - 1)} = 1 - \frac{6 \times 122}{8 \times (64 - 1)} = -0.452\dots$ </p> <p>Critical value for $n = 8$ and 1-tail test at 5% significance level is 0.6429</p> <p>Since $0.452 < 0.6429$ the result is not significant. (The null hypothesis is accepted. The data do not support Jessica's conjecture.)</p>	Country	Population	GDP	Pop rank	GDP rank	d	d^2	Angola	19 088 106	6300	3	6	-3	9	Mozambique	24 692 144	1200	2	8	-6	36	Kiribati	104 488	6400	8	5	3	9	Haiti	9 996 731	1300	4	7	-3	9	Hong Kong	7 112 688	52 700	5	2	3	9	Slovenia	1 998 292	27 400	7	3	4	16	Norway	5 147 792	55 400	6	1	5	25	Mexico	120 286 655	15 600	1	4	-3	9					Σ	0	122	<p>B1 Rankings</p> <p>B1 Finding d (FT)</p> <p>B1 Finding and summing d^2 (FT)</p> <p>M1 Attempt formula for r_s</p> <p>A1 FT their '122'</p> <p>M1 Correct procedure: Finding a critical value and comparing to their r_s</p> <p>A1 Correct CV and conclusion FT from H_1 (CV=0.7381)</p>
Country	Population	GDP	Pop rank	GDP rank	d	d^2																																																																		
Angola	19 088 106	6300	3	6	-3	9																																																																		
Mozambique	24 692 144	1200	2	8	-6	36																																																																		
Kiribati	104 488	6400	8	5	3	9																																																																		
Haiti	9 996 731	1300	4	7	-3	9																																																																		
Hong Kong	7 112 688	52 700	5	2	3	9																																																																		
Slovenia	1 998 292	27 400	7	3	4	16																																																																		
Norway	5 147 792	55 400	6	1	5	25																																																																		
Mexico	120 286 655	15 600	1	4	-3	9																																																																		
				Σ	0	122																																																																		
		[8]																																																																						

Question	Answer	Marks	Guidance
<p>6 (iv)</p>		<p>B1 B1 B1 [3]</p>	<p>Qatar Bermuda USA</p>
<p>6 (v)</p>	<p>The evidence (from the hypothesis test) does not support the conjecture.</p> <p>USA is an anomaly.</p> <p>All the other countries with high GDP per capita are small. (So conjecture is supported.)</p> <p>If the few (about 20) countries with the highest GDP per capita are not considered then the scatter diagram does not suggest any correlation.</p>	<p>B1 B1 [2]</p>	<p>Allow other sensible comments about the evidence</p> <p>‘Conjecture supported’ needs evidence</p>

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2016

