



H

GCSE (9–1)

Mathematics

J560/04: Paper 4 (Higher tier)

General Certificate of Secondary Education

Mark Scheme for November 2019

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2019

Annotations used in the detailed Mark Scheme.

Annotation	Meaning
✓	Correct
✗	Incorrect
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working (after correct answer obtained), provided method has been completed
M0	Method mark awarded 0
M1	Method mark awarded 1
M2	Method mark awarded 2
A1	Accuracy mark awarded 1
B1	Independent mark awarded 1
B2	Independent mark awarded 2
MR	Misread
SC	Special case
^K	Omission sign

These should be used whenever appropriate during your marking.

The **M**, **A**, **B** etc annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks.

It is vital that you annotate these scripts to show how the marks have been awarded.

It is not mandatory to use annotations for any other marking, though you may wish to use them in some circumstances.

Subject-Specific Marking Instructions

1. **M** marks are for using a correct method and are not lost for purely numerical errors.
A marks are for an accurate answer and depend on preceding **M** (method) marks. Therefore **M0 A1** cannot be awarded.
B marks are independent of **M** (method) marks and are for a correct final answer, a partially correct answer, or a correct intermediate stage.
SC marks are for special cases that are worthy of some credit.
2. Unless the answer and marks columns of the mark scheme specify **M** and **A** marks etc, or the mark scheme is 'banded', then if the correct answer is clearly given and is not from wrong working **full marks** should be awarded.
Do not award the marks if the answer was obtained from an incorrect method, ie incorrect working is seen and the correct answer clearly follows from it.
3. Where follow through (**FT**) is indicated in the mark scheme, marks can be awarded where the candidate's work follows correctly from a previous answer whether or not it was correct.
Figures or expressions that are being followed through are sometimes encompassed by single quotation marks after the word *their* for clarity, eg $FT 180 \times (\text{their } '37' + 16)$, or $FT 300 - \sqrt{(\text{their } '5^2 + 7^2)}$. Answers to part questions which are being followed through are indicated by eg $FT 3 \times \text{their } (a)$.
For questions with **FT** available you must ensure that you refer back to the relevant previous answer. You may find it easier to mark these questions candidate by candidate rather than question by question.
4. Where dependent (**dep**) marks are indicated in the mark scheme, you must check that the candidate has met all the criteria specified for the mark to be awarded.
5. The following abbreviations are commonly found in GCSE Mathematics mark schemes.
 - **cao** means **correct answer only**.
 - **figs 237**, for example, means any answer with only these digits. You should ignore leading or trailing zeros and any decimal point
eg
237000, 2.37, 2.370, 0.00237 would be acceptable but 23070 or 2374 would not.
 - **isw** means **ignore subsequent working** (after correct answer obtained).
 - **nfw** means **not from wrong working**.
 - **oe** means **or equivalent**.

- **rot** means **rounded or truncated**.
 - **seen** means that you should award the mark if that number/expression is seen anywhere in the answer space, including the answer line,
even if it is not in the method leading to the final answer.
 - **soi** means **seen or implied**.
6. Make no deductions for wrong work after an acceptable answer unless the mark scheme says otherwise, indicated for example by the instruction 'mark final answer'.
7. As a general principle, if two or more methods are offered, mark only the method that leads to the answer on the answer line. If two (or more) answers are offered, mark the poorer (poorest).
8. When the data of a question is consistently misread in such a way as not to alter the nature or difficulty of the question, please follow the candidate's work and allow follow through for **A** and **B** marks. Deduct 1 mark from any **A** or **B** marks earned and record this by using the **MR** annotation. **M** marks are not deducted for misreads.
9. Unless the question asks for an answer to a specific degree of accuracy, always mark at the greatest number of significant figures even if this is rounded or truncated on the answer line. For example, an answer in the mark scheme is 15.75, which is seen in the working. The candidate then rounds or truncates this to 15.8, 15 or 16 on the answer line. Allow full marks for the 15.75.
10. If the correct answer is seen in the body and the answer given in the answer space is a clear transcription error allow full marks unless the mark scheme says 'mark final answer' or 'cao'. Place the annotation **✓** next to the correct answer.

If the answer space is blank but the correct answer is seen in the body allow full marks. Place the annotation **✓** next to the correct answer.

If the correct answer is seen in the working but a completely different answer is seen in the answer space, then accuracy marks for the answer are lost. Method marks would still be awarded. Use the **M0**, **M1**, **M2** annotations as appropriate and place the annotation **✗** next to the wrong answer.
11. Ranges of answers given in the mark scheme are always inclusive.
12. For methods not provided for in the mark scheme give as far as possible equivalent marks for equivalent work. If in doubt, consult your Team Leader.
13. Anything in the mark scheme which is in square brackets [...] is not required for the mark to be earned, but if present it must be correct.

Question		Answer	Mark	Part Marks and Guidance
1		Correct working leading to correct reason e.g. yes and 21.[3..] and 24 or 1280 and 1440 or [80 and] 90 or 2.6[...] or 2.7 [and 3] or 7.[1..] and 8	5	<p>M1 for unit calculation e.g. $48 \div 3$ M1 for scale factor e.g. 80×16 <i>possibly in one stage</i> M1 for first conversion e.g. $1280 \div 60$ M1 for second conversion e.g. 3×8 <i>possibly in one stage</i> A1 dep on M3 or M4 for 'Yes' and two correct comparative figures</p>
2		1.5×10^3	4	<p>B3 for 1500 or $1.50[0] \times 10^3$ or $1.48[9..] \times 10^3$ or 1.49×10^3 or B2 for 1489.1... or M1 for $\frac{5.8 \times 10^6}{\sqrt{4.1 \times 10^8 \times 3.7 \times 10^{-2}}}$ and M1 for writing <i>their</i> answer correct to 2s.f. and in standard form</p>
3		11 45 [am]	4	<p>SC3 for 11 45 pm OR B2 for LCM as 225 and M1 for <i>their</i> LCM correctly converted to hours and minutes OR M1 for $25 = 5 \times 5$ and $45 = 3 \times 3 \times 5$ and M1 for [0] 8 00 plus <i>their</i> LCM OR B1 for listing [0] 8:25, [0] 8:50, [0] 9:15... or 25, 50, 75... and B1 for listing [0] 8:45, [0] 9:30, 10:15... or 45, 90, 135..</p>


Question		Answer	Mark	Part Marks and Guidance
4	(a)	50×30 $\frac{50 \times 30}{1+2+3} [\times 1, 2 \text{ or } 3]$ $2 \times 250 = 500$	1 1 1	Accept any correct method e.g. M1 for $\frac{2}{1+2+3} = \frac{2}{6}$ M1 for $\frac{2}{6} \times 50 \text{ or } 50/3$ M1 for $\frac{50}{3} \times 30 = 500$ Alternative method M1 for $30 \div 6 = 5$ M1 for <i>their</i> $(30 \div 6) \times 2$ M1 for $50 \times 10 = 500$
	(b)	32	5	M1 for $\frac{250}{25} \text{ or } \frac{500}{20} \text{ or } \frac{750}{15}$ M1 for <i>their</i> $10 \times 5.5[0] \text{ or } 25 \times 2[.00] \text{ or } 50 \times 3.9[0]$ M1 for <i>their</i> $10 \times 5.5[0] + 25 \times 2[.00] + 50 \times 3.9[0]$ M1 for $\frac{396}{300} \text{ or } \frac{396 - \text{their } 300}{300}$ Alternative method M1 for $\frac{\frac{30}{6} \times 1}{25} \text{ or } \frac{\frac{30}{6} \times 2}{20} \text{ or } \frac{\frac{30}{6} \times 3}{15}$ or $\frac{396}{50}$ M1 for $5.5[0] \times \text{their } [0].2 \text{ or } 2[.00] \times \text{their } [0].5 \text{ or } 3.9[0] \times \text{their } 1$ M1 for <i>their</i> $(5.5[0] \times [0].2) + (2[.00] \times [0].5) + (3.9[0] \times 1)$ M1 for $\frac{7.92}{\text{their } 6} \text{ or } \frac{7.92 - \text{their } 6}{\text{their } 6}$

Question		Answer	Marks	Part Marks and Guidance
5		$4x^2 - 11xy - 3y^2$ final answer	3	<p>M2 for three correct terms from $4x^2 - 12xy + [1]xy - 3y^2$ oe or M1 for two correct terms in the expansion above</p> <p>M1 implied by two correct terms in answer</p>
6		$\frac{21}{50}$	3	<p>M1 for two ratios with a common number of sherberts e.g. 14 : 21 or 21 : 15 or 50 and</p> <p>M1 for $\frac{\text{their } 21}{\text{their } (14+21+15)}$ only after valid attempt to create a triple ratio eg 0 for $\frac{3+7}{2+3+7+5}$ etc allow any correct method</p> <p>Alternative : M1 for $7 \div 3$ implied by 2.3[3...] oe or better and M1 for $5 \div 2.3[3...]$ oe implied by 2.1428..... (exact = $\frac{15}{7}$) or $2 \times 2.3[3..]$ oe and e.g. 2 : 3 : $\text{their } \frac{15}{7}$ (= 14 : 21 : 15) or $\text{their } \frac{14}{3} : 7 : 5$</p>
7	(a)	2 points plotted correctly	1	tolerance $\pm \frac{1}{2}$ small square
	(b)	positive	1	Ignore embellishments
	(c) (i)	acceptable ruled line	1	see overlay, it must be at least from $x = 10$ to $x = 45$ and between (10,4) to (10,12) and (45,40) to (45,50) if more than 1 line, both must be in tolerance, ignore horizontal and vertical lines.
	(ii)	35 to 44	1	for answers out of tolerance FT <i>their</i> ruled line with positive gradient with tolerance $\pm \frac{1}{2}$ small square
	(d)	42 or 41.7 or 41.66... or 41.67	4	<p>B1 for 5 M1 for $\frac{\text{their } 5}{12}$ M1 for $(\text{their } \frac{5}{12}) \times 100$</p> <p>If 0 scored SC2 for answer of 30 from $\frac{3}{10}$ or 36[.36...] or 36.4 from $\frac{4}{11}$</p> <p><i>their</i> 5 must be less than 12 implied by [0].4166...</p>

Question		Answer	Mark	Part Marks and Guidance	
8		[adult =] 12[.00] [child =] 4.5[0]	5	B1 for $5a + 4c = 78$ B1 for $3a + 6c = 63$ M1 for multiplying/dividing both equations to get the coefficient of one variable equal (allow one error) M1 for correctly adding or subtracting both equations to eliminate one variable (allow one error)	one error in total
9	(a)	6	3	B1 for $3(k - 4) = k$ oe M1 for solving <i>their</i> equation to get k on one side and numbers on the other e.g. $3k - k = 12$	also $\frac{k}{3} + 4 = k$ condone use of other letters
	(b)	$\frac{y}{3} + 4$ oe	2	M1 for $\div 3$ and $+ 4$ e.g. $\frac{y+4}{3}$	
	(c)	16	2	B1 for output of first function as 36 or $3(3(n - 4) - 4) = 96$ or better	e.g. $9n - 48 = 96$
10	(a)	165 000	1		
	(b)	3	1		
	(c)	$165\ 000 \times 1.03^7$ 202 929.1878 truncated or rounded to at least 3sf	1 1		

Question		Answer	Mark	Part Marks and Guidance	
11	(a)	Accept any correct reason e.g. No as lengths are unknown or lengths may be double each one or triangles are similar	1		See exemplars in appendix
	(b)	First correct reason Second correct reason Third correct reason and SAS	1 1 1	Reasons are Angle [I]G[H] = angle [L]J[K] GH = JK GI = JL	Reasons can be given in any order Allow same angle, (36), but lines must be identified
12	(a)	24	1		condone 24 000
	(b)	14 26 36 50 60	2	B1 for complete table with three correct	
	(c)	28.5	5	B1FT for frequencies 14, 12, 10, 14, 10 with one error (allow FT table in (b)). B1 for 5, 15, 25, 40, 65 (allow one error) M1 for $\sum(\text{their midpoint} \times \text{their frequency})$ M1 for $\text{their } 1710 \div 60$	implied by $70 + 180 + 250 + 560 + 650$ or 1710 <i>their midpoint</i> must be within the group range
	(d)	Acceptable answer e.g. for the mean the figures used are approximate	1		See exemplars in appendix

Question		Answer	Mark	Part marks and Guidance
13	(a)	241.[1...]	3	M2 for $90 \div 72^3 [\times 100^3]$ or B1 for 72^3 or 373 248 or 100^3 or 1 000 000 implied by e.g. 90 000 000
	(b)	392	3	M2 for $8 \times 7 \times 7$ or M1 for $8 \times 7 \times 8$ or 8, 7, 7 clearly identified (e.g. summed) if 0 scored award SC2 for $8 \times 7 \times 6$ or SC1 for $8 \times 8 \times 8$, $6 \times 7 \times 6$ e.g. 448 as answer implied by 336 implied by 512 or 252
14		25	5	M1 for $m_1 = \frac{16}{-12}$ oe M1 for $m_2 = \frac{-1}{\text{their } m_1}$ or $\frac{12}{16}$ oe M1 for $y = (\text{their } \frac{12}{16})x + p$ M1 for substituting (-12,16) into <i>their</i> equation accept any correct method gradient of radius gradient of tangent
15	(a)	6.0[1...] or 6 nfww	3	M2 for $DF^2 = 10^2 + 12^2 - 2 \times 10 \times 12 \cos 30$ oe and allow one error or M1 for $\cos 30 = \frac{10^2 + 12^2 - DF^2}{2 \times 10 \times 12}$, condone one error nfww i.e. not $12 \times \sin 30$ M2 implied by $DF^2 = 36.15$ to 36.16 if they draw a perpendicular from F to P on DG award M1 for correct method to find both FP = 5 and PG = 8.6[6...] or 8.67 or 8.7 M1 for correct use of Pythagoras' on triangle FPD
	(b)	57.3[...] or 57 nfww	4	M2 for $[\sin B =] \frac{12.4 \times \sin 63}{12.8}$ or M1 for $\frac{\sin B}{12.4} = \frac{\sin 63}{12.8}$ oe and M1 for $180 - 63 - \text{their } 59.67$ implied by [0].863... or 59.67 to 59.7

Question		Answer	Mark	Part Marks and Guidance
16		$\frac{4(x-1)}{x^2-1} \text{ oe}$ $x + 9 + 4x - 4 \text{ or } 5x + 5$ <p>numerator = $5(x+1)$</p> <p>denominator = $(x+1)(x-1)$</p>	M1 also allow method with denominator $(x^2-1)(x+1)$ e.g. M1 for $\frac{(x+9)(x+1)}{(x^2-1)(x+1)}$ or $\frac{4(x^2-1)}{(x^2-1)(x+1)}$ M1 for $x^2 + 9x + x + 9 + 4x^2 - 4$ or better A1 If 0 scored SC1 for $x^2 - 1 = (x+1)(x-1)$	
17			2	B1 for either an acceptable curve and no/incorrect y intercept marked or any curve with 1 marked at the y intercept
18	(a)	330	2	M1 for $\frac{1}{2} \times 10 \times 6 \times 11$
	(b)	$\tan [...] = \frac{6}{10} \text{ oe}$ $30.9[...]$	1 1	accept any correct method, equivalents include $\tan^{-1}(0.6)$
	(c)	$\sqrt{257}$ cao	4	M3 for $\sqrt{11^2 + 10^2 + 6^2}$ or better or M2 for $11^2 + 10^2 + 6^2$ or M1 for $11^2 + 10^2$ or $10^2 + 6^2$ or $[\text{FD}] = \frac{10}{\cos 31}$ or $\frac{6}{\sin 31}$ implied by 16.0[3...] implied by 257 implied by $[\text{FD}] = \sqrt{136}$ or 11.6 to 11.7 or $[\text{AC}] = \sqrt{221}$ or 14.8 to 14.9

Question		Answer	Mark	Part Marks and Guidance
19		1.0 to 1.4 dep. on a tangent drawn	4	<p>B3 for an answer within the given range with a slightly inaccurate tangent, i.e. touches curve in range $t = 4.5$ to 5.5 but not at $t = 5$ or an answer of 1.3 using a symmetric chord about $t = 5$ e.g. from (4, 2) to (6, 4.6).</p> <p>OR</p> <p>B2 for an answer from using a symmetric chord about $t = 5$, e.g. from (4, 2) to (6, 4.6), with at most one error</p> <p>OR</p> <p>M1 for an attempt to draw a tangent touching curve at $t = 5$</p> <p>and</p> <p>M2 for <i>their</i> distance \div <i>their</i> time e.g. $(7.4 - 1.2) \div (8 - 3)$ or M1 for an attempt at <i>their</i> distance \div <i>their</i> time with one error or a correct attempt to find the gradient of <i>their</i> line which is not a tangent e.g. line from (0, 0) to (5, 3) which gives the average speed of $\frac{3}{5}$ or 0.6</p>

APPENDIXExemplar responses for Q11(a)

Response	Mark
No, lengths unknown	1
No, lengths may be different	1
No as triangles are similar	1

Exemplar responses for Q12(d)

Response	Mark
Because the median estimate is only affected by mistakes in measuring	1
Because the mean relies on assumption that all values lie at midpoint	1
Median is accurately read from graph, mean is estimated	1
Number of people earning each amount is not evenly spread out	1(BOD)
Mean has no exact values ...	1
Accurate line graph instead of an assumption	1(BOD)

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2019

