Write your name here Surname	Other	names
Pearson Edexcel International GCSE	Centre Number	Candidate Number
Chemistry Unit: 4CH0 Paper: 2CR	y	
Wednesday 14 June 2017 Time: 1 hour	– Morning	Paper Reference 4CH0/2CR
You must have: Ruler, calculator		Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







THE PERIODIC TABLE

0	
7	
9	
2	
4	
က	
Group	
0	
	

Hydrogen

Period

ß

က

4 **H**eije 2

			Π			Г															
ຂ	Š	Neon 10	40	Ā	Argon 18	84	궃	Krypton	98	131	×	Xenon	¥	222	뜐	Radon	98				
40	ш	Fluorine 9	35.5	రె	Chlorine 17	80	ă	Bromine	35	127	_	lodine	Sa	210	¥	Astatine	85				
16	0	Oxygen 8	32	S	Sulfur 16	7.9	တိ	Selenium	35	128	<u>e</u>	Tellurium	52	210	S.	Polonium	84				
4	z	Nitrogen 7	31	۵	Phosphorus 15	75	As	Arsenic	83	122	S	Antimony	51	508	ã	Bismuth	83				
12	O	Carbon 6	28	Ö	Silicon 14	7.3	Ge	Germanium	35	119	Š	Т'n	50	207	ص م	Lead	82				
F	ω	Boron 5	27	₹	Aluminium 13	02	Ga	Gallium	3	115		Indium	49	204	F	Thallium	18				
			1			65	Zu	Zinc	8	112	8	Cadmium	48	201	몬	Mercury	80				
						63.5	ਹ	Copper	53	108	Aa	Silver	47	197	Αn	Gold	62				
						53	Z	Nickel	28	106	Pd	Palladium	94	195	ă	Platinum	78				
						59	රි	Cobalt	27	103	듄	Rhodium	45	192	_	Iridium	77				
						26	ē.	rou	92	101	2	Ruthenium	4	190	ő	Osmium	9/				
						55	Ž	Mandanese	25	8	ပ	Technetium	43	186	æ	Rhenium	75	ł			
										96											
						51	>	Vanadium	83	83	2	Niobium	14	181	Ę	Tantalum	73				
						48	j=	Titanium	22	91	Ž	Zirconium	40	179	Ĭ	Hafnium	75				
						45	တိ	Scandium	21	68	>	Yttrinm	33	139	2	Lanthanum	57	227	Ac	Actinium	
			Τ		<u> </u>		œ	E	8	88	Ų.	ntium	98	137	Ba	m m	26	526	g	Ē	
6	Be	Benyllium	24	Ž	Magne:	¥	O	S	.,	_	•	Stro			_	å		"	Œ	Rad	

Key

Relative atomic mass
Symbol
Name

4

2

မှ


^

BLANK PAGE



Answer ALL questions.

1 The diagram represents an atom of an element.



- (a) (i) What is the particle labelled X?
 - A an electron
 - **B** an ion

 - **D** a neutron
 - (ii) What is the mass number of this atom?

(1)

(1)


- A 4
- \bowtie B 5
- **C** 9
- (iii) Name the element that contains these atoms.

(1)

(Total for Question 1 = 5 n	narks)
different	
same	(2)
State, in terms of subatomic particles, one way in which these isotopes are the same and one way in which they are different.	(2)
(b) Hydrogen has three isotopes.	

	(Total for Question 2 = 4	marks)
	(c) Write a word equation for the reaction.	(1)
	(b) The reaction is exothermic. State what happens to the temperature of the acid during the reaction.	(1)
2		
l		(2)
	Hydrogen gas is produced. (a) State two observations that are seen during the reaction.	
2	A small piece of magnesium ribbon is added to dilute sulfuric acid in a test tube.	

3 This apparatus can be used to prepare carbon dioxide from reagents A and B.



(a) Calcium chloride and water are also products of the reaction between A and B. Identify reagent A and reagent B.

(2)

A	
---	--

B......

(b) In the diagram, the carbon dioxide is collected over water.

State another way of collecting the carbon dioxide.

(1)

(c) At the end of the experiment, the solution in the trough is weakly acidic.

- (i) State the colour of the solution when some Universal Indicator is added.
- (ii) Give the name and the formula of the acid that forms when carbon dioxide dissolves in water.

formula

(2)

(1)

name

(Total for Question 3 = 6 marks)




4 Cr		rude oil is a complex mixture containing many compounds and is a source of many chemicals.									
	(a) Most of the compounds in crude oil contain two elements.										
		Name these two elements.									
			(1)								
		and									
	(b)	Crude oil is separated into fractions in order to produce useful chemicals.									
		(i) State what is meant by the term fraction .									
			(1)								
		(ii) Describe the industrial process used to obtain fractions from crude oil.									
			(3)								
•••••											
•••••											

	bitumen	diesel	fuel oil	gasoline	
(i) Which of	these fractions co	ontains comp	ounds with th	e highest boiling p	points?
(ii) Which of	these fractions is	the most vol	atile?		(1)
d) Fuel oil can b	pe burned to heat	homes.			
If combustio	n is incomplete, a	dangerous g	jas is produced	d.	
(i) Name thi	s gas.				(1)
(ii) State wh	y this gas is dang	erous.			(1)
			(Tota	al for Question 4 :	= 9 marks)



5 Ammonia is a toxic gas that is very soluble in water.

A teacher uses this apparatus to investigate the solubility of ammonia in water at different temperatures.



This is the teacher's method.

- pour 100 cm³ of water into the beaker and measure the temperature of the water
- place the beaker on the balance and record the mass of the beaker and water
- bubble ammonia into the water until the mass is constant
- record the constant mass

The teacher repeats the experiment with the water at different temperatures.

The table shows the teacher's results.

Temperature of water in °C	Mass at start in g	Mass at end in g
15	145.0	204.5
20	145.0	198.1
25	145.0	191.6
30	145.0	185.1

(a) (i) Calculate the mass of ammonia dissolved in 100 cm ³ of water at 25 °C.	(1)
mass of ammonia =(ii) State the relationship between temperature and solubility of ammonia.	(1)
(b) Explain one safety precaution that the teacher should take when doing this experi	iment.
(c) When the teacher does the experiment at a higher temperature, the reading on the balance gradually increases but then slowly decreases. Suggest why the reading on the balance slowly decreases.	(1)
(d) Ammonia is an alkaline gas. Suggest a different method that the teacher could use to compare the mass of ammonia dissolved in the water at different temperatures.	(1)
(Total for Question 5 = 6 ma	nrks)




C Falson	al as a ha maada l	t	favorat va atla a da			
6 Ethan	ol can be made l					
	method 1	ferment	ation of glucose			
	method 2	reaction	of ethene with steam			
(a) Na	ame the catalyst	used in ea	ach method.	(
method 1						
method 2)					
/I- \ -		الدر ومنام	and for different com-			
			anol for different purposes.			
Th	ne table gives sor	ne inform	ation about each company.			
			Company A	Company B		
	Location of con	npany	large agricultural area	near an oil refinery		
	Use of ethanol		to obtain a dilute solution to convert into vinegar	as a solvent for perfumes		
E	xplain which me	thod of p	roduction each company is r			
_				(
Company	[,] A					
Company	В					



- (c) Most of the ethene used to make polymers is produced by the cracking of crude oil fractions.
 - (i) One of the polymers made from ethene is poly(ethene).

Complete the equation to show the formation of poly(ethene) from ethene.

(2)



(ii) The kerosene fraction obtained from crude oil contains a hydrocarbon with the formula $C_{16}H_{34}$

Complete the equation to show the formation of ethene and one molecule of another hydrocarbon from the cracking of $C_{16}H_{34}$

(1)

$$C_{16}H_{34} \rightarrow 2C_{2}H_{4} + \dots$$

(iii) Suggest why it may be necessary, in future, to make ethene from ethanol.

(1)


(Total for Question 6 = 10 marks)



BLANK PAGE



7 A student uses this apparatus to investigate the change in temperature when dilute hydrochloric acid is added to aqueous sodium hydroxide.



This is the student's method.

- pour some aqueous sodium hydroxide into the polystyrene cup
- record the temperature of the sodium hydroxide
- add some dilute hydrochloric acid and stir the mixture
- record the highest temperature of the mixture


The student repeats the experiment using different volumes of the two solutions.

(a) Explain why the student uses a polystyrene cup to contain the solution, rather than a beaker.

.....

(2)

(b) The diagram shows the thermometer readings for one experiment before and after adding the acid.



Record the temperatures before and after adding the acid.

(2)

before°C

after°C




(c) The table shows the results of a series of experiments.

The initial temperatures of the aqueous sodium hydroxide and the dilute hydrochloric acid are the same.

Experiment	Volume of aqueous sodium hydroxide in cm ³	Volume of dilute hydrochloric acid in cm ³	Highest temperature of mixture in °C
1	10	90	22.2
2	20	80	24.2
3	30	70	26.0
4	70	30	24.0
5	80	20	23.0
6	90	10	22.0

(i) Plot the results from the table on the grid.



Draw a straight line of best fit for experiments 1, 2 and 3.

Draw a second straight line of best fit for experiments 4, 5 and 6.

Extend both lines so that they cross.

(4)



(ii) The point where the two lines cross indicates when equal amounts, in moles, of sodium hydroxide and hydrochloric acid react.		
Use your graph to find the volumes that contain equal amounts of sodium hydroxide and hydrochloric acid.		
	(2)	
volume of sodium hydroxide		cm³
volume of hydrochloric acid		cm³
(iii) The equation for the reaction between sodium hydroxide and hydrochloric act	id is	
NaOH + HCl \rightarrow NaCl + H ₂ O		
Explain which solution, the sodium hydroxide or the hydrochloric acid, has the greater concentration.	į	
greater concentration.	(2)	

(Total for Question 7 = 12 marks)

8 A student does a titration to find the concentration of a solution of nitric acid.

This is the student's method.

- pipette 25.0 cm³ of the nitric acid into a conical flask
- add a few drops of indicator
- add aqueous potassium hydroxide from a burette until the indicator just changes colour
- determine the volume of alkali added from the burette

The concentration of the potassium hydroxide solution is 0.0200 mol/dm³.

The volume of potassium hydroxide required to neutralise the acid is 23.50 cm³.

The equation for the reaction between nitric acid and potassium hydroxide is

$$\mathsf{HNO_3} + \mathsf{KOH} \to \mathsf{KNO_3} + \mathsf{H_2O}$$

(a) (i) Calculate the amount, in moles, of KOH used in this titration.

(2)

(ii) Calculate the concentration, in mol/dm³, of the nitric acid.

(2)

concentration of nitric acid = mol/dm³

 b) The student makes a solution of potassium nitrate by neutralising aqu potassium hydroxide with dilute nitric acid. 	eous	
Describe how he could use crystallisation to obtain a pure, dry sample potassium nitrate crystals from the solution of potassium nitrate.	e of (4)	
	(=)	
(Total for Questi	on 8 = 8 marks)	
TOTAL FOR PAPER = 60 MARKS		



BLANK PAGE

Every effort has been made to contact copyright holders to obtain their permission for the use of copyright material. Pearson Education Ltd. will, if notified, be happy to rectify any errors or omissions and include any such rectifications in future editions.