Write your name here Surname	Other r	names
Pearson Edexcel International GCSE	Centre Number	Candidate Number
Chemistry Unit: 4CH0 Paper: 2C	y	
Wednesday 13 June 2018 Time: 1 hour	– Morning	Paper Reference 4CH0/2C
You must have: Calculator, ruler		Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







THE PERIODIC TABLE

/ ဖ S က Group a

0

4 Heligia 4

	32 Sulfur 16	79 Selenium 34	128 Te Tellurium 52	210 Polonium 84
Nitrogen 7	31 P Phosphorus 15	75 As Arsenic 33	122 Sb Antimony 51	209 Bismuth 83
12 Carbon 6	Silicon 14	73 Ge Germanium 32		
11 Boron 5	27 Al Atuminium 13	70 Ga 31	115 Indium 49	
		65 Zinc 30	112 Cd Cadmium 48	201 Hg Mercury 80
		Copper 29	Ag Silver 47	Au Gold 79

Hydrogen		
	I	Hydrogen 1

Pd Palladium 46 195 P†	28 Z S
Rhodium 45 192	59 Cobalt
Ruthenium 44 44 C	% Te
Tc Technetium A 43 186 Re	55 Mn Manganese 25
Molybdenum 42 184	_
No liobium	51 V Inadium 23

Niobium 41 181 Ta Tantalum 73 48
Tilanium
22
91
Zreonium
40
179
HAfrium
72 SC Scandium 21 89 Y Yttrium 39 139 139 139 139 139 ACActinium 89 Beeylilium 4 4 Mg Mg Magnessum 24 40 Caa Caalcium 20 Ca Strontium 20 St Strontium 38 Ba Banum 56 56 226 Raa Radium 88 88 88 88 88 88

ဖ

Key

Relative atomic mass Symbol Name Atomic number

Period


N

7
Lithium
3
Sodium
11
11
39
Sodium
11
11
39
Rb
Rubidium
37
133
CS
Caestum
55
55
78
Francium
87

S

Answer ALL questions.

1 The diagram shows a Bunsen burner.



(a) The Bunsen burner uses methane as a fuel.

Methane has the formula CH₄

Give the names of the two elements in methane.

(2)

(b) When methane burns it reacts with a gas in the air.

Give the name of this gas.

(1)

(c) (i) Name the two substances that form when methane burns in plenty of air.

(2)

Z......


(ii) Name the poisonous gas that forms when methane burns in a shortage of air.

(1)

(Total for Question 1 = 6 marks)



2 A student uses this apparatus to investigate the burning of four different liquid fuels, W, X, Y and Z.



The table shows the student's results.

Fuel	Initial temperature in °C	Final temperature in °C	Increase in temperature in °C
W	19.0	31.3	12.3
Х	18.4	28.7	
Υ	19.5	35.4	
Z	18.7	29.8	

(a) Complete the table by giving the increase in temperature for fuels X, Y and Z.

(1)

(b) The student uses the same mass of water and burns each fuel for the same period of time. Explain which fuel releases the most heat energy.

1	ľ	ø	ь.	١
-(l	d	_	J

(c) What is the name given to reactions that release heat energy?

(1)


- A decomposition
- **B** endothermic
- **D** reduction

(Total for Question 2 = 4 marks)

A student makes an alkali solution by dissolving a small volume of cleaning liquid in deionised water.

He then titrates a sample of this solution with an acid until neutralisation is complete.

(a) The diagram shows the burette readings for his titration.



Use the readings to complete the table, giving all values to the nearest 0.05 cm³.

(2)

Burette reading after adding the acid	20.80
Burette reading before adding the acid	
Volume of acid added in cm ³	

(b) Another student does a titration using a solution of a different cleaning liquid.

The table shows her results.

Burette reading after adding the acid	29.65	28.70	29.25	29.10	28.55
Burette reading before adding the acid	3.40	3.60	3.50	3.80	3.35
Volume of acid added in cm ³	26.25	25.10	25.75	25.30	25.20
Concordant results (✓)					

Concordant results are those that differ by 0.20 cm³ or less.

(i) Place ticks in the table to show which results are concordant.

(1)

(ii) Use the concordant results to calculate the average (mean) volume of acid added.

(1

(Total for Question 3 = 4 marks)

- **4** Bromine, chlorine, fluorine and iodine are elements in Group 7 of the Periodic Table.
 - (a) Which element is the most reactive?

(1)

- A bromine
- **B** chlorine
- C fluorine
- **D** iodine
- (b) Which element is a solid at room temperature?

(1)

- **A** bromine
- B chlorine
- C fluorine
- **D** iodine
- (c) Which element has the darkest colour at room temperature?

- A bromine
- B chlorine
- C fluorine
- **D** iodine

(d) Bromine reacts with hydrogen to form hydrogen bromide.

The equation for the reaction is

$$H_2(g) + Br_2(g) \rightarrow 2HBr(g)$$

The table shows some average bond energies.

Bond	Н—Н	Br—Br	H—Br
Average bond energy in kJ/mol	436	193	366

Use the values in the table to calculate the enthalpy change for the reaction between hydrogen and bromine.

(3)

(Total for Question 4 = 6 marks)



- **5** Ethanol can be manufactured by fermentation or by the direct hydration of ethene.
 - (a) In Brazil, the main source of sugar for fermentation is sugar cane.
 - sugar cane is added to water
 - sugar cane contains sucrose $(C_{12}H_{22}O_{11})$ that dissolves in the water
 - during the fermentation process the sucrose is broken down into glucose (C₆H₁₂O₆)
 - this glucose is then converted into ethanol (C₂H₅OH) and carbon dioxide
 - (i) Name the substance that is added to the sucrose solution to allow fermentation to take place.

(1)

(ii) Complete the equation for the conversion of sucrose into glucose.

(1)

$$C_{12}H_{22}O_{11} + H_2O \rightarrow \dots$$

(iii) Write a chemical equation for the conversion of glucose into ethanol and carbon dioxide.

(1)

(iv) Fermentation produces a solution that is a mixture of ethanol and water.

Which of these is the most effective method of obtaining ethanol from this mixture?

- A crystallisation
- **B** filtration
- C fractional distillation
- **D** simple distillation



(b) In the direct hydration method, ethene reacts with steam.

The equation for the reaction is

$$C_2H_4 + H_2O \rightarrow C_2H_5OH$$

(i) Name the catalyst used in this reaction.

(1)


(ii) State the temperature and pressure used in this reaction.

(2)

temperature

pressure.

(c) This apparatus is used to convert ethanol into ethene.



(i) Name the type of reaction taking place.

(1)

(ii) State the function of the aluminium oxide in this reaction.



	(ii)	State the colour change that is observed when bromine water is shaken with ethene in a test tube.	
/1\	 		(1)



- **6** Phosgene (COCl₂) is used in industry to make polymers.
 - (a) Phosgene is formed when carbon monoxide reacts with chlorine.

$$CO(g) + CI_{2}(g) \rightleftharpoons COCI_{2}(g)$$

$$\Delta H = -108 \,\mathrm{kJ/mol}$$

(i) The reaction mixture is kept at temperatures between 50 and 150 °C. If a temperature above 200 °C is used, only a small amount of phosgene is formed.

Suggest why only a small amount of phosgene is formed at temperatures above 200 °C.



(ii) Predict how the yield of phosgene will change if the reaction is carried out at a higher pressure.

Give a reason for your answer.
[assume the reaction reaches a position of equilibrium]



(b) Phosgene reacts with water to form hydrochloric acid and carbon dioxide.

Write a chemical equation for this reaction.



(c) The diagram shows the displayed formula of phosgene.

Draw a dot and cross diagram to show the arrangement of all the outer electrons in a molecule of phosgene.

(3)

(Total for Question 6 = 8 marks)



7 Magnesium carbonate decomposes when heated to form magnesium oxide and carbon dioxide. The equation for the reaction is

$$MgCO_{3}(s) \rightarrow MgO(s) + CO_{2}(g)$$

A student uses this method to investigate the reaction.

- Step 1 weigh a clean, dry crucible and record the mass
- Step 2 add some magnesium carbonate
- Step 3 reweigh the crucible and contents and record the new mass
- Step 4 heat the crucible and contents for five minutes
- Step 5 allow the crucible and contents to cool and then reweigh
- Step 6 repeat steps 4 and 5 until the mass of the crucible and contents does not change

The student does the experiment four times.

The table shows her results.

		Mass	s in g	
	Experiment 1	Experiment 2	Experiment 3	Experiment 4
mass of empty crucible	19.20	21.31	19.83	20.45
mass of crucible and magnesium carbonate before heating	23.40	24.94	24.65	26.92
mass of crucible and contents after heating for 5 minutes	22.85	23,21	22.13	24.02
mass of crucible and contents after heating for a total of 10 minutes	21.94	23.04	22.13	23.53
mass of crucible and contents after heating for a total of 15 minutes	21.60	23.04	22.13	23.53

(a) State why the mass of the crucible and contents decreases during heating.



(b) (i)	State the reason for Step 6.	(1)
(ii)	Explain in which experiment the student should have heated for a fourt of five minutes.	h period (2)
	(Total for Question 7	= 4 marks)



0	Acid indigestion i	ic caused by	having too	much hydre	ochloric ocid i	tha stamach
0	Acid indigestion	is caused by	y naving too	much nyur	ocinoric acid ii	i tile stornacii

A suspension of magnesium hydroxide, Mg(OH)₂, in water, can be used to cure acid indigestion.

The equation for the reaction between magnesium hydroxide and hydrochloric acid is

$$Mg(OH)_2(s) + 2HCI(aq) \rightarrow MgCI_2(aq) + 2H_2O(I)$$

A student investigates how much magnesium hydroxide is needed to neutralise 100 cm³ of hydrochloric acid with a concentration of 0.0968 mol/dm³.

He uses 0.29 g of magnesium hydroxide to neutralise the hydrochloric acid.

(a) Calculate the amount, in moles, of HCl in the hydrochloric acid.

(2)

amount of HCI mol

(b) Calculate the amount, in moles, of $Mg(OH)_2$ used by the student. $[M_r \text{ of } Mg(OH)_2 = 58]$

(2)

amount of Mg(OH)₂ mol


(c) Explain whether the student used the right amount of magnesium hydroxide to neutralise the hydrochloric acid.

(2)

(Total for Question 8 = 6 marks)



9 The flow chart shows the steps in the manufacture of sulfuric acid.



(a) The equation for the reaction in step 2 is

$$SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$$
 $\Delta H = -96 \text{ kJ/mol}$

(i) State what the symbols \rightleftharpoons and ΔH represent.

(2)

(ii) Name the catalyst used in step 2.

(1)

(iii) State the temperature and pressure used in the reaction in step 2.

(2)

temperature

pressure..



(b)	Sulfur trioxide reacts with water to form sulfuric acid.
	This reaction is very exothermic.

$$SO_3(g) + H_2O(I) \rightarrow H_2SO_4(aq)$$
 $\Delta H = -228 \text{ kJ/mol}$

- (i) State why the sulfur trioxide is not dissolved in water to form sulfuric acid in step 3. (1)
- (ii) Write chemical equations for the reactions that take place in step 3 and step 4.

(2)

(2)

step 3

step 4.....

(c) Give two industrial uses for sulfuric acid.

I

2

(Total for Question 9 = 10 marks)

TOTAL FOR PAPER = 60 MARKS







Every effort has been made to contact copyright holders to obtain their permission for the use of copyright material. Pearson Education Ltd. will, if notified, be happy to rectify any errors or omissions and include any such rectifications in future editions.

