



Pearson

Mark Scheme (Results)

January 2017

International GCSE
Physics (4PH0) Paper 2P

Pearson Edexcel Certificate in Physics (KPH0)
Paper 2P

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2017

Publications Code 4PH0_2P_1701_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks																
1 (a) (i)	<p>1 mark for each correct property;;</p> <table border="1" data-bbox="376 342 1158 898"> <thead> <tr> <th>Type of Radiation</th> <th>Nature</th> <th>Relative Charge</th> <th>Ionising Ability</th> </tr> </thead> <tbody> <tr> <td>alpha (α)</td> <td>helium nucleus</td> <td>(+)2</td> <td>high</td> </tr> <tr> <td>beta (β)</td> <td>(high energy) electron</td> <td>-1</td> <td>medium</td> </tr> <tr> <td>gamma (γ)</td> <td>electromagnetic wave</td> <td>0</td> <td>low</td> </tr> </tbody> </table> <p>(ii) alpha / α;</p> <p>(iii) alpha and beta / α and β;</p>	Type of Radiation	Nature	Relative Charge	Ionising Ability	alpha (α)	helium nucleus	(+)2	high	beta (β)	(high energy) electron	-1	medium	gamma (γ)	electromagnetic wave	0	low	<p>reject -2</p> <p>both required but can be in either order</p>	<p>2</p> <p>1</p> <p>1</p>
Type of Radiation	Nature	Relative Charge	Ionising Ability																
alpha (α)	helium nucleus	(+)2	high																
beta (β)	(high energy) electron	-1	medium																
gamma (γ)	electromagnetic wave	0	low																
(b)	<p>top line correct; bottom line correct;</p> <p>e.g.</p> $\begin{array}{ccc} \boxed{14} & & \boxed{14} \\ \text{C} & \longrightarrow & \text{N} \end{array} + \begin{array}{c} \boxed{0} \\ \boxed{-1} \end{array} \beta$	<p>e.g. 14, 0 e.g. 7</p>	2																


Total for question = 6 marks

Question number	Answer	Notes	Marks
2 (a)	(sum of) clockwise moments equals (sum of) anti-clockwise moments; (for a system) to be balanced / in equilibrium;		2
(b) (i)	measure <u>mass</u> of plastic strip (in kg); multiply mass by g;	allow multiply by 10 / 9.8 / 9.81 allow idea of setting balance to zero for 1 mark if no other mark scored	2
(ii)	any 1 of: use a ruler with smaller divisions; use a mm ruler; use a balance that measures to more decimal places;		1
(iii)	moment = force x (perpendicular) distance (from the pivot);	allow correct symbols e.g. moment = F x d condone use of M or m for moment	1
(iv)	calculates one moment correctly; correct use of principle of moments; evaluation; e.g. $0.2 \times 0.3 = 0.06$ $0.06 = 0.1 \times F$ (force =) 0.6 (N)	allow distances in cm throughout 0.4 (N) gets 1 mark max.	3
(v)	any 1 of: idea that calculated force includes weight of beaker / weight of beaker should be subtracted; mass of paperclip / string not considered; centre of mass of ruler may not be at 50 cm;	ignore references to mass/weight of rule allow mass/weight of beaker not considered	1

Total for question = 10 marks

Question number	Answer	Notes	Marks
3 (a)	B – sound waves are transverse;		1
(b) (i)	calculation of time period; substitution into correct frequency equation; evaluation; e.g. $(\text{time period} / T) = 0.02 \text{ (s)}$ $(f =) 1/0.02$ $(f =) 50 \text{ (Hz)}$	allow ecf for incorrect time period allow 0.02 seen anywhere 16.7, 100 (Hz) get 2 marks max.	3
(ii)	line drawn has smaller amplitude than existing line <u>throughout</u> ; line drawn has higher frequency (pitch) <u>throughout</u> ;	ignore vertical position of line	2

Total for question = 6 marks

Question number	Answer	Notes	Marks																		
4 (a) (i)	number of layers (of insulation);		1																		
(ii)	final temperature (of the water) / temperature after 15 minutes / rate at which the water cools down;	allow temperature difference (of the water) ignore unqualified 'temperature'	1																		
(b) (i)	<table border="1" data-bbox="370 562 1066 958"> <thead> <tr> <th data-bbox="370 562 587 734">Number of layers of insulation</th> <th data-bbox="587 562 820 734">Final temperature in °C</th> <th data-bbox="820 562 1066 734">Temperature difference in °C</th> </tr> </thead> <tbody> <tr> <td data-bbox="370 734 587 779">0</td> <td data-bbox="587 734 820 779">43</td> <td data-bbox="820 734 1066 779">42</td> </tr> <tr> <td data-bbox="370 779 587 824">1</td> <td data-bbox="587 779 820 824">47</td> <td data-bbox="820 779 1066 824">38</td> </tr> <tr> <td data-bbox="370 824 587 869">2</td> <td data-bbox="587 824 820 869">50</td> <td data-bbox="820 824 1066 869">35</td> </tr> <tr> <td data-bbox="370 869 587 913">3</td> <td data-bbox="587 869 820 913">50</td> <td data-bbox="820 869 1066 913">35</td> </tr> <tr> <td data-bbox="370 913 587 958">4</td> <td data-bbox="587 913 820 958">50</td> <td data-bbox="820 913 1066 958">35</td> </tr> </tbody> </table>	Number of layers of insulation	Final temperature in °C	Temperature difference in °C	0	43	42	1	47	38	2	50	35	3	50	35	4	50	35		2
Number of layers of insulation	Final temperature in °C	Temperature difference in °C																			
0	43	42																			
1	47	38																			
2	50	35																			
3	50	35																			
4	50	35																			
(ii)	<p>47 in first answer space; 50 in last three answer spaces;</p> <p>suitable scale chosen – longest bar occupies at least half of the grid;</p> <p>axes labelled correctly with quantities and temperature difference unit; all 5 bars correctly plotted;;</p> 	<p>must be seen in all three spaces</p> <p>ignore orientation</p> <p>temperature scale should be linear but need not start at 0</p> <p>reject both plotting marks if a line graph is drawn</p> <p>award 3 marks max. if graph is drawn using final temperature values instead of temperature difference values</p>	4																		

(iii)	any 2 of: MP1. idea of inverse relationship; MP2. idea that each additional layer gives a smaller decrease in temperature difference; MP3. idea of no effect on temperature difference with more than 2 layers;	allow pattern statements negative correlation	2
(iv)	repeat AND {average (mean)/discard anomalies};		1

Total for question = 11 marks

Question number	Answer	Notes	Marks
5 (a)	conversion of hours to seconds; substitution and rearrangement of equation; evaluation; e.g. time = $40 \times 60 \times 60$ (= 144 000 (s)) energy = $50 \times 144\,000$ (energy) = 7 200 000 (J)	no mark for equation as given in paper seen anywhere in working allow 2 000, 120 000 (J) for 2 marks	3
(b)	MP1. energy is wasted / lost (to the surroundings) as thermal energy; MP2. idea that light energy (output) is less than the electrical / input energy;	ignore statements about student being right/wrong allow heat allow RA e.g. 'heat is not useful' e.g. 'not all electrical energy is converted to light'	2
(c)	MP1. two coils of wire; MP2. iron core; MP3. more turns (of wire) on the primary coil than on the secondary coil;	marks can be awarded from diagram if clear allow 'magnetically soft ' core allow input for primary and output for secondary	3
(d) (i)	input power = output power;	allow $V_p I_p = V_s I_s$ rearrangements Use of 1,2 in place of P,S	1
(ii)	substitution into a correct equation; rearrangement; evaluation; e.g. $230 \times I_p = 12 \times 4.2$ ($I_p =$) $12 \times 4.2 / 230$ ($I_p =$) 0.22 (A)	0.21 (A) gets 2 marks only allow 0.2, 0.21913...	3

Total for question = 12 marks

Question number	Answer	Notes	Marks
6 (a)	B – energy;		1
(b) (i)	(resultant force =) 6750 (N);		1
(ii)	(resultant) force = mass x acceleration;	allow in standard symbols and rearrangements e.g. $F = m \times a$	1
(iii)	substitution OR rearrangement; evaluation; unit; e.g. acceleration = $6750/2500$ (acceleration =) 2.7 m/s^2	allow ecf from (b)(i) unit mark is independent allow $m s^{-2}$	3
(c)	any 5 of: MP1. there is a resultant force (to the right); MP2. (so) it accelerates (0 to 50 s); MP3. air resistance (and friction) increase as speed increases; MP4. so acceleration decreases; MP5. eventually air resistance (+ friction) = driving force; MP6. (hence) resultant force is zero (after 50 s); MP7. (hence) car travels at a constant speed (after 50 s);	allow idea that driving force is greater than air resistance and friction the speed/velocity increases forces are equal / balanced no acceleration / terminal velocity	5

Total for question = 11 marks

Question number	Answer	Notes	Marks
7 (a)	D – newtons per square metre (N/m ²);		1
(b)	any 3 of: MP1. air is heated / temperature of air increases; MP2. (air) molecules move faster / gain kinetic energy; MP3. more (frequent) collisions between molecules and walls ; MP4. molecules collide with walls with more force;	allow particles for molecules throughout allow pressure is proportional to (kelvin) temperature allow molecules collide harder with walls allow rate of change of momentum for force	3

Total for question = 4 marks

